



*Stiles*  
APR 25 1888

# CHECK-LIST

OF

# NORTH AMERICAN BIRDS

7-8  
QL  
677  
A52  
1895  
Bird Re

PREPARED BY A COMMITTEE

OF THE

American Ornithologists' Union


SECOND AND REVISED EDITION

Private property of

Dr. C. W. Stiles

Zoologist, B. A. I.

Zoölogical Nomenclature is a means, not an end, of Zoölogical Science



NEW YORK

AMERICAN ORNITHOLOGISTS' UNION

1895

*Copyright, 1895,*  
BY AMERICAN ORNITHOLOGISTS' UNION.

Cambridge, Mass. :  
PRESS OF EDWARD W. WHEELER.

## PREFACE TO THE FIRST EDITION.

---

AT the first Congress of the American Ornithologists' Union, held in New York, September 26-29, 1883, the following resolution was adopted:—

*“Resolved, That the Chairman appoint a Committee of five, including himself, to whom shall be referred the question of a Revision of the Classification and Nomenclature of the Birds of North America.”*

In pursuance of this resolution the following Committee was appointed: Messrs. Coues, Allen, Ridgway, Brewster, and Henshaw.

The Committee, having held numerous sessions in Washington and New York, presented its Report at the second Congress of the Union, held in New York, Sept. 30 to Oct. 2, 1884, when the following resolution was adopted:—

*“Resolved, That the Report of the Committee on the Revision of the Nomenclature and Classification of North American Birds be accepted and adopted, and that it be recommitted to the Committee, with instructions to complete and submit it to the Council as soon as practicable; and that the Council be empowered and instructed to accept and adopt the Report as finally rendered, with such modifications as they may deem necessary, and to publish the same, copyrighted, in part or in whole, and in one or more forms, in the name and under the auspices of the American Ornithologists' Union.”*

The Committee, having continued its sessions, presented its final report to the Council at a meeting held in Washington



on the 21st of April, 1885, when the Report of the Committee was accepted and adopted, and was referred again to the Committee for publication, the Committee to exercise such editorial revision as might seem necessary.

Pursuant to the foregoing resolutions of the Union and Council, the Committee now offers to the public, in the name and on behalf of the Union, the result of its labors, consisting of a List of North American Birds, preceded by the Code of Rules adopted by the Committee for its guidance in the preparation of the List.

The Committee ventures to hope that the new Code will find favor, not only with ornithologists, but among zoölogists generally.

ELLIOTT COUES.

J. A. ALLEN.

ROBERT RIDGWAY.

WILLIAM BREWSTER.

H. W. HENSHAW.

## PREFACE TO THE SECOND EDITION.

---

AT the Eleventh Congress of the American Ornithologists' Union, held in Cambridge, Mass., November 20-23, 1894, it was voted to publish, as early as practicable, a new edition of the Union's Check-List of North American Birds, to include the numerous additions and nomenclatural changes made in the several Supplements<sup>1</sup> to the Check-List since the publication of the original edition, together with a revision of the 'habitats' of the species and subspecies, but omitting the Code of Nomenclature.<sup>2</sup> The original Com-

<sup>1</sup> Supplement | to the | Code of Nomenclature and Check-List | of | North American Birds | adopted by the American Ornithologists' Union | Prepared by | a Committee of the Union | — | New York | American Ornithologists' Union | 1889. — 8vo, pp. 23.

Second Supplement to the American Ornithologists' Union Check-List of North American Birds. *The Auk*, VII, Jan. 1890, pp. 60-66. Also separate.

Third Supplement to the American Ornithologists' Union Check-List of North American Birds. *The Auk*, VIII, Jan. 1891, pp. 83-90. Also separate.

Fourth Supplement to the American Ornithologists' Union Check-List of North American Birds. *The Auk*, IX, Jan. 1892, pp. 105-108. Also separate.

Fifth Supplement to the American Ornithologists' Union Check-List of North American Birds. *The Auk*, X, Jan. 1893, pp. 59-63. Also separate.

Sixth Supplement to the American Ornithologists' Union Check-List of North American Birds. *The Auk*, XI, Jan. 1894, pp. 46-51. Also separate.

Seventh Supplement to the American Ornithologists' Union Check-List of North American Birds. *The Auk*, XII, April, 1894, pp. 163-169. Also separate.

Check-List | of | North American Birds | according to the Canons of Nomenclature | of the | American Ornithologists' Union | — | Abridged Edition | Revised | — | Published by the American Ornithologists' Union | 1889. — 8vo, pp. 71. Includes the additions to the original Check-List made in the first Supplement.

<sup>2</sup> Republished separately in 1892 as a pamphlet of 72 pages with the following title: The | Code of Nomenclature | adopted by the | American Ornithologists' Union | — | Zoölogical Nomenclature is a means, not an end, of Zoölogical Science | — | New York | American Ornithologists' Union | 1892.



mittee on Classification and Nomenclature of North American Birds was reappointed<sup>1</sup> to take charge of the work. The Committee held sessions in Washington, D. C., January 15-19, 1894, and February 12, 13, 1895, to outline the work and to rule on the questions involved in the publication of the revised List. The revision of the matter relating to the geographic distribution of the species and subspecies was undertaken by the Committee as a whole, each member in turn taking it in hand, while the incorporation of typographic and other rectifications made during the sessions of the Committee,<sup>2</sup> and the final preparation of the manuscript for the printer, was referred to a subcommittee consisting of the Editor of 'The Auk,' to whom was also assigned the general editorial supervision of the work.

The following extracts from the Introduction to the Code of Nomenclature (pp. 14, 15) will serve to explain the scope and plan of the Check-List, including the method of incorporating additions.

"1. That the term 'North American,' as applied to the proposed List of Birds, be held to include the continent of North America north of the present United States and Mexican boundary, and Greenland; and the peninsula of Lower California, with the islands naturally belonging thereto.

"2. That species be numbered consecutively, and that subspecies be enumerated by affixing the letters *a*, *b*, *c*, etc., to the number borne by their respective species; provided, that any subspecies of a species not included in the North American Fauna shall be separately numbered as if a species.

"3. That stragglers or accidental visitors, not regarded as components of the North American Fauna, be distinguished by having their respective numbers in brackets.

<sup>1</sup> With the exception of Mr. H. W. Henshaw, who was unable to serve, and Dr. C. Hart Merriam was appointed in his stead.

<sup>2</sup> The Committee desires to here acknowledge valuable assistance received, especially in the preparation of the geographic portions of the list, from Major Charles Bendire, Mr. Frank M. Chapman, Dr. Walter Faxon, Dr. A. K. Fisher, Mr. Gerrit S. Miller, Jr., and Dr. T. S. Palmer. The Committee is further indebted to Dr. Palmer for numerous corrections in the citations of original references.



"4. That any subsequent additions to the list be interpolated in systematic order, and bear the number of the species immediately preceding, with the addition of a figure (1, 2, etc., as the case may require), separated from the original number by a period or decimal point, thus giving the interpolated number a decimal form (*c. g.*, 243.1, etc.), in order that the original numbers may be permanent.

\* \* \* \* \*

"6. That Giraud's at present unconfirmed species of Texan birds be included in the List on Giraud's authority.

"7. That species and subspecies the zoölogical status of which cannot be satisfactorily determined, like, *e. g.*, *Regulus curvieri* and *Spiza townsendi* of Audubon, be referred to a hypothetical list, in each case with a brief statement of the reasons for such allocation.

"8. That a list of the fossil species of North American birds be added as an Appendix to the List proper.

"9. That the names of subgeneric and supergeneric groups of North American birds be included in the List in systematic order, to the end that the List may represent a classification as well as a nomenclature of the birds.

"10. That references be given to the original description of the species, and to the publication where the name as adopted in the List was first used; that the number borne by each species and subspecies in the Lists of Baird, 1858, of Coues, 1873, of Ridgway, 1880, and of Coues, 1882, be bracketed in chronological order after the synonymatic references.

"11. That a summary statement of the habitat of each species and subspecies, with special reference to its North American range, be included in the List.

"12. That the name of each bird shall consist of its generic without its subgeneric name, and of its specific with its subspecific name, if it have one, without the intervention of any other term.

\* \* \* \* \*

"14. That every technical name be followed by a vernacular name, selected with due regard to its desirability.

"15. That the name of each species and subspecies be followed by the name of the original describer of the same, to be enclosed in parentheses when it is not also the authority for the name adopted.

“16. That all specific and subspecific names shall begin with a lower-case letter.

“17. That the sequence in classification followed in previous Lists be reversed, the List to begin with the lowest or most generalized type, and end with the highest or most specialized.”

ELLIOTT COUES.

J. A. ALLEN.

WILLIAM BREWSTER.

C. HART MERRIAM.

ROBERT RIDGWAY.

# TABLE OF CONTENTS.

---

	PAGE
PREFACE TO FIRST EDITION . . . . .	iii
PREFACE TO SECOND EDITION . . . . .	v
CHECK-LIST . . . . .	I
✓ I. PYGOPODES . . . . .	I
<i>a.</i> Podicipedes . . . . .	I
1. Podicipidæ . . . . .	I
<i>b.</i> Cepphi . . . . .	3
2. Urinatoridæ . . . . .	3
3. Alcidæ . . . . .	5
✓ II. LONGIPENNES . . . . .	13
4. Stercorariidæ . . . . .	13
5. Laridæ . . . . .	15
6. Rynchopidæ . . . . .	27
III. TUBINARES . . . . .	28
7. Diomedeidæ . . . . .	28
8. Procellariidæ . . . . .	29
IV. STEGANOPODES . . . . .	39
9. Phaëthontidæ . . . . .	39
10. Sulidæ . . . . .	39
11. Anhingidæ . . . . .	41
12. Phalacrocoracidæ . . . . .	42
13. Pelecanidæ . . . . .	45
14. Fregatidæ . . . . .	46
✓ V. ANSERES . . . . .	47
15. Anatidæ . . . . .	47
✓ VI. ODONTOGLOSSÆ . . . . .	66
16. Phœnicopteridæ . . . . .	66
VII. HERODIONES . . . . .	66
<i>c.</i> Ibides . . . . .	66
17. Plataleidæ . . . . .	66
18. Ibididæ . . . . .	67



	PAGE
<i>d.</i> Ciconiæ . . . . .	68
19. Ciconiidae . . . . .	68
<i>e.</i> Herodii . . . . .	69
20. Ardeidae . . . . .	69
✓ VIII. PALUDICOLÆ . . . . .	75
<i>f.</i> Grues . . . . .	75
21. Gruidæ . . . . .	75
<i>g.</i> Ralli . . . . .	76
22. Aramidæ . . . . .	76
23. Rallidæ . . . . .	76
✓ IX. LIMICOLÆ . . . . .	82
24. Phalaropodidæ . . . . .	82
25. Recurvirostridæ . . . . .	83
26. Scolopacidæ . . . . .	84
27. Charadriidæ . . . . .	98
28. Aphrizidæ . . . . .	103
29. Hæmatopodidæ . . . . .	104
30. Jacanidæ . . . . .	105
✓ X. GALLINÆ . . . . .	106
<i>h.</i> Phasiani . . . . .	107
31. Tetraonidæ . . . . .	107
32. Phasianidæ . . . . .	117
<i>i.</i> Penelopes . . . . .	119
33. Cracidæ . . . . .	119
✓ XI. COLUMBÆ . . . . .	119
34. Columbidae . . . . .	119
XII. RAPTORES . . . . .	124
<i>j.</i> Sarcorhamphi . . . . .	124
35. Cathartidæ . . . . .	124
<i>k.</i> Falcones . . . . .	126
36. Falconidæ . . . . .	126
<i>l.</i> Striges . . . . .	142
37. Strigidæ . . . . .	142
38. Bubonidæ . . . . .	142
XIII. PSITTACI . . . . .	152
39. Psittacidæ . . . . .	152
XIV. COCCYGES . . . . .	153
<i>m.</i> Cuculi . . . . .	153
40. Cuculidæ . . . . .	153
<i>n.</i> Trogones . . . . .	156
41. Trogonidæ . . . . .	156
<i>o.</i> Alcyones . . . . .	156
42. Alcedinidæ . . . . .	156

# TABLE OF CONTENTS.

xi

	PAGE
✓ XV. PICI . . . . .	157
43. Picidæ . . . . .	157
✓ XVI. MACROCHIRES . . . . .	168
<i>p.</i> Caprimulgi . . . . .	168
44. Caprimulgidæ . . . . .	168
<i>q.</i> Cypseli . . . . .	171
45. Micropodidæ . . . . .	171
<i>r.</i> Trochili . . . . .	173
46. Trochilidæ . . . . .	173
✓ XVII. PASSERES . . . . .	179
<i>s.</i> Clamatores . . . . .	179
47. Cotingidæ . . . . .	179
48. Tyrannidæ . . . . .	179
<i>t.</i> Oscines . . . . .	191
49. Alaudidæ . . . . .	191
50. Corvidæ . . . . .	194
51. Sturnidæ . . . . .	202
52. Icteridæ . . . . .	202
53. Fringillidæ . . . . .	211
54. Tanagridæ . . . . .	254
55. Hirundinidæ . . . . .	256
56. Ampelidæ . . . . .	260
57. Laniidæ . . . . .	261
58. Vireonidæ . . . . .	262
59. Cœrebidæ . . . . .	267
60. Mniotiltidæ . . . . .	268
61. Motacillidæ . . . . .	288
62. Cinclidæ . . . . .	291
63. Troglodytidæ . . . . .	291
64. Certhiidæ . . . . .	303
65. Paridæ . . . . .	304
66. Sylviidæ . . . . .	313
67. Turdidæ . . . . .	316
HYPOTHETICAL LIST . . . . .	325
THE FOSSIL BIRDS OF NORTH AMERICA . . . . .	335
INDEX. . . . .	347





# CHECK-LIST.

---

## ORDER PYGOPODES. DIVING BIRDS.

### SUBORDER PODICIPEDES. GREBES.

#### FAMILY **PODICIPIDÆ**. GREBES.

#### GENUS **ÆCHMOPHORUS** COUES.

*Æchmophorus* COUES, Pr. Ac. Nat. Sci. Phila. April, 1862, 229.  
Type, *Podiceps occidentalis* LAWR.

1. **Æchmophorus occidentalis** (LAWR.).

Western Grebe.

*Podiceps occidentalis* LAWR. in BAIRD'S B. N. Am. 1858, 894.

*Æchmophorus occidentalis* COUES, Pr. Ac. Nat. Sci. Phila. 1862,  
229.

[B 704, C 608, R 729, C 845.]

GEOGRAPHIC DISTRIBUTION.—Western North America, eastward to Manitoba, south to central Mexico.

#### GENUS **COLYMBUS** LINNÆUS.

*Colymbus* LINN. S. N. ed. 10, I. 1758, 135. Type, by elimination,  
*Colymbus cristatus* LINN.

#### SUBGENUS **COLYMBUS**.

2. **Colymbus holbœllii** (REINH.).

Holbœll's Grebe.

*Podiceps holbœllii* REINH. Vid. Med. 1853, 76.

*Colymbus holbœllii* RIDGW. Water B. N. Am. II. 1884, 428.

[B 702, C 610, R 731, C 847.]

GEOG. DIST.—North America at large, including Greenland. Also eastern Siberia, and southward to Japan. Breeds in high latitudes, migrating south in winter.

SUBGENUS **DYTES** KAUP.

*Dytes* KAUP, Sk. Ent. Eur. Thierw. 1829, 41. Type, *Colymbus auritus* LINN.

3. **Colymbus auritus** LINN.

**Horned Grebe.**

*Colymbus auritus* LINN. S. N. ed. 10, I. 1758, 135.

[B 706, C 611, R 732, C 848.]

GEOG. DIST.—Northern hemisphere. Breeds from the northern United States northward.

4. **Colymbus nigricollis californicus** (HEERM.).

**American Eared Grebe.**

*Podiceps californicus* HEERM. Pr. Ac. Nat. Sci. Phila. 1854, 179.

*Colymbus nigricollis californicus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 356.

[B 707, C 612, R 733a, C 850.]

GEOG. DIST.—Northern and western North America, from the Mississippi Valley westward, southward to Guatemala.

SUBGENUS **PODICEPS** LATHAM.

*Podiceps* LATH. Ind. Orn. II. 1790, 780. Type, by elimination, *Colymbus fluviatilis* TUNST.

5. **Colymbus dominicus** LINN.

**St. Domingo Grebe.**

*Colymbus dominicus* LINN. S. N. ed. 12, I. 1766, 223.

[B 708a, C 613, R 734, C 851.]

GEOG. DIST.—Southern Texas and Lower California southward through tropical America to Paraguay, including the West Indies.

GENUS **PODILYMBUS** LESSON.

*Podilymbus* LESS. Traité, I. 1831, 595. Type, *Colymbus podiceps* LINN.

6. **Podilymbus podiceps** (LINN.).  
Pied-billed Grebe.

*Colymbus podiceps* LINN. S. N. ed. 10, I. 1758, 136.

*Podilymbus podiceps* LAWR. in BAIRD'S B. N. Am. 1858, 898.

[B 709, C 614, R 735, C 852.]

GEOG. DIST.—British Provinces southward to Brazil, Argentine Republic, and Chili, including the West Indies and Bermuda, breeding nearly throughout its range.

## SUBORDER CEPPHI. LOONS AND AUKS.

FAMILY **URINATORIDÆ**. LOONS.GENUS **URINATOR** CUVIER.

*Urinator* CUV. Anat. Comp. I. 1800, tabl. ii. Type, *Colymbus imber* GUNN.

7. **Urinator imber** (GUNN.).  
Loon.

*Colymbus imber* GUNNERUS, Trondh. Selsk. Skr. I. 1761, pl. iii.

*Urinator imber* STEJN. Orn. Expl. Kamchat. 1885, 313.

[B 698, C 605, R 736, C 840.]

GEOG. DIST.—Northern part of northern hemisphere. In North America breeds from the northern tier of States northward; ranges in winter south to the Gulf of Mexico and Lower California.

8. **Urinator adamsii** (GRAY).

Yellow-billed Loon.

*Colymbus adamsii* GRAY, P. Z. S. 1859, 167.*Urinator adamsii* STEJN. Pr. U. S. Nat. Mus. V. 1882, 43.

[B—, C 605a, R 737, C 841.]

GEOG. DIST.—Arctic America, west of Hudson Bay, and north-eastern Asia. Casual in northern Europe.

9. **Urinator arcticus** (LINN.).

Black-throated Loon.

*Colymbus arcticus* LINN. S. N. ed. 10, I. 1758, 135.*Urinator arcticus* STEJN. Pr. U. S. Nat. Mus. V. 1882, 43.

[B 699, C 606, R 738, C 842.]

GEOG. DIST.—Northern part of the northern hemisphere. In North America of casual occurrence in autumn and winter in the northern United States east of Rocky Mountains.

10. **Urinator pacificus** (LAWR.).

Pacific Loon.

*Colymbus pacificus* LAWR. in BAIRD'S B. N. Am. 1858, 889.*Urinator pacificus* STEJN. Pr. U. S. Nat. Mus. V. 1882, 43.

[B 700, C 606a, R 739, C 843.]

GEOG. DIST.—Pacific coast of North America, south in winter to Cape St. Lucas and Guadalupe Island.

11. **Urinator lumme** (GUNN.).

Red-throated Loon.

*Colymbus lumme* GUNN. Trond. Selsk. Skr. I. 1761, pl. ii. fig. 2.*Urinator lumme* STEJN. Pr. U. S. Nat. Mus. V. 1882, 43.

[B 701, C 607, R 740, C 844.]

GEOG. DIST.—Northern part of northern hemisphere, migrating southward in winter nearly across the United States.



FAMILY **ALCIDÆ**. AUKS, MURRES, AND PUFFINS.SUBFAMILY **FRATERCULINÆ**. PUFFINS.GENUS **LUNDA** PALLAS.*Lunda* PALL. Zoog. Rosso-As. II. 1826, 363. Type, *Alca cirrhata* PALL.12. **Lunda cirrhata** PALL.

Tufted Puffin.

*Alca cirrhata* PALL. Spic. Zool. V. 1769, 7, pl. i., pl. ii. figs. 1-3.*Lunda cirrhata* PALL. Zoog. Rosso-As. II. 1826, 363, pl. 82.

[B 712, 716, C 619, R 745, C 856.]

GEOG. DIST. — Coasts and islands of the North Pacific from the Santa Barbara Islands, California, to Alaska, and from Japan to Bering Strait. Accidental on the coast of Maine.

GENUS **FRATERCULA** BRISSON.*Fratercula* BRISS. Orn. VI. 1760, 81. Type, *Alca arctica* LINN.13. **Fratercula arctica** (LINN.).

Puffin.

*Alca arctica* LINN. S. N. ed. 10, I. 1758, 130.*Fratercula arctica* SCHÄFFER, Mus. Orn. 1789, 61.

[B 715, C 618, R 743, C 854.]

GEOG. DIST. — Coasts and islands of the North Atlantic, breeding on the North American coast from the Bay of Fundy northward. South in winter to Long Island, and casually further.

13 a. **Fratercula arctica glacialis** (TEMME.).

Large-billed Puffin.

*Fratercula glacialis* STEPHENS, Gen. Zool. Aves, XIII. pt. 1, 1826, 40.*Fratercula arctica*  $\beta$  *glacialis* BLASIUS, List B. Europ. 1862, 24.

[B 714, C 618a, R 743a, C 855.]

GEOG. DIST.—Coasts and islands of the Arctic Ocean, from Spitzbergen to northern and western Greenland.

14. **Fratercula corniculata** (NAUM.).

Horned Puffin.

*Mormon corniculata* NAUM. Isis, 1821, 782, pl. vii. figs. 3, 4.

*Fratercula corniculata* BRANDT, Bull. Ac. St. Pétersb. II. 1837, 348.

[B 713, C 617, R 744, C 853.]

GEOG. DIST.—Coasts and islands of the North Pacific, from the Kuril Islands to British Columbia.

SUBFAMILY **PHALERINÆ**. AUKLETS, MURRELETS, GUILLEMOTS.

GENUS **CERORHINCA** BONAPARTE.

*Cerorhinca* BONAP. Ann. Lyc. N. Y. 1828, 427. Type, *C. occidentalis* BP. = *Alca monocerata* PALL.

15. **Cerorhinca monocerata** (PALL.).

Rhinoceros Auklet.

*Alca monocerata* PALL. Zoog. Rosso-As. II. 1826, 362.

*Cerorhina monocerata* CASS. in BAIRD'S B. N. Am. 1858, 905.

[B 717, 718, C 620, R 746, C 857.]

GEOG. DIST.—Coasts and islands of the North Pacific, breeding southward (formerly) to the Farallones; in winter southward to Lower California and Japan.

GENUS **PTYCHORAMPHUS** BRANDT.

*Ptychoramphus* BRANDT, Bull. Ac. St. Pétersb. II. 1837, 347.

Type, *Uria aleutica* PALL.

16. **Ptychoramphus aleuticus** (PALL.).

Cassin's Auklet.

*Uria aleutica* PALL. Zoog. Rosso-As. II. 1826, 370.*Ptychoramphus aleuticus* BRANDT, Bull. Ac. St. Pétersb. II. 1837, 347.

[B 724, C 625, R. 751, C 862.]

GEOG. DIST.—Pacific coast of North America, from the Aleutian Islands to Lower California, breeding southward to San Geronimo Island (Lat. 30°).

GENUS **CYCLORRHYNCHUS** KAUP.

*Cyclorhynchus* KAUP, Sk. Ent. Eur. Thierw. 1829, 155. Type, *Alca psittacula* PALL.

17. **Cyclorhynchus psittaculus** (PALL.).

Paroquet Auklet.

*Alca psittacula* PALL. Spic. Zool. V. 1760, 13, pl. ii., pl. v. figs. 4-6.*Cyclorhynchus psittaculus* STEJN. Pr. U. S. Nat. Mus. VII. Aug. 5, 1884, 216.

[B 725, C 621, R 747, C 858.]

GEOG. DIST.—Coasts and islands of the North Pacific, from Sitka and the Kuril Islands northward.

GENUS **SIMORHYNCHUS** MERREM.SUBGENUS **SIMORHYNCHUS**.

*Simorhynchus* MERREM, in ERSCH & GRUBER'S Encycl. I sect. II. 1819, 405. Type, *Alca cristatella* PALL.

18. **Simorhynchus cristatellus** (PALL.).

Crested Auklet.

*Alca cristatella* PALL. Spic. Zool. V. 1769, 20, pl. iii., pl. v. figs. 7-9.*Simorhynchus cristatellus* BONAP. Compt. Rend. XLII. 1856, 774.

[B 719, 720, C 622, R 748, C 859.]

GEOG. DIST.—Coasts and islands of the North Pacific, from Kadiak and Japan northward.

SUBGENUS **PHALERIS** TEMMINCK.

*Phaleris* TEMM. Man. Orn. 1820, p. cxii. Type, by elimination,  
*Alca pygmæa* GMEL.

19. **Simorhynchus pygmæus** (GMEL.).

Whiskered Auklet.

*Alca pygmæa* GMEL. S. N. I. ii. 1788, 555.

*Simorhynchus pygmæus* BRANDT, Mém. Biol. VII. 1869, 222.

[B 721, C 623, R 749, C 860.]

GEOG. DIST.—Coasts and islands of the North Pacific, from Unalaska through the Aleutian chain to Kamchatka.

SUBGENUS **CICERONIA** REICHENBACH.

*Ciceronia* REICH. Syst. Av. 1852, p. iii. Type, *Phaleris microceros* BRANDT = *Uria pusilla* PALL.

20. **Simorhynchus pusillus** (PALL.).

Least Auklet.

*Uria pusilla* PALL. Zoog. Rosso-As. II. 1826, 373, pl. 70.

*Simorhynchus pusillus* COUES, Pr. Ac. Nat. Sci. Phila. 1862, 324.

[B 722, 723, C 624, R 750, C 861.]

GEOG. DIST.—Coasts and islands of the North Pacific, from Sitka and Japan northward to Bering Strait.

GENUS **SYNTHLIBORAMPHUS** BRANDT.

*Synthliboramphus* BRANDT, Bull. Ac. St. Pétersb. II. 1837, 347.  
Type, *Alca antiqua* GMEL.

21. **Synthliboramphus antiquus** (GMEL.).

Ancient Murrelet.

*Alca antiqua* GMEL. S. N. I. ii. 1788, 554.

*Synthliboramphus antiquus* BRANDT, Bull. Ac. St. Pétersb. II.  
1837, 347.



[B 734, 736, C 627, R 753, 759, C 864, 870.]

GEOG. DIST.—Coasts and islands of the North Pacific, from southern Vancouver Island and Japan northward. Accidental in Wisconsin.

GENUS **BRACHYRAMPHUS** BRANDT.

*Brachyramphus* BRANDT, Bull. Ac. St. Pétersb. II. 1837, 346.

Type, *Colymbus marmoratus* GMEL.

23. **Brachyramphus marmoratus** (GMEL.).

Marbled Murrelet.

*Colymbus marmoratus* GMEL. S. N. I. ii. 1788, 583.

*Brachyramphus marmoratus* BRANDT, Bull. Ac. St. Pétersb. II. 1837, 346.

[B 732, 733, C 629, R 755, C 866.]

GEOG. DIST.—Coasts and islands of the North Pacific; on the American coast from San Diego northward, and breeding as far south as Vancouver Island.

24. **Brachyramphus kittlitzii** BRANDT.

Kittlitz's Murrelet.

*Brachyramphus kittlitzii* BRANDT, Bull. Ac. St. Pétersb. II. 1837, 346.

[B 735, C 630, R 756, C 867.]

GEOG. DIST.—Northern Japan, Kamchatka and Aleutian Islands, east to Unalaska.

25. **Brachyramphus hypoleucus** XANTUS.

Xantus's Murrelet.

*Brachyrhamphus hypoleucus* XANTUS, Pr. Ac. Nat. Sci. Phila., Nov. 1859, 299.

[B —, C —, R 757, C 868.]

GEOG. DIST.—Coasts of southern and Lower California, from Santa Barbara Island to Cape St. Lucas.

26. **Brachyramphus craveri** (SALVAD.).

Craveri's Murrelet.

*Uria craveri* SALVAD. Atti Soc. It. Sc. Nat. VIII. 1866, Estr.  
p. 17.

*Brachyrhamphus craverii* COUES, Pr. Ac. Nat. Sci. Phila., 1868,  
66.

[B—, C—, R 758, C 869.]

GEOG. DIST.—Both coasts of Lower California, from Cape St. Lucas northward to Espiritu Santo Island in the Gulf of California, and to Natividad Island (lat. 28°) on the Pacific side.

GENUS **CEPPHUS** PALLAS.

*Cepphus* PALL. Spic. Zool. V. 1769, 33. Type, *C. lacteolus* PALL.  
= *C. grylle*, albino.

27. **Cepphus grylle** (LINN.).

Black Guillemot.

*Alca grylle* LINN. S. N. ed. 10, I. 1758, 130.

*Cepphus grylle* BREHM, Handb. Vög. Deutschl. 1831, 987.

[B 726, *part*, C 631, *part*, R 760, *part*, C 871, *part*.]

GEOG. DIST.—Coasts of northern Europe, south to Denmark and the British Islands. Newfoundland and coast of Maine, south in winter to Philadelphia.

28. **Cepphus mandtii** (LICHT.).

Mandt's Guillemot.

*Uria mandtii* LICHT. in MANDT'S Obs. Itin. Dissert. 1822, 30.

*Cepphus mandtii* BP. Cat. Parzud. 1856, 12.

[B 726, *part*, C 631, *part*, R 760, *part*, C 871, *part*.]

GEOG. DIST.—Arctic regions of both continents; south on the Atlantic coast of North America, in winter, to Massachusetts, breeding south to Hudson Bay and Labrador; Alaskan coast, south, in winter, to Norton Sound.

29. **Cepphus columba** PALL.

Pigeon Guillemot.

*Cepphus columba* PALL. Zoog. Rosso-As. II. 1826, 348.

[B 727, C 632, R 761, C 872.]

GEOG. DIST.—Coasts and islands of the North Pacific, southward from Bering Strait to northern Japan and the Santa Barbara Islands, California.

SUBFAMILY **ALCINÆ**. AUKS AND MURRES.GENUS **URIA** BRISSON.

*Uria* BRISS. Orn. VI. 1760, 70. Type, by elimination, *Colymbus troile* LINN.

30. **Uria troile** (LINN.).

Murre.

*Colymbus troile* LINN. Faun. Succ. ed. 1761, 52; S. N. ed. 12, I. 1766, 220.

*Uria troile* LATH. Ind. Orn. II. 1790, 796.

[B 729, 730, C 634, R 763, C 874.]

GEOG. DIST.—Coasts and islands of the North Atlantic, southward on the coast of North America, in winter, to southern New England; breeding from the Magdalen Islands northward.

30 a. **Uria troile californica** (BRYANT).

California Murre.

*Catarractes californicus* BRYANT, Pr. Bost. Soc. VIII. 1861, 11, figs. 3, 5.

*Uria troile californica* RIDGW. Water B. N. Am. II. 1884, 483.

[B—, C—, R 763a, C 875.]

GEOG. DIST.—Coasts and islands of the North Pacific, breeding from California north to the Pribilof Islands.

31. **Uria lomvia** (LINN.).

Brünnich's Murre.

*Alca lomvia* LINN. S. N. ed. 10, I. 1758, 130.*Uria lomvia* BRYANT, Proc. Bost. Soc. N. H. VIII. May, 1861, 75.

[B 731, C 635, R 764a, C 876.]

GEOG. DIST.—Coasts and islands of the North Atlantic and eastern Arctic Oceans; south to the lakes of northern New York and the coast of New Jersey. Breeding from the Gulf of St. Lawrence northward.

31 a. **Uria lomvia arra** (PALL.).

Pallas's Murre.

*Cephus arra* PALL. Zoog. Rosso-As. II. 1826, 347.*Uria lomvia arra* RIDGW. Water B. N. Am. II. Sept. 1884, 485.

[B —, C —, R 764, C —.]

GEOG. DIST.—Coasts and islands of the North Pacific and western Arctic Oceans, south to Kadiak and Kamchatka.

GENUS **ALCA** LINNÆUS.*Alca* LINN. S. N. ed. 10, I. 1758, 130. Type, by elimination,*Alca torda* LINN.32. **Alca torda** LINN.

Razor-billed Auk.

*Alca torda* LINN. S. N. ed. 10, I. 1758, 130.

[B 711, C 616, R 742, C 877.]

GEOG. DIST.—Coasts and islands of the North Atlantic, south in winter on the North American coast casually to North Carolina, breeding from eastern Maine northward.

GENUS **PLAUTUS** BRÜNNICH.*Plautus* BRÜNN. Zool. Fund. 1772, 78. Type, *Alca impennis* LINN.



33. **Plautus impennis** (LINN.).

Great Auk.

*Alca impennis* LINN. S. N. ed. 10, I. 1758, 130.*Plautus impennis* STEENSTR. Vid. Med. Nat. For. Kjob. 1855,  
114.

[B 710, C 615, R 741, C 878.]

GEOG. DIST.—Formerly the coasts and islands of the North Atlantic, from Massachusetts and Ireland northward nearly to the Arctic Circle. Now extinct.

SUBFAMILY **ALLINÆ**. DOVEKIES.GENUS **ALLE** LINK.*Alle* LINK, Besch. Nat. Samml. Univ. Rostock, I. 1806, 17.Type, *Alca alle* LINN.34. **Alle alle** (LINN.).

Dovekie.

*Alca alle* LINN. S. N. ed. 10, I. 1758, 131.*Alle alle* STEJNEGER, Stand. Nat. Hist. IV, 1885, 69.

[B 738, C 626, R 752, C 863.]

GEOG. DIST.—Coasts and islands of the North Atlantic and eastern Arctic Oceans; in North America south in winter to New Jersey; breeds in high northern latitudes. Accidental in Michigan.

ORDER LONGIPENNES. LONG-WINGED  
SWIMMERS.FAMILY **STERCORARIIDÆ**. SKUAS AND JAEGERES.GENUS **MEGALESTRIS** BONAPARTE.*Megalestris* BONAP. Cat. Parzudaki, 1856, 11. Type, *Catharacta skua* BRÜNN.

35. **Megalestris skua** (BRÜNN.).

Skua.

*Catharacta skua* BRÜNN. Orn. Bor. 1764, 33.*Megalestris skua* RIDGW. Pr. U. S. Nat. Mus. III. Sept. 4, 1880, 208.

[B 652, C 539, R 696, C 764.]

GEOG. DIST.—Coasts and islands of the North Atlantic, chiefly northward. South to Spain and North Carolina. Apparently rare on the coast of North America.

GENUS **STERCORARIUS** BRISSON.

*Stercorarius* BRISS. Orn. V. 1760, 149. Type, *Larus parasiticus* LINN.

36. **Stercorarius pomarinus** (TEMM.).

Pomarine Jaeger.

*Larus pomarinus* TEMM. Man. d'Orn. 1815, 514.• *Stercorarius pomarinus* VIEILL. Nouv. Dict. XXXII. 1819, 158.

[B 653, C 540, R 697, C 765.]

GEOG. DIST.—Seas and inland waters of northern portions of the northern hemisphere, south in winter to Africa, Australia, and probably South America.

37. **Stercorarius parasiticus** (LINN.).

Parasitic Jaeger.

*Larus parasiticus* LINN. S. N. ed. 10, I. 1758, 136.*Stercorarius parasiticus* SCHÄFF. Mus. Orn. 1789, 62, pl. 37.

[B 654, C 541, R 698, C 766.]

GEOG. DIST.—Northern part of northern hemisphere, southward in winter to South Africa and South America. Breeds in high northern districts, and winters from New York and California southward to Brazil.

38. **Stercorarius longicaudus** VIEILL.

Long-tailed Jaeger.

*Stercorarius longicaudus* VIEILL. Nouv. Dict. XXXII. 1819, 157.

[B 655, C 542, R 699, C 767.]

GEOG. DIST.—Northern part of northern hemisphere, breeding in high northern districts ; south in winter to the Gulf of Mexico.

FAMILY **LARIDÆ**. GULLS AND TERNS.SUBFAMILY **LARINÆ**. GULLS.GENUS **GAVIA** BOIE.

*Gavia* BOIE, Isis, 1822, 563. Type, *Larus eburneus* PHIPPS =  
*Larus albus* GUNN.

39. **Gavia alba** (GUNN.).

Ivory Gull.

*Larus albus* GUNN. in LEEM'S Beskr. Finn. Lapp. 1767, 285.*Gavia alba* STEJN. Pr. U. S. Nat. Mus. V. 1882, 39.

[B 676, 677, C 550, R 657, C 785.]

GEOG. DIST.—Arctic Seas, south in winter on the Atlantic coast of North America to Labrador and Newfoundland, casually to New Brunswick, and on the Pacific side to Bering Sea.

GENUS **RISSA** STEPHENS.

*Rissa* STEPHENS, Gen. Zool. XIII. 1825, 180. Type, *Larus tridactylus* LINN.

40. **Rissa tridactyla** (LINN.).

Kittiwake.

*Larus tridactylus* LINN. S. N. ed. 10, I. 1758, 136.*Rissa tridactyla* BONAP. Comp. List, 1838, 62.

[B 672, C 552, R 658, C 782.]

GEOG. DIST.—Arctic regions, south in eastern North America in winter to the Great Lakes and the Middle States.

40 a. **Rissa tridactyla pollicaris** RIDGW.

Pacific Kittiwake.

*Rissa tridactyla pollicaris* "STEJN. MS." RIDGW. Water B. N. Am. II. 1884, 202.

[B —, C 552a, R 658a, C 783.]

GEOG. DIST.—Coasts of North Pacific and Bering Sea, south in winter casually to southern California.

41. **Rissa brevirostris** (BRUCH).

Red-legged Kittiwake.

*Larus brevirostris* BRUCH, J. f. O. 1853, 103.

*Rissa brevirostris* LAWR. in BAIRD'S B. N. Am. 1858, 855.

[B 674, 675, C 553, R 659, C 784.]

GEOG. DIST.—Coasts and islands of Bering Sea.

GENUS **LARUS** LINNÆUS.

*Larus* LINN. S. N. ed. 10, I. 1758, 136. Type, by elimination,  
*L. canus* LINN.

42. **Larus glaucus** BRÜNN.

Glaucous Gull.

*Larus glaucus* BRÜNN. Orn. Bor. 1764, 44.

[B 656, C 543, R 660, C 768.]

GEOG. DIST.—Arctic regions, south in winter in North America to the Great Lakes and Long Island.

42.1. **Larus barrovianus** RIDGW.

Point Barrow Gull.

*Larus barrovianus* RIDGW. Auk, III. July, 1886, 330.



[B —, C —, R —, C —.]

GEOG. DIST.— Bering Sea and contiguous waters ; northeastward to Point Barrow, southwestward to Japan.

43. **Larus leucopterus** FABER.  
Iceland Gull.

*Larus leucopterus* FABER, Prodr. Isl. Orn. 1822, 91.

[B 658, C 544, R 661, C 769.]

GEOG. DIST.— Arctic regions, south in winter in North America to Massachusetts and the Great Lakes, occasionally much further south.

44. **Larus glaucescens** NAUM.  
Glaucous-winged Gull.

*Larus glaucescens* NAUM. Naturg. Vög. Deutschl. X. 1840, 351.

[B 657, 659, C 545, R 662, C 770.]

GEOG. DIST.— Pacific coast of North America, from Alaska south to California ; on the Asiatic side south to Japan.

45. **Larus kumlieni** BREWST.  
Kumlien's Gull.

*Larus kumlieni* BREWST. Bull. Nutt. Orn. Club, VIII. 1883, 216.

[B —, C —, R —, C —.]

GEOG. DIST.— North Atlantic coast of North America, breeding in Cumberland Gulf ; south in winter to the coast of the Middle States.

46. **Larus nelsoni** HENSH.  
Nelson's Gull.

*Larus nelsoni* HENSH. Auk, I. July, 1884, 250.

[B —, C —, R —, C —.]

GEOG. DIST.— Coast of Norton Sound, Alaska.

47. **Larus marinus** LINN.

Great Black-backed Gull.

*Larus marinus* LINN. S. N. ed. 10, I. 1758, 136.

[B 660, C 546, R 663, C 771.]

GEOG. DIST.—Coasts of the North Atlantic; south in winter to Long Island and Italy.

48. **Larus schistisagus** STEJN.

Slaty-backed Gull.

*Larus schistisagus* STEJN. Auk, I. July, 1884, 231.

[B—, C—, R—, C—.]

GEOG. DIST.—North Pacific, chiefly on the Asiatic side; Herald Island, Arctic Ocean, and Alaskan coast of Bering Sea.

49. **Larus occidentalis** AUD.

Western Gull.

*Larus occidentalis* AUD. Orn. Biog. V. 1839, 320.[B 662, C 547*b*, R 664, C 774.]

GEOG. DIST.—Pacific coast of North America, breeding from Lower California northward to British Columbia.

[50.] **Larus affinis** REINH.

Siberian Gull.

*Larus affinis* REINH. Vid. Med. 1853, 78.

[B—, C—, R 665, C 776.]

GEOG. DIST.—Northern Asia and Europe, southward in winter to northern Africa. Accidental in Greenland.

51. **Larus argentatus** BRÜNN.

Herring Gull.

*Larus argentatus* BRÜNN. Orn. Bor. 1764, 44.

[B—, C 547, R 666, C 772.]

GEOG. DIST.—Old World, south to the Azores; Cumberland Sound; occasional on the eastern coast of the United States.

51 *a.* **Larus argentatus smithsonianus** COUES.  
American Herring Gull.

*Larus smithsonianus* COUES, Pr. Ac. Nat. Sci. Phila. 1862, 296.

*Larus argentatus* var. *smithsonianus* COUES, Check-List, 1873,  
no. 547*a*.

[B 661, C 547*a*, R 666*a*, C 773.]

GEOG. DIST.—North America generally, breeding from Maine, northern New York, the Great Lakes, and Minnesota northward; in winter south to Cuba and Lower California.

52. **Larus vegæ** (PALMÉN).  
Vega Gull.

*Larus argentatus* var. *vegæ* PALM. Bidr. Sibirisk. Vega Exp. 1887,  
370.

*Larus vegæ* STEJN. Auk, V. July, 1888, 310.

[B—, C —, R 667, C 775.]

GEOG. DIST.—Bering Sea and adjacent waters, south in winter to California and Japan.

53. **Larus californicus** LAWR.  
California Gull.

*Larus californicus* LAWR. Ann. Lyc. N. Y. VI. 1854, 79.

[B 663, C 548*a*, R 668, C 777.]

GEOG. DIST.—Western North America, chiefly in the interior, from Alaska to Mexico.

54. **Larus delawarensis** ORD.  
Ring-billed Gull.

*Larus delawarensis* ORD, GUTHRIE'S Geog. 2d Am. ed. 1815, 319.

[B 664, C 548, R 669, C 778.]

GEOG. DIST.—North America at large; south in winter to Cuba and Mexico.

55. **Larus brachyrhynchus** RICH.

Short-billed Gull.

*Larus brachyrhynchus* RICH. F. B. A. II. 1831, 421.

[B 665, 673, C 549, R 670, C 780.]

GEOG. DIST.— Arctic America and Pacific coast, south in winter to southern California.

[56.] **Larus canus** LINN.

Mew Gull.

*Larus canus* LINN. S. N. ed. 10, I. 1758, 136.

[B —, C —, R 671, C 779.]

GEOG. DIST.— Europe and Asia; accidental in Labrador?

57. **Larus heermanni** CASS.

Heermann's Gull.

*Larus heermanni* CASS. Pr. Ac. Nat. Sci. Phila. VI. 1852, 187.

[B 666, C 551, R 672, C 781.]

GEOG. DIST.— Pacific coast of North America, from British Columbia to Panama.

58. **Larus atricilla** LINN.

Laughing Gull.

*Larus atricilla* LINN. S. N. ed. 10, I. 1758, 136.

[B 667, C 554, R 673, C 786.]

GEOG. DIST.— Atlantic and Gulf coasts of the United States, north to Maine and Nova Scotia; south in winter through West Indies, Mexico (both coasts), Central America, and northern South America (Atlantic side) to the Lower Amazon.

59. **Larus franklinii** SW. & RICH.

Franklin's Gull.

*Larus franklinii* SW. & RICH. F. B. A. II. 1831, 424, pl. 71.



[B 668, 669, C 555, R 674, C 787.]

GEOG. DIST.—Interior of North America, chiefly west of the Mississippi River and east of the Rocky Mountains, breeding from Iowa northward; south in winter through Mexico and Central America to Peru.

60. **Larus philadelphia** (ORD).

Bonaparte's Gull.

*Sterna philadelphia* ORD, GUTHRIE'S Geog. 2d Am. ed. II. 1815, 319.

*Larus philadelphia* GRAY, List Brit. B. 1863, 235.

[B 670, C 556, R 675, C 788.]

GEOG. DIST.—Whole of North America, breeding mostly north of the United States. Not yet recorded from south of the United States, though reported from the Bermudas.

[60.1.] **Larus minutus** PALL.

Little Gull.

*Larus minutus* PALL. Reis. Russ. Reichs. III. App. No. 35, 1771, 702.

[B —, C —, R —, C —.]

GEOG. DIST.—Europe and parts of Asia and Africa; accidental in Bermuda and on Long Island.

GENUS **RHODOSTETHIA** MACGILLIVRAY.

*Rhodostethia* MACGIL. Man. Brit. Orn. II. 1842, 253. Type,  
*Larus roseus* MACGIL.

61. **Rhodostethia rosea** (MACGIL.).

Ross's Gull.

*Larus roseus* MACGIL. Mem. Wern. Soc. V. 1824, 249.

*Rhodostethia rosea* BONAP. Rev. Crit. Orn. Eur. Degland, 1850, 201.

[B 678, C 557, R 676, C 789.]

GEOG. DIST.—Arctic regions; south in autumn and winter to Kamchatka, Point Barrow, Alaska, and Disco Bay, Greenland. Casual in England, Faroes, Helgoland, etc.

GENUS **XEMA** LEACH.

*Xema* LEACH, ROSS'S Voy. App. 1819, p. lvii. Type, *Larus sabinii* SAB.

62. **Xema sabinii** (SAB.).  
Sabine's Gull.

*Larus sabinii* J. SAB. Trans. Linn. Soc. XII. 1818, 520, pl. 29.  
*Xema sabini* LEACH, App. ROSS'S Voy. Baff. Bay. 4to ed. 1819, lvii.

[B 680, C 558, R 677, C 790.]

GEOG. DIST.—Arctic regions; in North America south in winter to New York, the Great Lakes, and Great Salt Lake; casual in Kansas, Bahama, and on coast of Peru.

SUBFAMILY **STERNINÆ**. TERNS.GENUS **GEOCHELIDON** BREHM.

*Gelochelidon* BREHM, Isis, 1830, 994. Type, *G. meridionalis* BREHM = *Sterna nilotica* HASSELQ.

63. **Gelochelidon nilotica** (HASSELQ.).  
Gull-billed Tern.

*Sterna nilotica* HASSELQ. Reise nach Pal. Deutsche Ausg. 1762, 325.  
*Gelochelidon nilotica* STEJN. Auk, I. Oct. 1884, 366.

[B 681, C 560, R 679, C 792.]

GEOG. DIST.—Nearly cosmopolitan; in North America chiefly along the Atlantic and Gulf coasts of the United States, breeding north to southern New Jersey, and wandering casually to Long Island and Massachusetts; in winter both coasts of Mexico and Central America and south to Brazil.

GENUS **STERNA** LINNÆUS.SUBGENUS **THALASSEUS** BOIE.

*Thalasseus* BOIE, Isis, 1822, 563. Type, *Sterna caspia* PALL. =  
*S. tschegrava* LEPECH.

64. **Sterna tschegrava** LEPECH.  
Caspian Tern.

*Sterna tschegrava* LEPECH. Nov. Comm. Petrop. XIV. 1770, 500,  
pl. 13, fig. 2.

[B 682, C 561, R 680, C 793.]

GEOG. DIST.—Nearly cosmopolitan; in North America breeding southward to Virginia, Lake Michigan, Texas, Nevada, and California.

SUBGENUS **ACTOCHELIDON** KAUP.

*Actochelidon* KAUP, Sk. Ent. Eur. Thierw. 1829, 31. Type,  
*Sterna cantiaca* GMEL. = *S. sandvicensis* GMEL.

65. **Sterna maxima** BODD.  
Royal Tern.

*Sterna maxima* BODD. Tabl. P. E. 1783, 58.

[B 683, C 562, R 681, C 794.]

GEOG. DIST.—Tropical America, and warmer parts of North America, casually northward to Massachusetts, the Great Lakes, and California. West coast of Africa, north to Tangiers.

66. **Sterna elegans** GAMB.  
Elegant Tern.

*Sterna elegans* GAMB. Pr. Ac. Nat. Sci. Phila. IV. 1848, 129.

[B 684, C 563, R 682, C 795.]

GEOG. DIST.—Pacific coast of America, from California to Chili.

67. ***Sterna sandvicensis acuflavida*** (CABOT).

Cabot's Tern.

*Sterna acuflavida* CABOT, Pr. Boston Soc. N. H. II. 1847, 257.*Sterna sandvicensis acuflavida* RIDGW. Water B. N. Am. II. 1884, 288.

[B 685, C 564, R 683, C 796.]

GEOG. DIST.—South Atlantic and Gulf coasts of the United States, north, casually, to southern New England; south, in winter, to Central America (both coasts) and West Indies.

SUBGENUS **STERNA.**

*Sterna* LINN. S. N. ed. 10, I. 1758, 137. Type, by elimination,  
*S. hirundo* LINN.

[68.] ***Sterna trudeaui*** AUD.

Trudeau's Tern.

*Sterna trudeaui* AUD. Orn. Biog. V. 1839, 125, pl. 409.

[B 687, C 571, R 684, C 802.]

GEOG. DIST.—Southern South America. Casual, or accidental, on the Atlantic coast of the United States (New Jersey, Long Island).

69. ***Sterna forsteri*** NUTT.

Forster's Tern.

*Sterna forsteri* NUTT. Man. II. 1834, 274.

[B 691, 686, C 566, R 685, C 798.]

GEOG. DIST.—North America generally, breeding from Manitoba southward to Virginia, Illinois, Texas, and California; in winter southward to Brazil.

70. ***Sterna hirundo*** LINN.

Common Tern.

*Sterna hirundo* LINN. S. N. ed. 10, I. 1758, 137.



[B 689, C 565, R 686, C 797.]

GEOG. DIST.—Greater part of the northern hemisphere and Africa. In North America chiefly east of the Plains, breeding from the Arctic coast, somewhat irregularly, to Florida, Texas, and Arizona, and wintering northward to Virginia. Also coast of Lower California.

71. **Sterna paradisæa** BRÜNN.

Arctic Tern.

*Sterna paradisæa* BRÜNN. Orn. Bor. 1764, 46.

[B 690, 693, C 567, 568, R 687, C 799.]

GEOG. DIST.—Northern hemisphere; in North America breeding from Massachusetts to the Arctic regions, and wintering southward to Virginia and California.

72. **Sterna dougalli** MONTAG.

Roseate Tern.

*Sterna dougalli* MONTAG. Orn. Dict. Suppl. 1813, —.

[B 692, C 569, R 688, C 800.]

GEOG. DIST.—Temperate and tropical regions; north on the Atlantic coast of North America to Massachusetts, and casually to Maine and Nova Scotia.

73. **Sterna aleutica** BAIRD.

Aleutian Tern.

*Sterna aleutica* BAIRD, Tr. Chicago Ac. Nat. Sci. I. 1869, 321, pl. 31, fig. 1.

[B—, C 572, R 689, C 803.]

GEOG. DIST.—Coast of Alaska from Kadiak to Norton Sound.

SUBGENUS **STERNULA** BOIE.

*Sternula* BOIE, Isis, 1822, 563. Type, *Sterna minuta* LINN.

74. ***Sterna antillarum*** (LESS.).

Least Tern.

*Sternula antillarum* LESS. Descr. Mam. et Ois. 1847, 256.*Sterna antillarum* COUES, Pr. Ac. Nat. Sci. Phila. 1862, 552.

[B 694, C 570, R 690, C 801.]

GEOG. DIST.—Northern South America, northward to California, Minnesota, and New England, and casually to Labrador, breeding nearly throughout its range.

SUBGENUS **HALIPLANA** WAGLER.*Haliplana* WAGL. Isis, 1832, 1224. Type, *Sterna fuliginosa* GMEL.75. ***Sterna fuliginosa*** GMEL.

Sooty Tern.

*Sterna fuliginosa* GMEL. S. N. I. ii. 1788, 605.

[B 688, C 573, R 691, C 804.]

GEOG. DIST.—Tropical and subtropical coasts of the globe. In America from Chili to western Mexico and the Carolinas, and casually to New England.

[76.] ***Sterna anæthetus*** SCOP.

Bridled Tern.

*Sterna anæthetus* SCOP. Del. Faun. et Flor. Ins. II. 1786, no. 72, 92.

[B —, C 574, R 692, C 805.]

GEOG. DIST.—Tropical regions generally. Casual in Florida.

GENUS **HYDROCHELIDON** BOIE.*Hydrochelidon* BOIE, Isis, 1822, 563. Type, *Sterna nigra* LINN.77. ***Hydrochelidon nigra surinamensis*** (GMEL.).

Black Tern.

*Sterna surinamensis* GMEL. S. N. I. ii. 1788, 604.*Hydrochelidon nigra surinamensis* STEJN. Pr. U. S. Nat. Mus. 1882, 40.

[B 695, C 575, R 693, C 806.]

GEOG. DIST.—Temperate and tropical America. From Alaska and the Fur Countries to Brazil and Chili, breeding from the middle United States, west of the Alleghanies, northward.

[78.] **Hydrochelidon leucoptera** (MEISN. & SCHINZ).

**White-winged Black Tern.**

*Sterna leucoptera* MEISN. & SCHINZ, Vög. Schweiz, 1815, 264.

*Hydrochelidon leucoptera* BOIE, Isis, 1822, 563.

[B —, C 575 *bis*, R 694, C 807.]

GEOG. DIST.—Eastern hemisphere, accidental in North America (Lake Koshkonong, Wisconsin).

GENUS **ANOUS** STEPHENS.

*Anous* STEPH. Gen. Zool. XIII. pt. i. 1826, 139. Type, *Sterna stolidus* LINN.

79. **Anous stolidus** (LINN.).

**Noddy.**

*Sterna stolidus* LINN. S. N. ed. 10, I. 1758, 137.

*Anous stolidus* GRAY, List Gen. B. 1841, 100.

[B 696, C 576, R 695, C 808.]

GEOG. DIST.—Tropical and subtropical regions; in America from Brazil and Chili north to the Gulf and South Atlantic States.

FAMILY **RYNCHOPIDÆ**. SKIMMERS.

GENUS **RYNCHOPS** LINN.

*Rynchops* LINN. S. N. ed. 10, I. 1758, 138. Type, *R. nigra* LINN.

80. **Rynchops nigra** LINN.

**Black Skimmer.**

*Rynchops nigra* LINN. S. N. ed. 10, I. 1758, 138.

[B 697, C 577, R 656, C 809.]

GEOG. DIST.—Warmer parts of America, north on the Atlantic coast to New Jersey, and casually to the Bay of Fundy.

ORDER TUBINARES. TUBE-NOSED  
SWIMMERS.

FAMILY **DIOMEDEIDÆ**. ALBATROSSES.

GENUS **DIOMEDEA** LINNÆUS.

*Diomedea* LINN. S. N. ed. 10, I. 1758, 132. Type, *D. exulans*  
LINN.

81. **Diomedea nigripes** AUD.

**Black-footed Albatross.**

*Diomedea nigripes* AUD. Orn. Biog. V. 1839, 327.

[B —, C 579, R 700, C 811.]

GEOG. DIST.—North Pacific, including west coast of North America.

82. **Diomedea albatrus** PALL.

**Short-tailed Albatross.**

*Diomedea albatrus* PALL. Spic. Zool. V. 1769, 28.

[B 631, C 578, R 701, C 810.]

GEOG. DIST.—Pacific Ocean, including western coast of America,  
northward to Bering Sea.

GENUS **THALASSOGERON** RIDGWAY.

*Thalassogeron* RIDGW. Water B. N. Am. II. 1884, 357. Type,  
*Diomedea culminata* GOULD.

[83.] **Thalassogeron culminatus** (GOULD).

**Yellow-nosed Albatross.**

*Diomedea culminata* GOULD, P. Z. S. 1843, 107.

*Thalassogeron culminatus* RIDGW. Water B. N. Am. II. 1884, 358.

[B 632, C —, R 702, C —.]

GEOG. DIST.— Indian and South Pacific Oceans; casual off the coast of Oregon; accidental in the Gulf of St. Lawrence.

GENUS **PHŒBETRIA** REICHENBACH.

*Phœbetria* REICH. Syst. Av. 1852, p. v. Type, *Diomedea fuliginosa* GMEL.

84. **Phœbetria fuliginosa** (GMEL.).

Sooty Albatross.

*Diomedea fuliginosa* GMEL. S. N. I. ii. 1788, 568.

*Phœbetria fuliginosa* REICH. Syst. Av. 1852, p. v.

[B 633, C 580, R 703, C 812.]

GEOG. DIST.— Oceans of the southern hemisphere, northward to the coast of Oregon.

FAMILY **PROCELLARIIDÆ**. FULMARS AND SHEARWATERS.SUBFAMILY **PROCELLARIINÆ**. FULMARS.GENUS **OSSIFRAGA** HOMBRON & JACQUINOT.

*Ossifraga* HOMB. & JACQ. Compt. Rend. XVIII. 1844, 356.

Type, *Procellaria gigantea* GMEL.

[85.] **Ossifraga gigantea** (GMEL.).

Giant Fulmar.

*Procellaria gigantea* GMEL. S. N. I. ii. 1788, 563.

*Ossifraga gigantea* REICH. Syst. Av. 1852, p. iv.

[B 634, C 581, R 704, C 813.]

GEOG. DIST.— Southern Oceans; casual off the coast of Oregon.



GENUS **FULMARUS** STEPHENS.SUBGENUS **FULMARUS**.

*Fulmarus* STEPHENS, Gen. Zool. XIII. pt. i. 1826, 233. Type,  
*Procellaria glacialis* LINN.

86. **Fulmarus glacialis** (LINN.)**Fulmar.**

*Procellaria glacialis* LINN. Faun. Suec. 2d ed. 1761, 51; S. N. ed.  
12, I. 1766, 213.

*Fulmarus glacialis* STEPH. Gen. Zool. XIII. pt. i. 1826, 234, pl. 27.

[B 635, C 582, R 705, C 814.]

GEOG. DIST.—North Atlantic, south on the American coast to  
Massachusetts, casually to New Jersey.

86 a. **Fulmarus glacialis minor** (KJÆRBØLLING).**Lesser Fulmar.**

*Procellaria minor* KJÆRB. Danm. Fugle, 1852, 324.

*Fulmarus glacialis* b. *minor* BONAP. Consp. II. 1856, 187.

[B —, C —, R —, C —.]

GEOG. DIST.—North Atlantic, south on American side to coast of  
Massachusetts.

86 b. **Fulmarus glacialis glupischa** STEJN.**Pacific Fulmar.**

*Fulmarus glacialis glupischa* STEJN. Auk, I. July, 1884, 234.

[B 636, C 582a, R 705a, C 815.]

GEOG. DIST.—North Pacific, south on the American coast to  
Mexico.

86 c. **Fulmarus glacialis rodgersii** (CASS.).**Rodgers's Fulmar.**

*Fulmarus rodgersii* CASS. Pr. Ac. Nat. Sci. Phila. 1862, 290.

*Fulmarus glacialis* var. *rodgersi* COUES, Key, 1872, 327.

[B —, C 582*b*, R 705*b*, C 816.]

GEOG. DIST.— Bering Sea and adjacent parts of the North Pacific.

SUBGENUS **PRIOCELLA**. HOMBRON & JACQUINOT.

*Priocella* HOMB. & JACQ. Compt. Rend. XVIII. 1844, 357.

Type, *P. garnoti* HOMB. & JACQ. = *Procellaria glacialis* SMITH.

87. **Fulmarus glacialis** (SMITH).

Slender-billed Fulmar.

*Procellaria glacialis* SMITH, Illustr. S. Afr. B. 1840, t. 51.

*Fulmarus glacialis* STEJN. Auk, 1884, p. 233.

[B 637, C 583, R 706, C 817.]

GEOG. DIST.— Seas of the southern hemisphere, and northward along Pacific coast of North America to Washington.

GENUS **PUFFINUS** BRISSON.

*Puffinus* BRISS. Orn. VI. 1760, 131. Type, *Procellaria puffinus* BRÜNN.

88. **Puffinus borealis** CORY.

Cory's Shearwater.

*Puffinus borealis* CORY, Bull. Nutt. Orn. Club, VI. April, 1881, 84.

[B —, C —, R —, C 888.]

GEOG. DIST.— Known only from off the coasts of Massachusetts, Rhode Island, and Long Island.

89. **Puffinus major** FABER.

Greater Shearwater.

*Puffinus major* FABER, Prodr. Isl. Orn. 1822, 56.

[B 647, C 597, R 709, C 832.]

GEOG. DIST.— Atlantic Ocean, from Cape Horn and Cape of Good Hope northward to the Arctic Circle.

[90.] **Puffinus puffinus** (BRÜNN.).**Manx Shearwater.***Procellaria puffinus* BRÜNN. Orn. Bor. 1764, 29.*Puffinus puffinus* LICHT. Nomencl. Mus. Berol. 1854, 100.

[B 649, C 599, R 711, C 834.]

GEOG. DIST.—North Atlantic, chiefly on the eastern side; accidental in Greenland, and rare or casual off the North American coast(?).

91. **Puffinus creatopus** COUES.**Pink-footed Shearwater.**

*Puffinus creatopus* "COOPER, MS.," COUES, Pr. Ac. Nat. Sci. Phila. April, 1864, 131.

[B —, C 598, R 710, C 833.]

GEOG. DIST.—Pacific Ocean; on the American coast from California to Juan Fernandez Islands.

92. **Puffinus auduboni** FINSCH.**Audubon's Shearwater.***Puffinus auduboni* FINSCH, P. Z. S. 1872, 111.

[B 650, C 600, R 712, C 835.]

GEOG. DIST.—Warmer parts of the Atlantic, north casually to New Jersey and Long Island.

93. **Puffinus gavia** (FORST.).**Black-vented Shearwater.***Procellaria gavia* FORST. Descr. An. 1844, 148.*Puffinus gavia* FINSCH, J. f. O. 1872, 256.

[B —, C 601, R 713, C 836.]

GEOG. DIST.—Pacific Ocean, chiefly southward; coast of Lower California, north to Santa Cruz, Cal.

94. **Puffinus stricklandi** RIDGW.**Sooty Shearwater.***Puffinus stricklandi* RIDGW. Water B. N. Am. II. 1884, 390.

[B 648, C 602, R 714, C 837.]

GEOG. DIST.—Atlantic Ocean, breeding in the southern hemisphere; a summer visitor off our coast, from South Carolina northward.

95. **Puffinus griseus** (GMEL.).

Dark-bodied Shearwater.

*Procellaria grisea* GMEL. S. N. I. ii. 1788, 564.*Puffinus griseus* FINSCH, J. f. O. 1874, 209.

[B —, C 603, R 715, C 838.]

GEOG. DIST.—South Pacific, north on the American coast, casually to San Francisco, California.

96. **Puffinus tenuirostris** (TEMM.).

Slender-billed Shearwater.

*Procellaria tenuirostris* TEMM. Pl. Col. 1828, 587.*Puffinus tenuirostris* TEMM. & SCHLEG. Faun. Jap. Aves, 1849, 131, pl. 86.

[B —, C 604, R 716, C 839.]

GEOG. DIST.—Both coasts of the North Pacific, from Japan and Kotzebue Sound south to Australia and New Zealand; on the American coast to British Columbia.

SUBGENUS **PRIOFINUS** HOMBRON & JACQUINOT.*Priofinus* HOMBR. & JACQ. Compt. Rend. XVIII. 1844, 355.Type, *Procellaria cinerea* GMEL.[97.] **Puffinus cinereus** (GMEL.).

Black-tailed Shearwater.

*Procellaria cinerea* GMEL. S. N. I. ii. 1788, 563.*Puffinus cinereus* LAWR. in BAIRD'S B. N. Am. 1858, 835.

[B 651, C 595, R 707, C 830.]

GEOG. DIST.—South Pacific; accidental off the coast of California.

GENUS **ÆSTRELATA** BONAPARTE.

*Æstrelata* BONAP. Cons. II. 1856, 188. Type, *Procellaria hasitata* KUHL.

[98.] **Æstrelata hasitata** (KUHL).

Black-capped Petrel.

*Procellaria hasitata* KUHL, Mon. Proc. Beitr. Zool. 1 Abt. 1820, 142.

*Æstrelata hasitata* COUES, Pr. Ac. Nat. Sci. Phila. 1866, 139.

[B 638, C 585, R 717, C 819.]

GEOG. DIST.—Warmer parts of the Atlantic Ocean, straying to Florida, Virginia, New York (Ulster County, Oneida Lake and Long Island), Vermont and Ontario. Also England and France.

[99.] **Æstrelata scalaris** BREWST.

Scaled Petrel.

*Æstrelata scalaris* BREWST. Auk, III. July, 1886, 300.

[B—, C—, R—, C—.]

GEOG. DIST.—Unknown, the single specimen obtained having been taken in Livingston Co., N. Y., in April, 1880.

100. **Æstrelata fisheri** RIDGW.

Fisher's Petrel.

*Æstrelata fisheri* RIDGW. Pr. U. S. Nat. Mus. V. June 26, 1883, 656.

[B—, C—, R—, C—.]

GEOG. DIST.—The only known specimen was taken at Kadiak, Alaska.

GENUS **BULWERIA** BONAPARTE.

*Bulweria* BONAP. Cat. Met. Ucc. Eur. 1842, 81. Type, *Procellaria bulweri* JARD. & SELBY.



[101.] **Bulweria bulweri** (JARD. & SELBY).**Bulwer's Petrel.***Procellaria bulweri* JARD. & SELBY, Illustr. Orn. —, pl. 65.*Bulweria bulweri* BOUCARD, Cat. Av. 1876, 69.

[B —, C —, R 718, C 820.]

GEOG. DIST. — Eastern Atlantic, including coasts of Europe and Africa. Accidental in Greenland. Also various parts of the Pacific Ocean.

GENUS **DAPTION** STEPHENS.

*Daption* STEPH. Gen. Zool. XIII. 1825, 239. Type, *Procellaria capensis* LINN.

[102.] **Daption capensis** (LINN.).**Pintado Petrel.***Procellaria capensis* LINN. S. N. ed. 10, I. 1758, 132.*Daption capensis* STEPH. Gen. Zool. XIII. pt. i. 1825, 241.

[B 639, C 584, R 719, C 818.]

GEOG. DIST. — Oceans of the southern hemisphere, north to about latitude 25°. Accidental on the coasts of California and England.

GENUS **HALOCYPTENA** COUES.*Halocyptena* COUES, Pr. Ac. Nat. Sci. Phila. March, 1864, 78.Type, *H. microsoma* COUES.103. **Halocyptena microsoma** COUES.**Least Petrel.***Halocyptena microsoma* COUES, Pr. Ac. Nat. Sci. Phila. 1864, 79.

[B —, C 586, R 720, C 821.]

GEOG. DIST. — Coast of Lower California and south to Panama.

GENUS **PROCELLARIA** LINNÆUS.

*Procellaria* LINN. S. N. ed. 10, I. 1758, 131. Type, by elimination, *P. pelagica* LINN.

104. ***Procellaria pelagica*** LINN.

Stormy Petrel.

*Procellaria pelagica* LINN. S. N. ed. 10, I. 1758, 131.

[B 645, C 587, R 721, C 822.]

GEOG. DIST.—Atlantic Ocean, south on the American side to the Newfoundland Banks. West coast of Africa and coast of Europe.

GENUS **OCEANODROMA** REICHENBACH.

*Oceanodroma* REICH. Syst. Av. 1852, p. iv. Type, *Procellaria furcata* GMEL.

105. ***Oceanodroma furcata*** (GMEL.).

Forked-tailed Petrel.

*Procellaria furcata* GMEL. S. N. I. ii. 1788, 561.

*Oceanodroma furcata* REICH. Syst. Av. 1852, p. iv.

[B 640, C 591, R 726, C 826.]

GEOG. DIST.—North Pacific, south on the American coast, in winter, to Humboldt Bay, California. Breeds on the Aleutian Islands and on Copper Island.

106. ***Oceanodroma leucorhoa*** (VIEILL.).

Leach's Petrel.

*Procellaria leucorhoa* VIEILL. Nouv. Dict. d'Hist. Nat. XXV.

1817, 422.

*Oceanodroma leucorhoa* STEJN. Orn. Expl. Kamtsch. 1885, 97.

[B 642, C 588, R 723, C 823.]

GEOG. DIST.—North Atlantic and North Pacific Oceans; south on the coast of the United States to Virginia and California; breeds from Maine and the Hebrides northward on the coasts of the Atlantic; also on the Aleutian Islands and on Copper Island, Bering Sea.

106.1. **Oceanodroma macrodactyla** (BRYANT).

Guadalupe Petrel.

*Oceanodroma leucorhoa macrodactyla* BRYANT, Bull. Cal. Ac. Sci. II. No. 8, July 23, 1887, 450.*Oceanodroma macrodactyla* A. O. U. COMM. Suppl. Check-List N. A. Birds, 1889, 5.

[B—, C—, R—, C—.]

GEOG. DIST.—Guadalupe Island, Lower California.

107. **Oceanodroma melania** (BONAP.).

Black Petrel.

*Procellaria melania* BONAP. Compt. Rend. XXVIII. 1854, 662.*Oceanodroma melania* STEJN. Orn. Expl. Kamtsch. 1885, 371.

[B—, C 589, R 724, C 824.]

GEOG. DIST.—South Pacific, northward to southern California.

108. **Oceanodroma homochroa** (COUES).

Ashy Petrel.

*Cymochorea homochroa* COUES, Pr. Ac. Nat. Sci. Phila. 1864, 77.*Oceanodroma homochroa* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 356.

[B 643, C 590, R 725, C 825.]

GEOG. DIST.—Coast of California. Breeds on the Santa Barbara and Farallone Islands.

## SUBFAMILY OCEANITINÆ.

## GENUS OCEANITES KEYSERLING &amp; BLASIUS.

*Oceanites* KEYS. & BLAS. Wirb. Eur. I. 1840, xciii, 131, 238.  
Type, *Procellaria oceanica* KUHL.109. **Oceanites oceanicus** (KUHL).

Wilson's Petrel.

*Procellaria oceanica* KUHL, Beitr. Zool. Mon. Proc. 1820, 136,  
pl. 10, fig. 1.*Oceanites oceanica* LICHT. Nomencl. Mus. Berol. 1854, 99.

[B 644, C 593, R 722, C 828.]

GEOG. DIST.—North and South Atlantic and Southern Oceans, breeding on Kerguelen Island in February. Abundant off the eastern coast of the United States during the entire summer.

GENUS **CYMODROMA** RIDGWAY.

*Cymodroma* RIDGW. Water B. N. Am. II. 1884, 418. Type, *Procellaria grallaria* VIEILL.

[110.] **Cymodroma grallaria** (VIEILL.).

White-bellied Petrel.

*Procellaria grallaria* VIEILL. Nouv. Dict. d'Hist. Nat. XXVI. 1817, 418.

*Cymodroma grallaria* RIDGW. Water B. N. Am. II. 1884, 419.

[B 646, C 594, R 728, C 829.]

GEOG. DIST.—Tropical oceans generally; accidental on the coast of Florida.

GENUS **PELAGODROMA** REICHENBACH.

*Pelagodroma* REICH. Syst. Av. 1852, p. iv. Type, *Procellaria marina* LATH.

[111.] **Pelagodroma marina** (LATH.).

White-faced Petrel.

*Procellaria marina* LATH. Ind. Orn. II. ii. 1790, 826.

*Pelagodroma marina* REICH. Syst. Av. 1852, p. iv.

[B —, C —, R —, C —.]

GEOG. DIST.—South Atlantic, and Southern Seas. Accidental off the coast of Massachusetts.

# ORDER STEGANOPODES. TOTIPALMATE SWIMMERS.

FAMILY **PHAËTHONTIDÆ**. TROPIC BIRDS.

GENUS **PHAËTHON** LINNÆUS.

*Phaëthon* LINN. S. N. ed. 10, I. 1758, 134. Type, *P. æthereus* LINN.

## 112. *Phaëthon flavirostris* BRANDT.

Yellow-billed Tropic Bird.

*Phaëthon flavirostris* BRANDT, Bull. Ac. St. Pétersb. II. 1837, 349.

[B 629, C 538, R 654, C 763.]

GEOG. DIST.—West Indies and Atlantic coast of Central America, north to Florida and Bermuda; accidental in western New York. South Pacific and Indian Oceans.

## 113. *Phaëthon æthereus* LINN.

Red-billed Tropic Bird.

*Phaëthon æthereus* LINN. S. N. ed. 10, I. 1758, 134.

[B —, C —, R 655, C 762.]

GEOG. DIST.—Coasts of tropical America, north on the Pacific coast to Cape Colnett, Lower California; accidental on the Newfoundland Banks. Breeds on San Pedro Martir and other islands in the Gulf of California.

FAMILY **SULIDÆ**. GANNETS.

GENUS **SULA** BRISSON.

SUBGENUS **SULA**.

*Sula* BRISS. Orn. VI. 1760, 495. Type, by elimination, *Pelecanus sula* LINN.



[114.] ***Sula cyanops* SUND.****Blue-faced Booby.***Dysporus cyanops* SUND. Phys. Tidskr. Lund, 1837, pt. 5.*Sula cyanops* SUND. Isis, 1842, 858.

[B—, C—, R 651, C—.]

GEOG. DIST.—South Atlantic, South Pacific, and Indian Oceans;  
West Indies, and northward to southern Florida.

114.1. ***Sula gossi* GOSS.****Blue-footed Booby.***Sula gossi* (RIDGW. MS.) GOSS, Auk, V. July, 1888, 241.

[B—, C—, R—, C—.]

GEOG. DIST.—Islands in the Gulf of California, and south to the  
Galapagos.

115. ***Sula sula* (LINN.).****Booby.***Pelecanus sula* LINN. Syst. Nat. ed. 12, I. 1766, 218.*Sula sula* RIDGW. Pr. U. S. Nat. Mus. VIII, 1885, 356.

[B 618, C 525, R 652, C 747.]

GEOG. DIST.—Atlantic coasts of tropical and subtropical America,  
north to Georgia. Also, western Pacific and Indian Oceans.

115.1. ***Sula brewsteri* GOSS.****Brewster's Booby.***Sula brewsteri* GOSS, Auk, V. July, 1888, 242.

[B—, C—, R—, C—.]

GEOG. DIST.—Coasts and islands of the eastern South Pacific  
Ocean, north to Lower California; breeding as far north as Georges  
Island at the head of the Gulf of California.

[116.] ***Sula piscator* (LINN.).****Red-footed Booby.***Pelecanus piscator* LINN. S. N. ed. 10, I. 1758, 134.*Sula piscator* GRAY, Gen. Bds. III. 1845, 666.

[B —, C —, R 653, C —.]

GEOG. DIST.—Coasts and islands of tropical and subtropical seas, north to western Mexico and Florida.

SUBGENUS **DYSPORUS** ILLIGER.

*Dysporus* ILLIG. Prodr. 1811, 279. Type, by elimination, *Pelecanus bassanus* LINN.

117. **Sula bassana** (LINN.).

Gannet.

*Pelecanus bassanus* LINN. S. N. ed. 10, I. 1758, 133.

*Sula bassana* BOIE, Isis, 1822, p. 563.

[B 617, C 524, R 650, C 746.]

GEOG. DIST.—Coasts of the North Atlantic, south in winter to the Gulf of Mexico and Africa; breeds from Nova Scotia and the British Islands northward.

FAMILY **ANHINGIDÆ**. DARTERS.GENUS **ANHINGA** BRISSON.

*Anhinga* BRISSON, Orn. VI. 1760, 476. Type, *Anhinga* MARCGR.  
= *Plotus anhinga* LINN.

118. **Anhinga anhinga** (LINN.).

Anhinga.

*Plotus anhinga* LINN. S. N. ed. 12, I. 1766, 218.

*Anhinga anhinga* STEJN. Stand. Nat. Hist. IV. 1885, 193.

[B 628, C 536, R 649, C 760.]

GEOG. DIST.—Tropical and subtropical America, north in the United States to the Carolinas, the mouth of the Ohio River, and southern Kansas.

FAMILY **PHALACROCORACIDÆ**. CORMORANTS.GENUS **PHALACROCORAX** BRISSON.SUBGENUS **PHALACROCORAX**.

*Phalacrocorax* BRISS. Orn. VI. 1760, 511. Type, *Pelecanus carbo* LINN.

119. **Phalacrocorax carbo** (LINN.).**Cormorant.**

*Pelecanus carbo* LINN. S. N. ed. 10, I. 1758, 133.

*Phalacrocorax carbo* LEACH, Syst. Cat. Mam. and Bds. Brit. Mus. 1816, 34.

[B 620, C 528, R 642, C 750.]

GEOG. DIST.—Coasts of the North Atlantic, south in winter on the coast of the United States, casually, to the Carolinas; breeding (formerly) from Massachusetts northward.

120. **Phalacrocorax dilophus** (SWAIN.).**Double-crested Cormorant.**

*Pelecanus (Carbo) dilophus* SWAIN. in SW. & RICH. F. B. A. II. 1831, 473.

*Phalacrocorax dilophus* NUTT. Man. II. 1834, 483.

[B 623, C 530, R 643, C 751.]

GEOG. DIST.—Eastern North America, breeding from the Bay of Fundy, the Great Lakes, Minnesota and Dakota northward; south in winter to the Southern States.

120 a. **Phalacrocorax dilophus floridanus** (AUD.).**Florida Cormorant.**

*Phalacrocorax floridanus* AUD. Orn. Biog. III. 1835, 387.

*Phalacrocorax dilophus floridanus* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880, 205.

[B 624, C 530a, R 643a, C 753.]

GEOG. DIST.—South Atlantic and Gulf States, northward in the Mississippi Valley to southern Illinois.

120 *b*. **Phalacrocorax dilophus cincinatus** (BRANDT).

White-crested Cormorant.

*Carbo cincinatus* BRANDT, Bull. Sc. Ac. St. Pétersb. III. 1838, 55.

*Phalacrocorax dilophus cincinnatus* RIDGW. Pr. U. S. Nat. Mus.

III. Aug. 24, 1880, 205.

[B 622, C 529, R 643b, C 752.]

GEOG. DIST.—West coast of North America, south in winter to California.

120 *c*. **Phalacrocorax dilophus albociliatus** RIDGW.

Farallone Cormorant.

*Phalacrocorax dilophus albociliatus* RIDGW. Proc. Biol. Soc.

Wash. II. Apr. 10, 1884, 94.

[B —, C —, R —, C —.]

GEOG. DIST.—California, south to Cape St. Lucas and the Revillagigedo Islands.

121. **Phalacrocorax mexicanus** (BRANDT).

Mexican Cormorant.

*Carbo mexicanus* BRANDT, Bull. Sc. Ac. St. Pétersb. III. 1838, 55.

*Phalacrocorax mexicanus* SCL. & SALV. Nom. Neotr. 1873, 124.

[B 625, C 531, R 644, C 754.]

GEOG. DIST.—West Indies and Central America to southern United States; north in the interior to Kansas and southern Illinois.

SUBGENUS **COMPSOHALIEUS** RIDGWAY.

*Compsohalieus* RIDGW. Water B. N. Am. II. 1884, 145. Type,

*Carbo penicillatus* BRANDT.

122. **Phalacrocorax penicillatus** (BRANDT).

Brandt's Cormorant.

*Carbo penicillatus* BRANDT, Bull. Sc. Ac. St. Pétersb. III. 1838, 55.*Phalacrocorax penicillatus* HEERM. Pr. Ac. Nat. Sci. Phila. VII.  
1854, 178.

[B 626, C 532, R 645, C 755.]

GEOG. DIST.—Pacific coast of North America, from Cape St. Lucas to Washington.

SUBGENUS **URILE** BONAPARTE.*Urile* BONAP. Consp. II. 1855, 175. Type, *Pelecanus urile* GMEL.123. **Phalacrocorax pelagicus** PALL.

Pelagic Cormorant.

*Phalacrocorax pelagicus* PALL. Zoog. Rosso-As. II. 1826, 303.

[B —, C —, R —, C —.]

GEOG. DIST.—Aleutian and Kuril Islands, and Kamchatka, south to Japan.

123 a. **Phalacrocorax pelagicus robustus** RIDGW.

Violet-green Cormorant.

*Phalacrocorax pelagicus robustus* RIDGW. Water B. N. Am. II.  
1884, 160.

[B 627, C 535, R 646, C 758.]

GEOG. DIST.—Coast of Alaska, from Norton Sound south to Washington.

123 b. **Phalacrocorax pelagicus resplendens** (AUD.).

Baird's Cormorant.

*Phalacrocorax resplendens* AUD. Orn. Biog. V. 1839, 148.*Phalacrocorax pelagicus resplendens* RIDGW. Water B. N. Am. I.  
1884, 160.



[B —, C —, R 646a, C 759.]

GEOG. DIST.—Pacific coast of North America, from Washington south to Cape St. Lucas and Mazatlan, Mexico.

124. **Phalacrocorax urile** (GMEL.).

Red-faced Cormorant.

*Pelecanus urile* GMEL. S. N. I. ii. 1788, 575.

*Phalacrocorax urile* RIDGW. Water B. N. Am. II. 1884, 162.

[B —, C 534, R 647, C 757.]

GEOG. DIST.—Pribilof, Aleutian and Kuril Islands, and coast of Kamchatka. South in winter to northern Japan.

FAMILY **PELECANIDÆ**. PELICANS.

GENUS **PELECANUS** LINNÆUS.

*Pelecanus* LINN. S. N. ed. 10, I. 1758, 132. Type, by elimination, *P. onocrotalus* LINN.

SUBGENUS **CYRTOPELICANUS** REICHENBACH.

*Cyrtopelicanus* REICH. Syst. Av. 1852, p. vii. Type, *Pelecanus erythrorhynchos* GMEL.

125. **Pelecanus erythrorhynchos** GMEL.

American White Pelican.

*Pelecanus erythrorhynchos* GMEL. S. N. I. ii. 1788, 571.

[B 615, C 526, R 640, C 748.]

GEOG. DIST.—Temperate North America, north in the interior to about Lat. 61°, south in winter to western Mexico and Guatemala; now rare or accidental in the northeastern States; abundant in the interior and along the Gulf coast; common on the coast of California.

SUBGENUS **LEPTOPELICANUS** REICHENBACH.

*Leptopelicanus* REICH. Syst. Av. 1852, p. vii. Type, *Pelecanus fuscus* LINN.

126. **Pelecanus fuscus** LINN.

Brown Pelican.

*Pelecanus fuscus* LINN. S. N. ed. 12, I. 1766, 215.

[B 616, C 527, R 641, C 749.]

GEOG. DIST.—Atlantic coast of tropical and subtropical America, north on the Atlantic coast to North Carolina ; accidental in Illinois.

127. **Pelecanus californicus** RIDGW.

California Brown Pelican.

*Pelecanus (fuscus ?) californicus* RIDGW. Water B. N. Am. II. 1884, 143.

*P[elecanus] californicus* RIDGW. l. c.

[B —, C —, R —, C —.]

GEOG. DIST.—Pacific coast, from Burrard Inlet, British Columbia, to the Galapagos.

FAMILY **FREGATIDÆ**. MAN-O'-WAR BIRDS.GENUS **FREGATA** BRISSON.

*Fregata* BRISS. Orn. VI. 1760, 506. Type, *Pelecanus aquilus* LINN.

128. **Fregata aquila** LINN.

Man-o'-War Bird.

*Pelecanus aquilus* LINN. S. N. ed. 10, I. 1758, 133.

*Fregata aquila* REICH. Syst. Av. 1852, p. vi.

[B 619, C 537, R 639, C 761.]

GEOG. DIST.—Tropical and subtropical coasts generally ; in America, north to Florida and Texas, and casually to Nova Scotia, Ohio, Wisconsin, Kansas, and Humboldt Bay, California.

# ORDER ANSERES. LAMELLIROSTRAL SWIMMERS.

FAMILY **ANATIDÆ**. DUCKS, GEESE, AND SWANS.

SUBFAMILY **MERGINÆ**. MERGANSERS.

GENUS **MERGANSER** BRISSON.

*Merganser* BRISS. Orn. VI. 1760, 230. Type, *Mergus merganser*  
LINN.

## 129. **Merganser americanus** (CASS.).

American Merganser.

*Mergus americanus* CASSIN, Pr. Ac. Nat. Sci. Phila. VI. 1853, 187.

*Merganser americanus* STEJN. Orn. Expl. Kamtsch. 1885, 177.

[B —, C —, R —, C —.]

GEOG. DIST.—North America generally, breeding south in the United States to Pennsylvania and the mountains of Colorado and California. Bermuda.

## 130. **Merganser serrator** (LINN.).

Red-breasted Merganser.

*Mergus serrator* LINN. S. N. ed. 10, I. 1758, 129.

*Merganser serrator* SCHÄFFER, Mus. Orn. 1789, 66.

[B 612, C 522, R 637, C 744.]

GEOG. DIST.—Northern portions of northern hemisphere; south, in winter, throughout the United States.

GENUS **LOPHODYTES** REICHENBACH.

*Lophodytes* REICHENBACH, Syst. Av. 1852, p. ix. Type, *Mergus cucullatus* LINN.

131. **Lophodytes cucullatus** (LINN.).

Hooded Merganser.

*Mergus cucullatus* LINN. S. N. ed. 10, I. 1758, 129.*Lophodytes cucullatus* REICHENBACH, Syst. Av. 1852, p. ix.

[B 613, C 523, R 638, C 745.]

GEOG. DIST.—North America generally, south to Mexico and Cuba, breeding nearly throughout its range. Casual in Europe.

SUBFAMILY **ANATINÆ**. RIVER DUCKS.GENUS **ANAS** LINNÆUS.*Anas* LINN. S. N. ed. 10, I, 1758, 122. Type, *A. boschas* LINN.132. **Anas boschas** LINN.

Mallard.

*Anas boschas* LINN. S. N. ed. 10, I, 1758, 127.

[B 576, C 488, R 601, C 707.]

GEOG. DIST.—Northern parts of northern hemisphere; in America south to Panama and Cuba, breeding southward to the southern United States; less common in the East.

133. **Anas obscura** GMEL.

Black Duck.

*Anas obscura* GMEL. S. N. I. 1788, 541.

[B 577, C 489, R 602, C 708.]

GEOG. DIST.—Eastern North America, west to the Mississippi Valley, north to Labrador, breeding southward to the northern parts of the United States.

134. **Anas fulvigula** RIDGW.

Florida Duck.

*Anas obscura* var. *fulvigula* RIDGW. Am. Nat. VIII. Feb. 1874, 111.*Anas fulvigula* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880,

203.

[B —, C 489a, R 603, C 709.]

GEOG. DIST. — Florida.

134a. **Anas fulvigula maculosa** (SENN.).

Mottled Duck.

*Anas maculosa* SENNETT, Auk, VI, July, 1889, 263.[B 577, *part*, C 489, *part*, R 602, *part*, C 708, *part*.]

GEOG. DIST. — Eastern Texas and north to Kansas.

SUBGENUS **CHAULELASMUS** BONAPARTE.*Chaulelasmus* BONAP. Comp. List, 1838, 56. Type, *Anas strepera* LINN.135. **Anas strepera** LINN.

Gadwall.

*Anas strepera* LINN. S. N. ed. 10, I. 1758, 125.

[B 584, C 491, R 604, C 711.]

GEOG. DIST. — Nearly cosmopolitan. In North America breeds chiefly within the United States.

SUBGENUS **MARECA** STEPHENS.*Mareca* STEPHENS, Gen. Zool. XII. pt. ii. 1824, 130. Type, *Anas penelope* LINN.136. **Anas penelope** LINN.

Widgeon.

*Anas penelope* LINN. S. N. ed. 10, I. 1758, 126.

[B 586, C 492, R 606, C 712.]

GEOG. DIST. — Northern parts of the Old World. In North America breeds in the Aleutian Islands, and occurs frequently in the eastern United States, and occasionally in California.

137. **Anas americana** GMEL.

Baldpate.

*Anas americana* GMELIN, S. N. I. 1788, 526.



[B 585, C 493, R 607, C 713.]

GEOG. DIST.—North America, from the Arctic Ocean south, in winter, to Guatemala and Cuba. Breeds chiefly north of United States.

SUBGENUS **NETTION** KAUP.

*Nettion* KAUP, Sk. Ent. Europ. Thierw. 1829, 95. Type, *Anas crecca* LINN.

[138.] **Anas crecca** LINN.  
European Teal.

*Anas crecca* LINN. S. N. ed. 10, I. 1758, 126.

[B 580, C 494, R 611, C 714.]

GEOG. DIST.—Northern parts of the Old World. Occasional in eastern North America, the Aleutian Islands, and California.

139. **Anas carolinensis** GMEL.  
Green-winged Teal.

*Anas carolinensis* GMEL. S. N. I. 1788, 533.

[B 579, C 495, R 612, C 715.]

GEOG. DIST.—North America, breeding chiefly north of the United States, and migrating south to Honduras and Cuba.

SUBGENUS **QUERQUEDULA** STEPHENS.

*Querquedula* STEPHENS, Gen. Zool. XII. pt. ii. 1824, 142. Type, *Anas querquedula* LINN.

140. **Anas discors** LINN.  
Blue-winged Teal.

*Anas discors* LINN. S. N. ed. 12, I. 1766, 205.

[B 581, C 496, R 609, C 716.]

GEOG. DIST.—North America in general, but chiefly eastward; north to Alaska, and south to the West Indies, Lower California, and northern South America. Casual in California. Breeds from Kansas and southern Illinois northward.

141. **Anas cyanoptera** VIEILL.  
Cinnamon Teal.

*Anas cyanoptera* VIEILLOT, Nouv. Dict. d'Hist. Nat. V. 1816, 104.

[B 582, C 497, R 610, C 717.]

GEOG. DIST.—Western America from British Columbia south to Chili, Patagonia, and Falkland Islands; east in North America to the Rocky Mountains and southern Texas; casual in the Mississippi Valley and Florida.

GENUS **SPATULA** BOIE.

*Spatula* BOIE, Isis, 1822, 564. Type, *Anas clypeata* LINN.

142. **Spatula clypeata** (LINN.).  
Shoveller.

*Anas clypeata* LINN. S. N. ed. 10, I. 1758, 124.

*Spatula clypeata* BOIE, Isis, 1822, 564.

[B 583, C 498, R 608, C 718.]

GEOG. DIST.—Northern hemisphere. In North America breeding from Alaska to Texas; not abundant on the Atlantic coast north of the Carolinas.

GENUS **DAFILA** STEPHENS.

*Dafila* STEPHENS, Gen. Zool. XII. pt. ii. 1824, 126.

143. **Dafila acuta** (LINN.).  
Pintail.

*Anas acuta* LINN. S. N. ed. 10, I. 1758, 126.

*Dafila acuta* BONAP. Comp. List, 1838, 56.

[B 578, C 490, R 605, C 710.]

GEOG. DIST.—Northern hemisphere. In North America breeds from the northern parts of the United States northward, and migrates south to Panama and Cuba.

GENUS **AIX** BOIE.

*Aix* BOIE, Isis, 1828, 329. Type, *Anas sponsa* LINN.

144. **Aix sponsa** (LINN.).

Wood Duck.

*Anas sponsa* LINN. S. N. ed. 10, I. 1758, 128.

*Aix sponsa* BONAP. Comp. List, 1838, 57.

[B 587, C 499, R 613, C 719.]

GEOG. DIST.— Temperate North America, breeding throughout its range. Cuba. Accidental in Europe.

SUBFAMILY **FULIGULINÆ**. SEA DUCKS.GENUS **NETTA** KAUP.

*Netta* KAUP, Sk. Ent. Europ. Thierw. 1829, 102. Type, *Anas rufina* PALL.

[145.] **Netta rufina** (PALL.).

Rufous-crested Duck.

*Anas rufina* PALL. It. II. App. 1773, 713.

*Netta rufina* KAUP, Sk. Ent. Europ. Thierw. 1829, 102.

[B —, C —, R —, C 886.]

GEOG. DIST.— Eastern hemisphere; accidental in eastern United States.

GENUS **AYTHYA** BOIE.

*Aythya* BOIE, Isis, 1822, 564. Type, by elimination, *Anas ferina* LINN.

146. **Aythya americana** (EYT.).

Redhead.

*Fuligula americana* EYTON, Monogr. Anat. 1838, 155.

*Aythya americana* BAIRD, B. N. Am. 1858, 793.

[B 591, C 503, R 618, C 723.]

GEOG. DIST.—North America, breeding from California, southern Michigan, and Maine northward.

147. **Aythya vallisneria** (WILS.).

Canvas-back.

*Anas vallisneria* WILSON, Am. Orn. VIII. 1814, 103.

*Aythya vallisneria* BOIE, Isis, 1826, 980.

[B 592, C 504, R 617, C 724.]

GEOG. DIST.—Nearly all of North America, breeding from the northwestern States northward to Alaska.

SUBGENUS **FULIGULA** STEPHENS.

*Fuligula* STEPHENS, Gen. Zool. XII. pt. ii. 1824, 187. Type, by elimination, *Anas fuligula* LINN.

148. **Aythya marila nearctica** STEJN.

American Scaup Duck.

*Aythya marila nearctica* STEJN. Orn. Expl. Kamtsch. 1885, 161.

[B 588, C 500, R 614, C 720.]

GEOG. DIST.—North America, breeding far north. South in winter to Guatemala.

149. **Aythya affinis** (EYT.).

Lesser Scaup Duck.

*Fuligula affinis* EYT. Mon. Anat. 1838, 157.

*Aythya affinis* STEJN. Orn. Expl. Kamtsch. 1885, 161.

[B 589, C 501, R 615, C 721.]

GEOG. DIST.—North America in general, breeding chiefly north of the United States, migrating south to Guatemala and the West Indies.

150. **Aythya collaris** (DONOV.).

Ring-necked Duck.

*Anas collaris* DONOV. Br. Birds, VI. 1809, pl. 147.

*Aythya collaris* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 356.

[B 590, C 502, R 616, C 722.]

GEOG. DIST.—North America, breeding far north and migrating south to Guatemala and the West Indies.

GENUS **GLAUCIONETTA** STEJNEGER.

*Glaucionetta* STEJN. Pr. U. S. Nat. Mus. VIII. 1885, 409. Type,  
*Anas clangula* LINN.

151. **Glaucionetta clangula americana** (BONAP.).

American Golden-eye.

*Clangula americana* BONAP. Comp. List, 1838, 58.

*Glaucionetta clangula americana* STEJN. Pr. U. S. Nat. Mus.  
VIII. 1885, 409.

GEOG. DIST.—North America, breeding from Maine and the British Provinces northward; in winter, south to Cuba and Mexico.

152. **Glaucionetta islandica** (GMEL.).

Barrow's Golden-eye.

*Anas islandica* GMEL. S. N. I. 1788, 541.

*Glaucionetta islandica* STEJN. Pr. U. S. Nat. Mus. VIII. 1885,  
409.

[B 594, C 506, R 619, C 726.]

GEOG. DIST.—Northern North America, south in winter to New York, Illinois, and Utah; breeding from the Gulf of St. Lawrence northward, and south in the Rocky Mountains to Colorado; Greenland; Iceland. Accidental in Europe.

GENUS **CHARITONETTA** STEJNEGER.

*Charitonetta* STEJN. Orn. Expl. Kamtsch. 1885, 163. Type,  
*Anas albeola* LINN.

153. **Charitonetta albeola** (LINN.).

Buffle-head.

*Anas albeola* LINN. Syst. Nat. ed. 10, I. 1758, 124.

*Charitonetta albeola* STEJN. Orn. Expl. Kamtsch. 1885, 166.



[B 595, C 507, R 621, C 727.]

GEOG. DIST.—North America ; south in winter to Cuba and Mexico. Breeds from Maine and Montana northward, through the Fur Countries and Alaska.

GENUS **CLANGULA** LEACH.

*Clangula* LEACH, in Ross's Voy. Disc. 1819, App. p. xlviii. Type, *Anas glacialis* LINN.

154. **Clangula hyemalis** (LINN.).

Old-squaw.

*Anas hyemalis* LINN. S. N. ed. 10, I. 1758, 126.

*Clangula hiemalis* BREHM, Handb. Vög. Deutschl. 1831, 933.

[B 597, C 508, R 623, C 728.]

GEOG. DIST.—Northern hemisphere ; in North America south to the Potomac and the Ohio (more rarely to Florida and Texas) and California ; breeds far northward.

GENUS **HISTRIONICUS** LESSON.

*Histrionicus* LESSON, Man. d'Orn. II. 1828, 415. Type, *Anas histrionica* LINN.

155. **Histrionicus histrionicus** (LINN.).

Harlequin Duck.

*Anas histrionica* LINN. S. N. ed. 10, I. 1758, 127.

*Histrionicus histrionicus* BOUCARD, Cat. Av. 1876, 60.

[B 596, C 510, R 622, C 730.]

GEOG. DIST.—Northern North America, breeding from Newfoundland, the northern Rocky Mountains, and the Sierra Nevada (lat. 38°), northward ; south in winter to the Middle States and California ; Eastern Asia ; Iceland.

GENUS **CAMPTOLAIMUS** GRAY.

*Camptolaimus* GRAY, List Gen. 1841, 95. Type, *Anas labradoria* GMEL.

156. **Camptolaimus labradorius** (GMEL.).

Labrador Duck.

*Anas labradoria* GMEL. S. N. I. 1788, 537.

*Camptolaimus labradorus* GRAY, List Gen. 1841, 95.

[B 600, C 510, R 624, C 730.]

GEOG. DIST.—Formerly Northern Atlantic coast, from New Jersey (in winter) northward, breeding from Labrador northward. Now extinct.

GENUS **ENICONETTA** GRAY.

*Eniconetta* GRAY, List Gen. 1840, 75. Type, *Anas stelleri* PALL.

157. **Eniconetta stelleri** (PALL.).

Steller's Duck.

*Anas stelleri* PALL. Spicil. Zool. VI. 1769, 35.

*Eniconetta stelleri* GRAY, List Gen. 1840, 75.

[B 598, C 511, R 625, C 731.]

GEOG. DIST.—Arctic and subarctic coasts of the northern hemisphere. Aleutian Islands, east to Unalaska and Kadiak; Kenai Peninsula.

GENUS **ARCTONETTA** GRAY.

*Arctonetta* GRAY, P. Z. S. 1855 (Feb. 1856), 212. Type, *Fuligula fischeri* BRANDT.

158. **Arctonetta fischeri** (BRANDT).

Spectacled Eider.

*Fuligula fischeri* BRANDT, Mém. Acad. St. Pétersb. VI. 1849, 6, 10.

*Arctonetta fischeri* BLAKISTON, Ibis, 1863, 150.

[B 599, C 512, R 626, C 732.]

GEOG. DIST.—Alaskan coast of Bering Sea and north to Point Barrow.

GENUS **SOMATERIA** LEACH.

SUBGENUS **SOMATERIA**.

*Somateria* LEACH, in ROSS'S Voy. Disc. 1819, App. p. xlviii.

Type, *Anas mollissima* LINN.

159. **Somateria mollissima borealis** A. E. BREHM.

Northern Eider.

*Somateria mollissima borealis* A. E. BREHM, Verz. Samml. Eur. Vög. 1866, 14.

[B 606, *part*, C 513, *part*, R 627, C 733.]

GEOG. DIST.—Northeastern North America, including Greenland; south, in winter, to coast of Massachusetts.

160. **Somateria dresseri** SHARPE.

American Eider.

*Somateria dresseri* SHARPE, Ann. Mag. Nat. Hist. July, 1871, 51.

[B 606, *part*, C 513, *part*, R 627a, C 734.]

GEOG. DIST.—Atlantic coast of North America, from Maine to Labrador; south in winter to the Delaware and west to the Great Lakes.

161. **Somateria v-nigra** GRAY.

Pacific Eider.

*Somateria v-nigra* GRAY, P. Z. S. 1855 (Feb. 1856), 212, pl. cvii.

[B 607, C 514, R 628, C 735.]

GEOG. DIST.—Coasts of the North Pacific; in the interior to the Great Slave Lake district; eastern Siberia.

SUBGENUS **ERIONETTA** COUES.

*Erionetta* COUES, Key N. A. Birds, ed. 2, 1884, 709. Type,

*Anas spectabilis* LINN.

162. **Somateria spectabilis** (LINN.).

King Eider.

*Anas spectabilis* LINN. S. N. ed. 10, I. 1758, 123.*Somateria spectabilis* LEACH, in ROSS'S Voy. Disc. 1819, App. p. xlviii.

[B 608, C 515, R 629, C 736.]

GEOG. DIST.—Northern parts of northern hemisphere, breeding in the Arctic regions; in North America south casually in winter to Georgia and the Great Lakes.

GENUS **OIDEMIA** FLEMING.SUBGENUS **OIDEMIA**.

*Oidemia* FLEMING, Philos. Zool. II. 1822, 260. Type, by elimination, *Anas nigra* LINN.

163. **Oidemia americana** SW. & RICH.

American Scoter.

*Oidemia americana* SW. & RICH. Faun. Bor. Amer. II. 1831, 450.

[B 604, C 516, R 630, C 737.]

GEOG. DIST.—Coasts and larger inland waters of northern North America; breeds in Labrador and the northern interior; south in winter to New Jersey, the Great Lakes, Colorado, and California.

SUBGENUS **MELANITTA** BOIE.

*Melanitta* BOIE, Isis, 1822, 564. Type, by elimination, *Anas fusca* LINN.

[164.] **Oidemia fusca** (LINN.).

Velvet Scoter.

*Anas fusca* LINN. S. N. ed. 10, I. 1758, 123.*Oidemia fusca* STEPHENS, Gen. Zool. XII. pt. ii. 1824, 216.

[B—, C—, R 631, C—.]

GEOG. DIST.—Northern Old World; accidental (?) in Greenland.

165. **Oidemia deglandi** BONAP.

White-winged Scoter.

*Oidemia deglandi* BONAP. Rev. Crit. de l'Orn. Europ. de Dr. Degl., 1850, 108.

[B 601, C 517, R 632, C 738.]

GEOG. DIST.—Northern North America, breeding in Labrador and the Fur Countries; south in winter to Chesapeake Bay, Southern Illinois, and San Quentin Bay, Lower California.

SUBGENUS **PELIONETTA** KAUP.

*Pelionetta* KAUP, Sk. Ent. Eur. Thierw. 1829, 107. Type, *Anas perspicillata* LINN.

166. **Oidemia perspicillata** (LINN.).

Surf Scoter.

*Anas perspicillata* LINN. S. N. ed. 10, I. 1758, 125.

*Oidemia perspicillata* STEPHENS, Gen. Zool. XII. pt. ii. 1824, 219.

[B 602, 603, C 518, 518a, R 633, C 739, 740.]

GEOG. DIST.—Coasts and larger inland waters of northern North America; in winter south to Florida, the Ohio River, and San Quentin Bay, Lower California. Accidental in Europe.

GENUS **ERISMATURA** BONAPARTE.

*Erismatura* BONAP. Saggio Distr. Meth. 1832, 143. Type, *Anas rubidus* WILS.

167. **Erismatura rubida** (WILS.).

Ruddy Duck.

*Anas rubidus* WILSON, Am. Orn. VIII. 1814, 128.

*Erismatura rubida* BONAP. Comp. List, 1838, 59.

[B 609, C 519, R 634, C 741.]

GEOG. DIST.—North America in general, south to the West Indies and through Central America to Colombia; breeds throughout much of its North American range and south to Guatemala.



GENUS **NOMONYX** RIDGWAY.

*Nomonyx* RIDGW. Pr. U. S. Nat. Mus. III. 1880, 15. Type, *Anas dominica* LINN.

[168.] **Nomonyx dominicus** (LINN.).

Masked Duck.

*Anas dominica* LINN. S. N. ed. 12, 1766, 201.

*Nomonyx dominicus* RIDGW. Pr. U. S. Nat. Mus. III. 1880, 15.

[B 610, C 520, R 635, C 742.]

GEOG. DIST.—Tropical America in general, including West Indies, north on the Gulf coast to the Lower Rio Grande; accidental in eastern North America (Wisconsin; Lake Champlain; Massachusetts).

SUBFAMILY **ANSERINÆ**. GEESE.GENUS **CHEN** BOIE.

*Chen* BOIE, Isis, 1822, 563. Type, *Anser hyperboreus* PALL.

169. **Chen hyperborea** (PALL.).

Lesser Snow Goose.

*Anser hyperboreus* PALL. Spicil. Zool. VI. 1769, 25.

*Chen hyperborea* BOIE, Isis, 1822, 563.

[B —, C 480a, R 591a, C 696.]

GEOG. DIST.—Pacific coast to the Mississippi Valley, breeding in Alaska; south in winter to southern Illinois and southern California; casually to New England. Northeastern Asia.

169 a. **Chen hyperborea nivalis** (FORST.).

Greater Snow Goose.

*Anas nivalis* FORSTER, Philos. Trans. LXII. 1772, 413.

*Chen hyperborea nivalis* RIDGW. Pr. Biol. Soc. Wash. II. 1884, 107.

[B 563, C 480, R 591, C 695.]

GEOG. DIST.—North America, breeding far north (east of Mackenzie basin) and migrating south in winter, chiefly along the Atlantic coast, reaching Cuba.

169.1. **Chen cærulescens** (LINN.).

Blue Goose.

*Anas cærulescens* LINN. S. N. ed. 10, I. 1758, 124.*Chen cærulescens* GUNDL. in POEY's Rept. Fis.-nat. Isla Cuba, I, 1865-66, 387.

[B 564, C 479, R 590, C 694.]

GEOG. DIST.—Interior of North America, breeding on eastern shores of Hudson Bay and migrating south, in winter, through Mississippi Valley to Gulf coast; occasional on Atlantic coast.

170. **Chen rossii** (CASSIN).

Ross's Snow Goose.

*Anser rossii* "BAIRD MSS.," CASS. Pr. Ac. Nat. Sci. Phila. 1861, 73.*Chen rossii* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880, 203.

[B —, C 481, R 592, C 697.]

GEOG. DIST.—Arctic America in summer, Pacific coast to southern California and east to Montana in winter.

GENUS **ANSER** BRISSON.*Anser* BRISSON, Orn. VI. 1760, 261. Type, *Anas anser* LINN.[171.] **Anser albifrons** (GMEL.).

White-fronted Goose.

*Anas albifrons* GMEL. S. N. I. 1788, 509.*Anser albifrons* BECHST. Gem. Naturg. Deutschl. IV. 1809, 898.

[B —, C —, R 593, C 692.]

GEOG. DIST.—Northern parts of eastern hemisphere. Eastern Greenland?

171 a. **Anser albifrons gambeli** (HARTL.).

American White-fronted Goose.

*Anser gambeli* HARTLAUB, Rev. Mag. Zool. 1852, 7.*Anser albifrons* var. *gambeli* COUES, Key, 1872, 282.

[B 565, 566, C 478, R 593*a*, C 693.]

GEOG. DIST.—North America (rare on the Atlantic coast), breeding far northward; in winter south to Cape St. Lucas, Mexico, and Cuba.

GENUS **BRANTA** SCOPOLI.

*Branta* SCOPOLI, Ann. I. Hist. Nat. 1769, 67. Type, *Anas bernicla* LINN.

172. ***Branta canadensis*** (LINN.).

Canada Goose.

*Anas canadensis* LINN. S. N. ed. 10, I. 1758, 123.

*Branta canadensis* BANNISTER, Pr. Ac. Nat. Sci. Phila. 1870, 131.

[B 567, C 485, R 594, C 702.]

GEOG. DIST.—Temperate North America, breeding in the northern United States and British Provinces; south in winter to Mexico.

172 *a*. ***Branta canadensis hutchinsii*** (RICH.).

Hutchins's Goose.

*Anser hutchinsii* RICH. in Sw. & RICH. Faun. Bor. Am. II. 1831, 470.

*Branta canadensis* var. *hutchinsii* COUES, Key, 1872, 284.

[B 569, C 485*b*, R 594*a*, C 704.]

GEOG. DIST.—North America, breeding in the Arctic regions, and migrating south in winter, chiefly through the western United States and Mississippi Valley; northeastern Asia.

172 *b*. ***Branta canadensis occidentalis*** (BAIRD).

White-cheeked Goose.

*Bernicla occidentalis* BAIRD, B. N. Am. 1858, 766.

*Branta canadensis occidentalis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 567*a*, C —, R 594*c*, C —.]

GEOG. DIST.—Pacific coast region, from Sitka south, in winter, to California.

172 c. **Branta canadensis minima** RIDGW.

Cackling Goose.

*Branta minima* RIDGW. Pr. U. S. Nat. Mus. VIII. No. 2, April 20, 1885, 23.*Branta canadensis minima* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 568, C 485a, R 594b, C 703.]

GEOG. DIST.—Coast of Alaska, chiefly about Norton Sound and Lower Yukon, migrating southward into the western United States, east to Wisconsin.

173. **Branta bernicla** (LINN.).

Brant.

*Anas bernicla* LINN. S. N. ed. 10, I. 1758, 124.*Branta bernicla* SCOPOLI, Ann. I. Hist. Nat. 1769, 67.

[B 570, C 484, R 595, C 700.]

GEOG. DIST.—Northern parts of the northern hemisphere; in North America chiefly on the Atlantic coast; rare in the interior, or away from salt water.

174. **Branta nigricans** (LAWR.).

Black Brant.

*Anser nigricans* LAWRENCE, Ann. Lyc. N. Y. IV. 1846, 171.*Branta nigricans* BANNISTER, Pr. Ac. Nat. Sci. Phila. 1870, 131.

[B 571, C —, R 596, C 701.]

GEOG. DIST.—Arctic and western North America, south in winter to Lower California; casual in the Atlantic States.

[175.] **Branta leucopsis** (BECHST.).

Barnacle Goose.

*Anas leucopsis* BECHSTEIN, Orn. Taschb. Deutschl. 1803, 424.*Branta leucopsis* BANNISTER, Pr. Ac. Nat. Sci. Phila. 1870, 131.

[B 572, C 483, R 597, C 699.]

GEOG. DIST.—Northern parts of the Old World; casual in eastern North America.

GENUS **PHILACTE** BANNISTER.

*Philacte* BANNISTER, Pr. Ac. Nat. Sci. Phila. 1870, 131. Type,  
*Anas canagica* SEVAST.

176. **Philacte canagica** (SEVAST.).

Emperor Goose.

*Anas canagica* SEVASTIANOFF, N. Act. Petrop. XIII. 1800, 346.

*Philacte canagica* BANNISTER, Pr. Ac. Nat. Sci. Phila. 1870, 131.

[B 573, C 482, R 598, C 698.]

GEOG. DIST.—Coast and islands of Alaska north of the Peninsula; chiefly about Norton Sound and valley of the Lower Yukon; Commander Islands, Kamchatka; casually southward to Humboldt Bay, California.

GENUS **DENDROCYGNA** SWAINSON.

*Dendrocygna* SWAINSON, Classif. Birds, II. 1837, 365. Type,  
*Anas arcuata* CUV.

177. **Dendrocygna autumnalis** (LINN.).

Black-bellied Tree-duck.

*Anas autumnalis* LINN. S. N. ed. 10, I. 1758, 127.

*Dendrocygna autumnalis* EYTON, Monogr. Anat. 1838, 109.

[B 574, C 487, R 599, C 706.]

GEOG. DIST.—Southwestern border of the United States and southward (Mexico, West Indies, etc.).

178. **Dendrocygna fulva** (GMEL.).

Fulvous Tree-duck.

*Anas fulva* GMEL. S. N. I. 1788, 530.

*Dendrocygna fulva* BURMEISTER, Reise durch die La Plata Staaten, 1856, 515.



[B 575, C 486, R 600, C 705.]

GEOG. DIST.—Southern border of the United States (Louisiana, Texas, Nevada, California), south to Mexico, reappearing in southern Brazil and Argentine Republic. Casual in North Carolina and Missouri.

SUBFAMILY **CYGNINÆ**. SWANS.GENUS **OLOR** WAGLER.

*Olor* WAGLER, Isis, 1832, 1234. Type, *Anas cygnus* LINN.

[179.] **Olor cygnus** (LINN.).**Whooping Swan.**

*Anas cygnus* LINN. S. N. ed. 10, I. 1758, 122.

*Olor cygnus* BONAPARTE, Catal. Parzudaki, 1856, 15.

[B—, C—, R 586, C 690.]

GEOG. DIST.—Europe and Asia; occasional in southern Greenland.

180. **Olor columbianus** (ORD).**Whistling Swan.**

*Anas columbianus* ORD, in GUTHRIE'S Geogr. 2d Am. ed. 1815, 319.

*Olor columbianus* STEJN. Pr. U. S. Nat. Mus. V. 1882, 210.

[B 561 *bis*, C 477, R 588, C 689.]

GEOG. DIST.—The whole of North America, breeding far north. Commander Islands, Kamchatka; accidental in Scotland.

181. **Olor buccinator** (RICH.).**Trumpeter Swan.**

*Cygnus buccinator* RICH. in Sw. & RICH. Fauna Bor. Am. II. 1831, 464.

*Olor buccinator* WAGLER, Isis, 1832, 1234.

[B 562, C 476, R 589, C 688.]

GEOG. DIST.—Chiefly the interior of North America, from the Gulf coast to the Fur Countries, breeding from Iowa and the Dakotas northward; west to the Pacific coast; rare or casual on the Atlantic.

ORDER ODONTOGLOSSÆ. LAMELLIROSTRAL GRALLATORES.

FAMILY **PHŒNICOPTERIDÆ**. FLAMINGOES.

GENUS **PHŒNICOPTERUS** LINN.

*Phœnicopterus* LINN. S. N. ed. 10, I. 1758, 139. Type, *P. ruber* LINN.

182. **Phœnicopterus ruber** LINN.  
American Flamingo.

*Phœnicopterus ruber* LINN. S. N. ed. 10, I. 1758, 139.

[B 502, C 475, R 585, C 687.]

GEOG. DIST.—Atlantic coasts of subtropical and tropical America ; southern Florida. Galapagos.

ORDER HERODIONES. HERONS, STORKS,  
IBISES, ETC.

SUBORDER IBIDES. SPOONBILLS AND IBISES.

FAMILY **PLATALEIDÆ**. SPOONBILLS.

GENUS **AJAJA** REICH.

*Ajaja* REICH. Handb. 1852, p. xvi. Type, *Platalea ajaja* LINN.

183. **Ajaja ajaja** (LINN.).

Roseate Spoonbill.

*Platalea ajaja* LINN. S. N. ed. 10, I. 1758, 140.*Ajaja ajaja* BOUCARD, Cat. Av. 1876, 54.

[B 501, C 488, R 505, C 653.]

GEOG. DIST.—Southern Atlantic and Gulf States southward to the Falkland Islands and Patagonia. Formerly north to southern Illinois.

FAMILY **IBIDIDÆ**. IBISES.GENUS **GUARA** REICHENBACH.

*Guara* REICH. Syst. Av. 1852, p. xiv. Type, *Tantalus ruber* LINN.

184. **Guara alba** (LINN.).

White Ibis.

*Scolopax alba*, LINN. S. N. ed. 10, I. 1758, 145.*Guara alba*, STEJN, Stand. Nat. Hist. IV. 1885, 9.

[B 499, C 446, R 501, C 651.]

GEOG. DIST.—South Atlantic and Gulf States southward to the West Indies and northern South America; north to North Carolina, southern Illinois, Great Salt Lake, and Lower California; casually to Long Island, Connecticut, and South Dakota.

[185.] **Guara rubra** (LINN.).

Scarlet Ibis.

*Tantalus ruber* LINN. S. N. ed. 12, I. 1766, 241.*Guara rubra* REICH. Syst. Av. 1852, p. xiv.

[B 498, C 447, R 502, C 652.]

GEOG. DIST.—Florida, Louisiana, and Texas, southward to the West Indies and northern South America. No record of its recent occurrence in the United States.

GENUS **PLEGADIS** KAUP.

*Plegadis* KAUP, Skizz. Entw. Gesch. Eur. Thierw. 1829, 82.

Type, *Tantalus falcinellus* LINN. = *Tringa autumnalis* HASSELQ.

186. **Plegadis autumnalis** (HASSELQ.).

Glossy Ibis.

*Tringa autumnalis* HASSELQUIST, Reise nach Paläst. Deutsche Aug. 1762, 306.

*Plegadis autumnalis* STEJN. Stand. Nat. Hist. IV. 1885, 160.

[B 500, C 445, R 503, C 649.]

GEOG. DIST.—Warmer parts of eastern hemisphere, West Indies and southern portions of eastern United States, wandering northward to New England and Illinois. In America only locally abundant and of irregular distribution.

187. **Plegadis guarauna** (LINN.).

White-faced Glossy Ibis.

*Scolopax guarauna* LINN. S. N. ed. 12, I. 1766, 242.

*Plegadis guarauna* RIDGW. Pr. U. S. Nat. Mus. I. Oct. 2, 1878, 163.

[B —, C 445 *bis*, 445 *ter*, R 504, C 650.]

GEOG. DIST.—Western United States, from Texas to California, Oregon, and casually to southern British Columbia, Kansas and Florida (breeding); southward through the West Indies and Mexico to South America.

## SUBORDER CICONIÆ. STORKS, ETC.

FAMILY **CICONIIDÆ**. STORKS AND WOOD IBISES.

SUBFAMILY **TANTALINÆ**. WOOD IBISES.

GENUS **TANTALUS** LINNÆUS.

*Tantalus* LINN. S. N. ed. 10, I. 1758, 140. Type, *T. loculator* LINN.

188. **Tantalus loculator** LINN.

Wood Ibis.

*Tantalus loculator* LINN. S. N. ed. 10, I. 1758, 140.

[B 497, C 444, R 500, C 648.]

GEOG. DIST.—Southern United States, from the Ohio Valley, Colorado, Utah, southeastern California, etc., south to Argentine Republic; casually northward to Pennsylvania and New York.

SUBFAMILY **CICONIINÆ**. STORKS.GENUS **MYCTERIA** LINNÆUS.

*Mycteria* LINN. S. N. ed. 10, I. 1758, 140. Type, *M. americana* LINN.

[189.] **Mycteria americana** LINN.

Jabiru.

*Mycteria americana* LINN. S. N. ed. 10, I. 1758, 140.[B—, C 448 *bis*, R 499, C 654.]

GEOG. DIST.—Tropical America, north casually to southern Texas.

SUBORDER HERODII. HERONS, EGRETS,  
BITTERNS, ETC.

FAMILY **ARDEIDÆ**. HERONS, BITTERNS, ETC.SUBFAMILY **BOTAURINÆ**. BITTERNS.GENUS **BOTAURUS** HERMANN.

*Botaurus* HERMANN, Tabl. Affin. Anim. 1783, 135. Type,  
*Ardea stellaris* LINN.

190. ***Botaurus lentiginosus*** (MONTAG.).**American Bittern.***Ardea lentiginosa* MONTAG. Orn. Dict. Suppl. 1813, —.*Botaurus lentiginosus* STEPH. Gen. Zool. XI. ii. 1819, 592.

[B 492, C 460, R 497, C 666.]

GEOG. DIST.—Temperate North America, south to Guatemala, Cuba, Jamaica, and Bermuda ; occasional in British Islands.

GENUS **ARDETTA** GRAY.

*Ardetta* GRAY, List Gen. B. App. 1842, 13. Type, *Ardea minuta* LINN.

191. ***Ardetta exilis*** (GMEL.).**Least Bittern.***Ardea exilis* GMEL. S. N. I. ii. 1788, 645.*Ardetta exilis* GUNDL. J. f. O. 1856, 345.

[B 491, C 461, R 498, C 667.]

GEOG. DIST.—Temperate North America, north to the British Provinces and south to the West Indies and Brazil. Less common west of the Rocky Mountains ; on the Pacific coast north to northern California.

191.1. ***Ardetta neoxena*** CORY.**Cory's Least Bittern.***Ardetta neoxena* CORY, Auk, III. April, 1886, 262.

[B —, C —, R —, C —.]

GEOG. DIST.—Southern Florida (Caloosahatchie River, near Lake Okeechobee) ; Ontario ; Michigan.

SUBFAMILY **ARDEINÆ**. HERONS AND EGRETS.GENUS **ARDEA** LINN.SUBGENUS **ARDEA**.

*Ardea* LINN. S. N. ed. 10, I. 1758, 141. Type, by elimination, *A. cinerea* LINN.



192. **Ardea occidentalis** AUD.

Great White Heron.

*Ardea occidentalis* AUD. Orn. Biog. III. 1835, 542.[B 489, C 451, R 486, C 656, *part.*]

GEOG. DIST.—Florida; Cuba; Jamaica.

193. **Ardea wardi** RIDGW.

Ward's Heron.

*Ardea wardi* RIDGW. Bull. Nutt. Orn. Club, VII. Jan. 1882, 5.

[B —, C —, R —, C —.]

GEOG. DIST.—Florida.

194. **Ardea herodias** LINN.

Great Blue Heron.

*Ardea herodias* LINN. S. N. ed. 10, I. 1758, 143.

[B 487, C 449, R 487, C 655.]

GEOG. DIST.—North America, from the Arctic regions southward to the West Indies and northern South America. Bermudas; Galapagos.

[195.] **Ardea cinerea** LINN.

European Blue Heron.

*Ardea cinerea* LINN. S. N. ed. 10, I. 1758, 143.

[B —, C —, R 488, C 657.]

GEOG. DIST.—Most of the eastern hemisphere; accidental in southern Greenland.

SUBGENUS **HERODIAS** BOIE.*Herodias* BOIE, Isis, 1822, 559. Type, by elimination, *Ardea egretta* GMEL.196. **Ardea egretta** GMEL.

American Egret.

*Ardea egretta* GMEL. S. N. I. ii. 1788, 629.

[B 486, 486\*, C 452, R 489, C 658.]

GEOG. DIST.—Temperate and tropical America, from New Jersey, Minnesota, and Oregon south to Patagonia; casually on the Atlantic coast to Nova Scotia.

SUBGENUS **GARZETTA** KAUP.

*Garzetta* KAUP, Skizz. Entw. Gesch. Eur. Thierw. 1829, 76.

Type, *Ardea garzetta* LINN.

197. ***Ardea candidissima*** GMEL.

Snowy Heron.

*Ardea candidissima* GMEL. S. N. I. ii. 1788, 633.

[B 485, C 453, R 490, C 659.]

GEOG. DIST.—Temperate and tropical America, from Long Island and Oregon south to Argentine Republic and Chili; casually to Nova Scotia and southern British Columbia.

SUBGENUS **DICHROMANASSA** RIDGWAY.

*Dichromanassa* RIDGW. Bull. U. S. Geol. & Geog. Surv. Terr. IV.

Feb. 5, 1878, 246. Type, *Ardea rufa* BODD.

198. ***Ardea rufescens*** GMEL.

Reddish Egret.

*Ardea rufescens* GMEL. S. N. I. ii. 1788, 628.

[B 483, 482, C 455, R 491, C 661.]

GEOG. DIST.—Gulf States, north to southern Illinois; Lower California and Mexico (both coasts), south to Guatemala; Jamaica, and Cuba.

SUBGENUS **HYDRANASSA** BAIRD.

*Hydranassa* BAIRD, B. N. Am. 1858, 660 (in text). Type, *Ardea*

*ludoviciana* WILS. = *Egretta ruficollis* GOSSE.

199. ***Ardea tricolor ruficollis*** (GOSSE).

Louisiana Heron.

*Egretta ruficollis* GOSSE, B. Jamaica, 1847, 338.

*Ardea tricolor ruficollis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885,

355.

[B 484, C 454, R 492, C 660.]

GEOG. DIST.—Gulf States, Mexico (both coasts), Central America, and West Indies ; casually northward to New Jersey and Indiana.

SUBGENUS **FLORIDA** BAIRD.

*Florida* BAIRD, B. N. Am. 1858, 671. Type, *Ardea cærulea* LINN.

200. **Ardea cærulea** LINN.

Little Blue Heron.

*Ardea cærulea* LINN. S. N. ed. 10, I. 1758, 143.

[B 490, C 456, R 493, C 662.]

GEOG. DIST.—Eastern United States, from New Jersey, Illinois, and Kansas, southward through Central America and the West Indies to Guiana and Colombia ; casually north on the Atlantic coast to Massachusetts and Maine.

SUBGENUS **BUTORIDES** BLYTH.

*Butorides* BLYTH, Cat. B. Mus. As. Soc. 1849, 281. Type, *Ardea javanica* HORSE.

201. **Ardea virescens** LINN.

Green Heron.

*Ardea virescens* LINN. S. N. ed. 10, I. 1758, 144.

[B 493, C 457, R 494, C 663.]

GEOG. DIST.—Temperate North America, from Ontario and Oregon, southward to Colombia, Venezuela, and the West Indies. Bermuda.

201 a. **Ardea virescens frazari** BREWST.

Frazar's Green Heron.

*Ardea virescens frazari* BREWST. Auk, V. Jan. 1888, 83.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California (vicinity of La Paz).

GENUS **NYCTICORAX** STEPHENS.SUBGENUS **NYCTICORAX**.

*Nycticorax* STEPH. Gen. Zool. XI. ii. 1819, 608. Type, *Ardea nycticorax* LINN.

202. **Nycticorax nycticorax nævius** (BODD.).

Black-crowned Night Heron.

*Ardea nævia* BODD. Tabl. Pl. Enl. 1783, 56.

*Nycticorax nycticorax nævius* ZELEDON, Pr. U. S. Nat. Mus. VIII. 1885, 113.

[B 495, C 458, R 495, C 664.]

GEOG. DIST.—America, from Ontario and Manitoba southward to the Falkland Islands, including part of the West Indies.

SUBGENUS **NYCTANASSA** STEJNEGER.

*Nyctanassa* STEJN. Proc. U. S. Nat. Mus. X. 1887, 295. Type, *Ardea violacea* LINN.

203. **Nycticorax violaceus** (LINN.).

Yellow-crowned Night Heron.

*Ardea violacea* LINN. S. N. ed. 10, I. 1758, 143.

*Nycticorax violacea* VIGORS, Zool. Journ. III. 1827, 446.

[B 496, C 459, R 496, C 665.]

GEOG. DIST.—Tropical and warm temperate North America, from the Carolinas, the lower Ohio Valley and Lower California, south to Brazil; casually north to Massachusetts and Colorado.

## ORDER PALUDICOLÆ. CRANES, RAILS, ETC.

## SUBORDER GRUES. CRANES.

FAMILY **GRUIDÆ**. CRANES.GENUS **GRUS** PALLAS.

*Grus* PALL. Misc. Zool. 1766, 66. Type, *Ardea grus* LINN.

204. **Grus americana** (LINN.).

Whooping Crane.

*Ardea americana* LINN. S. N. ed. 10, I. 1758, 142.

*Grus americana* VIEILL. Nouv. Dict. d'Hist. Nat. XIII. 1817, 557.

[B 478, C 462, R 582, C 668.]

GEOG. DIST.—Interior of North America, from the Fur Countries to Florida, Texas, and Mexico, and from Ohio to Colorado. Formerly on the Atlantic coast, at least casually, to New England.

205. **Grus canadensis** (LINN.).

Little Brown Crane.

*Ardea canadensis* LINN. S. N. ed. 10, I. 1758, 141.

*Grus canadensis* TEMM. Man. I. 1820, p. c.

[B 480, C 463, R 584, C 669.]

GEOG. DIST.—Arctic and subarctic America, breeding from the Fur Countries and Alaska to the Arctic coast, migrating south in winter into the western United States.

206. **Grus mexicana** (MÜLL.).

Sandhill Crane.

*Ardea (grus) mexicana* MÜLL. S. N. Suppl. 1776, 110.

*Grus mexicana* VIEILL. Nouv. Dict. d'Hist. Nat. XIII. 1817, 561.

[B 479, C —, R 583, C 670.]

GEOG. DIST.—Southern half of North America; now rare near the Atlantic coast, except in Georgia and Florida.

## SUBORDER RALLI. RAILS, GALLINULES, COOTS, ETC.

FAMILY **ARAMIDÆ**. COURLANS.

GENUS **ARAMUS** VIEILLOT.

*Aramus* VIEILL. Analyse, 1816, 58. Type, *Ardea scolopacea* GMEL.

### 207. **Aramus giganteus** (BONAP.).

**Limpkin.**

*Rallus giganteus* BONAP. Jour. Ac. Nat. Sci. Phila. V. 1825, 31.

*Aramus giganteus* BAIRD, B. N. Am. 1858, 657.

[B 481, C 464, R 581, C 671.]

GEOG. DIST.—Florida, Greater Antilles, and both coasts of Central America.

FAMILY **RALLIDÆ**. RAILS, GALLINULES, AND COOTS.

SUBFAMILY **RALLINÆ**. RAILS.

GENUS **RALLUS** LINNÆUS.

*Rallus* LINN. S. N. ed. 10, I. 1758, 153. Type, *R. aquaticus* LINN.

### 208. **Rallus elegans** AUD.

**King Rail.**

*Rallus elegans* AUD. Orn. Biog. III. 1835, 27, pl. 203.



[B 552, C 466, R 569, C 676.]

GEOG. DIST.—Fresh-water marshes of the eastern United States, north to the Middle States, northern Illinois, Wisconsin, and Kansas, casually to Massachusetts, Maine, and Ontario.

209. **Rallus beldingi** RIDGW.

**Belding's Rail.**

*Rallus beldingi* RIDGW. Pr. U. S. Nat. Mus. V. 1882, 345.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California (Espiritu Santo Island and vicinity of La Paz).

210. **Rallus obsoletus** RIDGW.

**California Clapper Rail.**

*Rallus elegans* var. *obsoletus* RIDGW. Am. Nat. VIII. 1871, 111.

*Rallus obsoletus* RIDGW. Bull. Nutt. Orn. Club, V. July, 1880, 139.

[B —, C 466a, R 570, C 674.]

GEOG. DIST.—Salt marshes of the Pacific coast, from Washington (?) to Lower California.

211. **Rallus crepitans** GMEL.

**Clapper Rail.**

*Rallus crepitans* GMEL. S. N. I. ii. 1788, 713.

[B 553, C 465, R 571, C 673.]

GEOG. DIST.—Salt marshes of the Atlantic and Gulf coasts of the United States, breeding from southern Connecticut southward; resident from the Potomac southward, occasionally wintering further north. Casual north to Massachusetts.

211 a. **Rallus crepitans saturatus** RIDGW.

**Louisiana Clapper Rail.**

*Rallus longirostris saturatus* RIDGW. Bull. Nutt. Orn. Club, V. July, 1880, 140.

*Rallus crepitans saturatus* SENNETT, Auk, VI. April, 1889, pp. 164, 166.

[B—, C—, R 571a, C 675.]

GEOG. DIST.—Coast of Louisiana.

211.1. **Rallus scottii** (SENN.).

**Florida Clapper Rail.**

*Rallus longirostris scottii* SENN. Auk, V. July, 1888, 305.

*Rallus scottii* SENNETT, Auk, VI. April, 1889, pp. 165, 166.

[B—, C—, R—, C—.]

GEOG. DIST.—Western Florida (salt marshes).

[211.2.] **Rallus longirostris caribæus** RIDGW.

**Caribbean Clapper Rail.**

*Rallus longirostris caribæus* RIDGW. Bull. Nutt. Orn. Club, V. July, 1880, 140.

[B—, C—, R—, C—.]

GEOG. DIST.—West Indies and Gulf coast of Mexico, north to Texas (Corpus Christi and Galveston).

212. **Rallus virginianus** LINN.

**Virginia Rail.**

*Rallus virginianus* LINN. S. N. ed. 12, I. 1766, 263.

[B 554, C 467, R 572, C 677.]

GEOG. DIST.—North America, from the British Provinces south to Guatemala and Cuba.

GENUS **PORZANA** VIEILLOT.

SUBGENUS **PORZANA.**

*Porzana* VIEILL. Analyse, 1816, 61. Type, *Rallus porzana* LINN.

[213.] **Porzana porzana** (LINN.).

**Spotted Crane.**

*Rallus porzana* LINN. S. N. ed. 12, I. 1766, 262.

*Porzana porzana* BOUCARD, Cat. Av. 1876, 7.

[B—, C—, R 573, C 678.]

GEOG. DIST.—Northern parts of the Old World; occasional in Greenland.

214. **Porzana carolina** (LINN.).

**Sora.**

*Rallus carolinus* LINN. S. N. ed. 10, I. 1758, 153.

*Porzana carolina* BAIRD, Lit. Rec. & Jour. Linn. Assoc. Penn. Coll. Oct. 1845, 255.

[B 555, C 648, R 574, C 679.]

GEOG. DIST.—Temperate North America, breeding chiefly northward, but less commonly on the Pacific coast. Casually north to South Greenland. South to the West Indies and northern South America.

SUBGENUS **COTURNICOPS** BONAPARTE.

*Coturnicops* BONAP. Compt. Rend. XLIII. 1856, 599. Type, *Fulica novaboracensis* GMEL.

215. **Porzana noveboracensis** (GMEL.).

**Yellow Rail.**

*Fulica noveboracensis* GMEL. S. N. I. ii. 1788, 701.

*Porzana noveboracensis* BAIRD, Lit. Rec. & Jour. Linn. Assoc. Penn. Coll. Oct. 1845, 255.

[B 557, C 469, R 575, C 680.]

GEOG. DIST.—Chiefly eastern North America, north to Nova Scotia, Hudson Bay, etc., less commonly west to Nevada and California. No extralimital records except for Cuba and Bermuda.

SUBGENUS **CRECISCUS** CABANIS.

*Creciscus* CAB. J. f. O. 1856, 428. Type, *Rallus jamaicensis* LINN.

216. **Porzana jamaicensis** (GMEL.).

**Black Rail.**

*Rallus jamaicensis* GMEL. S. N. I. ii. 1788, 718.

*Porzana jamaicensis* BAIRD, Lit. Rec. & Jour. Linn. Assoc. Penn. Coll. Oct. 1845, 257.

[B 556, C 470, R 576, C 681.]

GEOG. DIST.— Temperate North America, north to Massachusetts, northern Illinois, and Oregon : south to West Indies and Guatemala.

216.1. **Porzana coturniculus** (RIDGW.).

Farallone Rail.

*Porzana jamaicensis* var. *coturniculus* "BAIRD, MS." RIDGW. Am. Nat. VIII. Feb. 1874, 111.

*Porzana coturniculus* RIDGW. Proc. U. S. Nat. Mus. XIII. 1890, 311.

[B —, C 470a, R 576a, C 682.]

GEOG. DIST.— Known only from the type specimen, from Farallone Islands, California.

GENUS **CREX** BECHSTEIN.

*Crex* BECHST. Orn. Taschb. Deutschl. 1802, 336. Type, *Rallus crex* LINN.

[217.] **Crex crex** (LINN.).

Corn Crane.

*Rallus crex* LINN. S. N. ed. 10, I. 1758, 153.

*Crex crex* SHARPE, LAYARD'S Bds. S. Africa, 1884, 611.

[B 558, C 471, R 577, C 683.]

GEOG. DIST.— Europe and northern Asia ; casual in Greenland, Bermuda, and eastern North America.

SUBFAMILY **GALLINULINÆ**. GALLINULES.

GENUS **IONORNIS** REICHENBACH.

*Ionornis* REICH. Syst. Av. 1852, p. xxi. Type, *Fulica martinica* LINN.

218. **Ionornis martinica** (LINN.).

Purple Gallinule.

*Fulica martinica* LINN. S. N. ed. 12, I. 1766, 259.

*Ionornis martinica* REICH. Syst. Av. 1852, p. xxi.

[B 561, C 473, R 578, C 685.]

GEOG. DIST.—South Atlantic and Gulf States, casually northward to Maine, New York, Wisconsin, etc.; south throughout the West Indies, Mexico, Central America and northern South America to Brazil.

GENUS **GALLINULA** BRISSON.

*Gallinula* BRISS. Orn. VI. 1760, 2. Type, *Fulica chloropus* LINN.

219. **Gallinula galeata** (LICHT.).

Florida Gallinule.

*Crex galeata* LICHT. Verz. Doubl. 1823, 80.

*Gallinula galeata* BONAP. Am. Orn. IV. 1832, 128.

[B 560, C 472, R 579, C 685.]

GEOG. DIST.—Temperate and tropical America, from Canada to Brazil and Chili.

SUBFAMILY **FULICINÆ**. COOTS.GENUS **FULICA** LINNÆUS.

*Fulica* LINN. S. N. ed. 10, I. 1758, 152. Type, *F. atra* LINN.

[220.] **Fulica atra** LINN.

European Coot.

*Fulica atra* LINN. S. N. ed. 10, I. 1758, 152.

[B —, C —, R —, C 885.]

GEOG. DIST.—Northern parts of the eastern hemisphere in general; accidental in Greenland.

221. **Fulica americana** GMEL.

American Coot.

*Fulica americana* GMEL. S. N. I. ii. 1788, 704.

[B 559, C 474, R 580, C 686.]

GEOG. DIST.—North America, from Greenland and Alaska southward to the West Indies and Veragua.

## ORDER LIMICOLÆ. SHORE BIRDS.

## FAMILY PHALAROPODIDÆ. PHALAROPES.

## GENUS CRYMOPHILUS VIEILLOT.

*Crymophilus* VIEILL. Anal. 1816, 62. Type, *Tringa fulicaria* LINN.

222. *Crymophilus fulicarius* (LINN.).

Red Phalarope.

*Tringa fulicaria* LINN. S. N. ed. 10, I. 1758, 148.

*Crymophilus fulicarius* STEJN. Auk, II. 1885, 183.

[B 521, C 411, R 563, C 604.]

GEOG. DIST.—Northern parts of northern hemisphere, breeding in the Arctic regions and migrating south in winter; in the United States south to the Middle States, Ohio Valley, and Cape St. Lucas; chiefly maritime.

## GENUS PHALAROPUS BRISSON.

## SUBGENUS PHALAROPUS.

*Phalaropus* BRISS. Orn. VI. 1760, 12. Type, by elimination, *Tringa lobata* LINN.

223. *Phalaropus lobatus* (LINN.).

Northern Phalarope.

*Tringa lobata* LINN. S. N. ed. 10, I. 1758, 148, 824.

*Phalaropus lobatus* SALVAD. Ucc. d'Italia, II. 1872, 210 (nec LATHAM qui *Crymophilus fulicarius*, nec WILSON qui *Ph. tricolor*).

[B 520, C 410, R 564, C 603.]

GEOG. DIST.—Northern portions of northern hemisphere, breeding in arctic latitudes; south in winter to the tropics.



SUBGENUS **STEGANOPUS** VIEILLOT.

*Steganopus* VIEILL. Nouv. Dict. d'Hist. Nat. XXXII. 1819, 136.

Type, *S. tricolor* VIEILL.

224. **Phalaropus tricolor** (VIEILL.).

Wilson's Phalarope.

*Steganopus tricolor* VIEILL. Nouv. Dict. d'Hist. Nat. XXXII. 1819, 136.

*Phalaropus tricolor* STEJN. Auk, II. 1885, 183.

[B 519, C 409, R 565, C 602.]

GEOG. DIST.—Temperate North America, chiefly the interior, breeding from northern Illinois and Utah northward to the Saskatchewan region; south in winter to Brazil and Patagonia.

FAMILY **RECURVIROSTRIDÆ**. AVOCETS AND STILTS.GENUS **RECURVIROSTRA** LINNÆUS.

*Recurvirostra* LINN. S. N. ed. 10, I. 1758, 151. Type, *R. avosetta* LINN.

225. **Recurvirostra americana** GMEL.

American Avocet.

*Recurvirostra americana* GMEL. S. N. I. ii. 1788, 693.

[B 517, C 407, R 566, C 600.]

GEOG. DIST.—Temperate North America, north to the Saskatchewan and Great Slave Lake; in winter, south to Guatemala and the West Indies. Rare in the eastern United States.

GENUS **HIMANTOPUS** BRISSON.

*Himantopus* BRISS. Orn. VI. 1760, 33. Type, *Charadrius himantopus* LINN.

226. **Himantopus mexicanus** (MÜLL.).

Black-necked Stilt.

*Charadrius mexicanus* MÜLL. S. N. Suppl. 1776, 117.*Himantopus mexicanus* ORD, WILS. Orn. VII. 1824, 52.

[B 518, C 408, R 567, C 601.]

GEOG. DIST.—Temperate North America, from the northern United States southward to the West Indies, northern Brazil, and Peru. Rare in eastern United States, except in Florida.

FAMILY **SCOLOPACIDÆ**. SNIPES, SANDPIPERS, ETC.GENUS **SCOLOPAX** LINNÆUS.

*Scolopax* LINN. S. N. ed. 10, I. 1758, 145. Type, *S. rusticola* LINN.

[227.] **Scolopax rusticola** LINN.

European Woodcock.

*Scolopax rusticola* LINN. S. N. ed. 10, I. 1758, 146.

[B—, C 413, R 524, C 606.]

GEOG. DIST.—Northern parts of the Old World; occasional in eastern North America.

GENUS **PHILOHELA** GRAY.*Philohela* GRAY, List Gen. B. 1841, 90. Type, *Scolopax minor* GM.228. **Philohela minor** (GMEL.).

American Woodcock.

*Scolopax minor* GMEL. S. N. I. ii. 1788, 661.*Philohela minor* GRAY, List Gen. B. 1841, 90.

[B 522, C 412, R 525, C 605.]

GEOG. DIST.—Eastern North America, north to the British Provinces, west to Dakota, Kansas, etc.; breeding throughout its range. No extralimital record except Bermuda.

GENUS **GALLINAGO** LEACH.

*Gallinago* LEACH, Syst. Cat. Brit. Mam. & Birds, 1816, 31.  
Type, *Scolopax major* LINN.

[229.] **Gallinago gallinago** (LINN.).

European Snipe.

*Scolopax gallinago* LINN. S. N. ed. 10, I. 1758, 147.

*Gallinago gallinago* LICHT. Nom. Mus. Berol. 1854, 93.

[B —, C —, R 526, C 607.]

GEOG. DIST.—Northern parts of the Old World; frequent in Greenland, accidental in Bermuda.

230. **Gallinago delicata** (ORD).

Wilson's Snipe.

*Scolopax delicata* ORD, WILS. Orn. IX. 1825, p. ccxviii.

*Gallinago delicata* A. O. U. Check-List, 1886, 148.

[B 523, C 414, R 526a, C 608.]

GEOG. DIST.—North and Middle America, breeding from the northern United States northward; south in winter to the West Indies and northern South America.

GENUS **MACRORHAMPHUS** LEACH.

*Macrorhamphus* LEACH, Syst. Cat. Brit. Mam. & B. 1816, 31.

Type, *Scolopax grisea* GMEL.

231. **Macrorhamphus griseus** (GMEL.).

Dowitcher.

*Scolopax grisea* GMEL. S. N. I. ii. 1788, 658.

*Macrorhamphus griseus* LEACH, Syst. Cat. Brit. Mam. & B. 1816, 31.

[B 524, C 415, R 527, C 609.]

GEOG. DIST.—Eastern North America, breeding far north; south in winter to the West Indies and Brazil. Casual (?) in Alaska (Nushagak River), Bermuda, Great Britain, and Europe.

232. **Macrorhamphus scolopaceus** (SAY).

Long-billed Dowitcher.

*Limosa scolopacea* SAY, LONG'S Exp. II. 1823, 170.*Macrorhamphus scolopaceus* LAWR. Ann. Lyc. N. Y. V. 1852, 4, pl. 1.

[B 525, C 415a, R 527a, C 610.]

GEOG. DIST.—Western North America, breeding in Alaska to the Arctic coast, migrating south in winter through the western United States (including Mississippi Valley) to Mexico, and, less commonly, along Atlantic coast.

GENUS **MICROPALAMA** BAIRD.

*Micropalama* BAIRD, B. N. Am. 1858, 726. Type, *Tringa himantopus* BONAP.

233. **Micropalama himantopus** (BONAP.).

Stilt Sandpiper.

*Tringa himantopus* BONAP. Ann. Lyc. N. Y. II. 1826, 157.*Micropalama himantopus* BAIRD, B. N. Am. 1858, 726.

[B 536, C 416, R 528, C 611.]

GEOG. DIST.—Eastern North America, breeding north of the United States, and migrating in winter to Bermuda, West Indies and Central and South America.

GENUS **TRINGA** LINNÆUS.SUBGENUS **TRINGA**.

*Tringa* LINN. S. N. ed. 10, I. 1758, 148. Type, by elimination, *T. canutus* LINN.

234. **Tringa canutus** LINN.

Knot.

*Tringa canutus* LINN. S. N. ed. 10, I. 1758, 149.

[B 526, C 426, R 529, C 626.]

GEOG. DIST.—Nearly cosmopolitan. Breeds in high northern latitudes, but visits the southern hemisphere during its migrations.

SUBGENUS **ARQUATELLA** BAIRD.

*Arquatella* BAIRD, B. N. Am. 1858, 714, 717. Type, *Tringa maritima* BRÜNN.

235. ***Tringa maritima*** BRÜNN.

Purple Sandpiper.

*Tringa maritima* BRÜNN. Orn. Bor. 1764, 54.

[B 528, C 423, R 530, C 620.]

GEOG. DIST.—Northern portions of the northern hemisphere; in North America chiefly the northeastern portions, breeding in the high north, migrating in winter to the Eastern and Middle States (casually to Florida), the Great Lakes, and the shores of the larger streams in the upper Mississippi Valley.

236. ***Tringa couesi*** (RIDGW.).

Aleutian Sandpiper.

*Arquatella couesi* RIDGW. Bull. Nutt. Orn. Club, V. July, 1880, 160.

*Tringa couesi* HARTLAUB, Journ. f. Orn. 1883, 280.

[B —, C —, R 531, C 621.]

GEOG. DIST.—Aleutian Islands and coast of Alaska, north to Kowak River, west to Commander Islands, Kamchatka.

237. ***Tringa ptilocnemis*** COUES.

Pribilof Sandpiper.

*Tringa ptilocnemis* COUES, ELLIOTT's Rep. Seal Isl. Alaska, 1873 (not paged).

[B —, C 426 *bis*, R 532, C 622.]

GEOG. DIST.—Breeding in the Pribilof Islands, Alaska, and migrating to coast of adjacent mainland south of Norton Sound.

SUBGENUS **ACTODROMAS** KAUP.

*Actodromas* KAUP, SKIZZ. Entw.-Gesch. Eur. Thierw. 1829, 37.

Type, *Tringa minuta* LEISL.

238. ***Tringa acuminata*** (HORSF.).

Sharp-tailed Sandpiper.

*Totanus acuminatus* HORSF. Linn. Trans. XIII. 1821, 192.*Tringa acuminata* SWINH. P. Z. S. 1863, 316.

[B —, C —, R 533, C 619.]

GEOG. DIST.—Eastern Asia, and coast of Alaska, migrating south to Java and Australia.

239. ***Tringa maculata*** VIEILL.

Pectoral Sandpiper.

*Tringa maculata* VIEILL. Nouv. Dict. d'Hist. Nat. XXXIV.  
1819, 465.

[B 531, C 420, R 534, C 616.]

GEOG. DIST.—The whole of North America, the West Indies, and the greater part of South America. Breeds in the Arctic regions. Of frequent occurrence in Europe.

240. ***Tringa fuscicollis*** VIEILL.

White-rumped Sandpiper.

*Tringa fuscicollis* VIEILL. Nouv. Dict. d'Hist. Nat. XXXIV.  
1819, 461.

[B 533, C 421, R 536, C 617.]

GEOG. DIST.—Eastern North America, breeding in the high north. In winter, the West Indies, Central and South America, south to the Falkland Islands. Occasional in Europe.

241. ***Tringa bairdii*** (COUES).

Baird's Sandpiper.

*Actodromas bairdii* COUES, Pr. Ac. Nat. Sci. Phila. 1861, 194.*Tringa bairdii* SCL. P. Z. S. 1867, 332.

[B —, C 419, R 537, C 615.]

GEOG. DIST.—Nearly the whole of North and South America, but chiefly the interior of North and the western portions of South America, south to Chili and Patagonia. Breeds in Alaska and on the Barren Grounds. Rare along the Atlantic coast, and not yet recorded from the Pacific coast of the United States.



242. **Tringa minutilla** VIEILL.

Least Sandpiper.

*Tringa minutilla* VIEILL. Nouv. Dict. d'Hist. Nat. XXXIV.  
1819, 452.

[B 532, C 418, R 538, C 614.]

GEOG. DIST.—The whole of North and South America, breeding north of the United States. Accidental in Europe.

[242.1.] **Tringa damacensis** (HORSF.).

Long-toed Stint.

*Totanus damacensis* HORSF. Trans. Linn. Soc. XIII. 1821, 129.*Tringa damacensis* SWINH. Ibis, Oct. 1863, 413.

[B —, C —, R —, C —.]

GEOG. DIST.—Asia, breeding toward Arctic coast; accidental in Alaska (Otter Island, Bering Sea).

SUBGENUS **PELIDNA** CUVIER.*Pelidna* CUV. Règne An. 1817, 490. Type, *Tringa alpina* LINN.[243.] **Tringa alpina** LINN.

Dunlin.

*Tringa alpina* LINN. S. N. ed. 10, I. 1758, 149.

[B —, C —, R 539, C 623.]

GEOG. DIST.—Northern parts of the Old World; accidental in eastern North America (west side of Hudson Bay and Long Island).

243a. **Tringa alpina pacifica** (COUES).

Red-backed Sandpiper.

*Pelidna pacifica* COUES, Pr. Ac. Nat. Sci. Phila. 1861, 189.*Tringa alpina pacifica* A. O. U. Check-List, 1886, 152.

[B 530, C 424, R 539a, C 624.]

GEOG. DIST.—North America in general, breeding far north. Eastern Asia.

SUBGENUS **ANCYLOCHEILUS** KAUP.

*Ancylocheilus* KAUP, Skizz. Entw.-Gesch. Eur. Thierw. 1829, 50.  
Type, *Tringa subarquata* TEMM. = *T. ferruginea* BRÜNN.

244. **Tringa ferruginea** BRÜNN.

Curlew Sandpiper.

*Tringa ferruginea* BRÜNN. Orn. Bor. 1764, 53.

[B 529, C 425, R 540, C 625.]

GEOG. DIST.— Old World in general; occasional in eastern North America and Alaska.

GENUS **EURYNORHYNCHUS** NILSSON.

*Eurynorhynchus* NILSS. Orn. Suec. II. 1821, 29. Type, *Platalea pygmæa* LINN.

[245.] **Eurynorhynchus pygmæus** (LINN.).

Spoon-bill Sandpiper.

*Platalea pygmæa* LINN. S. N. ed. 10, I. 1758, 140.

*Eurynorhynchus pygmæus* PEARSON, Jour. As. Soc. Beng. V. 1836, 127.

[B —, C —, R 542\*, C 884.]

GEOG. DIST.— Asia; in summer along the Arctic coast, in winter southern and southeastern Asia. Accidental on the coast of Alaska (Choris Peninsula).

GENUS **EREUNETES** ILLIGER.

*Ereunetes* ILLIG. Prodr. 1811, 262. Type, *E. petrificatus* ILLIG.  
= *Tringa pusilla* LINN.

246. **Ereunetes pusillus** (LINN.).

Semipalmated Sandpiper.

*Tringa pusilla* LINN. S. N. ed. 12, I. 1766, 252.

*Ereunetes pusillus* CASS. Pr. Ac. Nat. Sci. Phila. 1860, 195.

[B 535, C 417, R 541, C 612.]

GEOG. DIST.—Eastern North America, breeding north of the United States; south in winter to the West Indies and South America.

247. **Ereunetes occidentalis** LAWR.

Western Sandpiper.

*Ereunetes occidentalis* LAWR. Pr. Ac. Nat. Sci. Phila. 1864, 107.

[B —, C 417a, R 541a, C 613.]

GEOG. DIST.—Chiefly western United States, frequent eastward to the Atlantic coast; breeding far north and migrating in winter to Central and South America.

GENUS **CALIDRIS** CUVIER.

*Calidris* CUV. Leç. Anat. Comp. I. 1799–1800, tabl. ii. Type,  
*Tringa arenaria* LINN.

248. **Calidris arenaria** (LINN.).

Sanderling.

*Tringa arenaria* LINN. S. N. ed. 12, I. 1766, 251.

*Calidris arenaria* LEACH, Syst. Cat. Brit. Mam. & B. 1816, 28.

[B 534, C 427, R 542, C 627.]

GEOG. DIST.—Nearly cosmopolitan, breeding in the arctic and subarctic regions, migrating, in America, south to Chili and Patagonia.

GENUS **LIMOSA** BRISSON.

*Limosa* BRISS. Orn. V. 1760, 261. Type, *Scolopax limosa* LINN.

249. **Limosa fedoa** (LINN.).

Marbled Godwit.

*Scolopax fedoa* LINN. S. N. ed. 10, I. 1758, 146.

*Limosa fedoa* SABINE, FRANKLIN'S Journ. Polar Sea, 1823, 689.

[B 547, C 428, R 543, C 628.]

GEOG. DIST.—North America; breeding in the interior (from Iowa and Nebraska, northward to Manitoba and the Saskatchewan), migrating in winter to Guatemala, Yucatan, etc., and Cuba.

250. ***Limosa lapponica baueri*** (NAUM.).

Pacific Godwit.

*Limosa baueri* NAUMANN, Vög. Deutschl. VIII. 1834, 429.

*Limosa lapponica baueri* STEJN. Orn. Expl. Kamtsch. 1885, 122.

[B —, C 430, R 544, C 631.]

GEOG. DIST.—Shores and islands of the Pacific Ocean, from New Zealand and Australia to Kamchatka and Alaska. On the American coast recorded south of Alaska only from La Paz, Lower California.

251. ***Limosa hæmastica*** (LINN.).

Hudsonian Godwit.

*Scolopax hæmastica* LINN. S. N. ed. 10, I. 1758, 147.

*Limosa hæmastica* COUES, Birds Northwest, 1874, 760.

[B 548, C 429, R 545, C 629.]

GEOG. DIST.—Eastern North America and the whole of Middle and South America. Breeds only in the high north.

[252.] ***Limosa limosa*** (LINN.).

Black-tailed Godwit.

*Scolopax limosa* LINN. S. N. ed. 10, I. 1758, 147.

*Limosa limosa* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 356.

[B —, C —, R 546, C 630.]

GEOG. DIST.—Northern parts of the Old World; accidental in Greenland.

GENUS **TOTANUS** BECHSTEIN.

*Totanus* BECH. Orn. Tasch. Deutschl. 1803, 282. Type, *Scolopax totanus* LINN.

SUBGENUS **GLOTTIS** KOCH.

*Glottis* KOCH, Baier. Zool. 1816, 304. Type, *Totanus glottis* BECHST. = *Scolopax nebularius* GUNNER.

[253.] **Totanus nebularius** (GUNN.).

Green-shank.

*Scolopax nebularius* GUNNER. in LEEM, Lapp. Beskr. 1767, 251.

*Totanus nebularius* STEJN. Pr. U. S. Nat. Mus. V. 1882, 37.

[B 538, C 434, R 547, C 635.]

GEOG. DIST.—Eastern hemisphere; accidental in Florida.

254. **Totanus melanoleucus** (GMEL.).

Greater Yellow-legs.

*Scolopax melanoleuca* GMEL. S. N. I. ii. 1788, 659.

*Totanus melanoleucus* VIEILL. Nouv. Dict. d'Hist. Nat. VI. 1816, 398.

[B 539, C 432, R 548, C 633.]

GEOG. DIST.—America in general, breeding from Iowa and northern Illinois, etc., northward, and migrating south to Chili and Argentine Republic.

255. **Totanus flavipes** (GMEL.).

Yellow-legs.

*Scolopax flavipes* GMEL. S. N. I. ii. 1788, 659.

*Totanus flavipes* VIEILL. Nouv. Dict. d'Hist. Nat. VI. 1816, 410.

[B 540, C 433, R 549, C 634.]

GEOG. DIST.—America in general, breeding in the cold temperate and subarctic districts, and migrating south in winter to southern South America. Less common in western than in eastern North America.

SUBGENUS **HELODROMAS** KAUP.

*Helodromas* KAUP, Skizz. Entw.-Gesch. Eur. Thierw. 1829, 144.

Type, *Tringa ochropus* LINN.

256. **Totanus solitarius** (WILS.).

Solitary Sandpiper.

*Tringa solitaria* WILS. Am. Orn. VII. 1813, 53, pl. 58, fig. 3.*Totanus solitarius* BONAP. Journ. Ac. Nat. Sci. Phila. V. 1825, 86.

[B 541, C 435, R 550, C 637.]

GEOG. DIST.—North America, breeding occasionally in the northern United States, more commonly northward, and migrating southward as far as the Argentine Republic and Peru.

256 a. **Totanus solitarius cinnamomeus** BREWSTER.

Western Solitary Sandpiper.

*Totanus solitarius cinnamomeus* BREWSTER, Auk, VII. Oct. 1890, 377.

[B 541, *part*, C 435, *part*, R 550, *part*, C 637, *part*.]

GEOG. DIST.—Pacific coast region, eastward to the Plains.

[257.] **Totanus ochropus** (LINN.).

Green Sandpiper.

*Tringa ocrophus* (err. typ.) LINN. S. N. ed. 10, I. 1758, 149.*Totanus ochropus* TEMM. Man. 1815, 420.

[B —, C —, R 551, C 636.]

GEOG. DIST.—Northern parts of the Old World. Accidental in Nova Scotia.

GENUS **SYMPHEMIA** RAFINESQUE.*Symphemia* RAFINESQUE, Jour. de Phys. LXXXVIII. 1819, 418.Type, *Scolopax semipalmata* GMEL.258. **Symphemia semipalmata** (GMEL.).

Willet.

*Scolopax semipalmata* GMEL. S. N. I. ii. 1788, 659.*Symphemia semipalmata* HARTL. Rev. Zool. 1845, 342.

[B 537, C 431, R 552, C 632.]

GEOG. DIST.—Eastern temperate North America, south to the West Indies and Brazil. Breeds from Florida to New Jersey and locally and rarely to Maine. Accidental in Bermuda and Europe.



258 a. **Symphemia semipalmata inornata** BREWST.

Western Willet.

*Symphemia semipalmata inornata* BREWST. Auk, IV. April, 1887, 145.[B 537, *part*, R 431, *part*, R 552, *part*, C 632, *part*.]

GEOG. DIST.—Western North America, east to Mississippi Valley and Gulf States; in winter, south to Mexico, and, during migrations, sparingly along coast of southern Atlantic States. Breeds from coast of Texas to Manitoba.

GENUS **HETERACTITIS** STEJNEGER.*Heteractitis* STEJN. Auk, I. July, 1884, 236. Type, *Scolopax incanus* GMEL.259. **Heteractitis incanus** (GMEL.).

Wandering Tatler.

*Scolopax incanus* GMEL. S. N. I. ii. 1788, 658.*Heteractitis incanus* STEJN. Auk, I. July, 1884, 236.

[B 542, C 440, R 553, C 642.]

GEOG. DIST.—Pacific coast of America, from Norton Sound, Alaska, to the Galapagos, and west to Kamchatka and Hawaiian Islands; also the more eastern island groups of Polynesia.

GENUS **PAVONCELLA** LEACH.*Pavoncella* LEACH, Syst. Cat. Brit. Mam. & B. 1816, 29. Type, *Tringa pugnax* LINN.[260.] **Pavoncella pugnax** (LINN.).

Ruff.

*Tringa pugnax* LINN. S. N. ed. 10, I. 1758, 148.*Pavoncella pugnax* LEACH, Syst. Cat. Brit. Mam. & B. 1816, 29.

[B 544, C 437, R 554, C 639.]

GEOG. DIST.—Northern parts of the Old World, straying occasionally to eastern North America.

GENUS **BARTRAMIA** LESSON.

*Bartramia* LESS. Traité, 1831, 553. Type, *B. laticauda* LESS.  
= *Tringa longicauda* BECHST.

261. **Bartramia longicauda** (BECHST.).

**Bartramian Sandpiper.**

*Tringa longicauda* BECHST. Uebers. Lath. Ind. Orn. II. 1812,  
453.

*Bartramius longicaudus* BONAP. Rev. et Mag. Zool. XX. 1857, 59.

[B 545, C 438, R 555, C 640.]

GEOG. DIST.—North America, mainly east of the Rocky Mountains, north to Nova Scotia and Alaska, breeding throughout most of its North American range; migrating in winter southward, as far as Brazil and Peru. Occasional in Europe.

GENUS **TRYNGITES** CABANIS.

*Tryngites* CAB. Jour. für Orn. 1856, 418. Type, *Tringa rufescens* VIEILL. = *T. subruficollis* VIEILL.

262. **Tryngites subruficollis** (VIEILL.).

**Buff-breasted Sandpiper.**

*Tringa subruficollis* VIEILL. Nouv. Dict. d'Hist. Nat. XXXIV.  
1819, 465.

*Tryngites subruficollis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885,  
356.

[B 546, C 439, R 556, C 641.]

GEOG. DIST.—North America, especially in the interior; breeds in the Yukon district and the interior of British America, northward to the Arctic coast; South America in winter as far as Uruguay and Peru. Of frequent occurrence in Europe.

GENUS **ACTITIS** ILLIGER.

*Actitis* ILLIG. Prodr. 1811, 262. Type, *Tringa hypoleucos* LINN.

263. **Actitis macularia** (LINN.).

Spotted Sandpiper.

*Tringa macularia* LINN. S. N. ed. 12, I. 1766, 249.*Actitis macularia* NAUMANN, Vög. Deutschl. VIII. 1836, 34.

[B 543, C 436, R 557, C 638.]

GEOG. DIST.—North and South America, from Alaska south to southern Brazil. Breeds throughout temperate North America, less commonly on the Pacific coast. Occasional in Europe.

GENUS **NUMENIUS** BRISSON.

*Numenius* BRISS. Orn. VI. 1760, 311. Type, *Scolopax arquata* LINN.

264. **Numenius longirostris** WILS.

Long-billed Curlew.

*Numenius longirostris* WILS. Am. Orn. VIII. 1814, 24, pl. 64, fig. 4.

[B 549, C 441, R 558, C 643.]

GEOG. DIST.—Temperate North America, migrating south to Guatemala, Cuba, and Jamaica. Breeds in the South Atlantic States, and in the interior through most of its North American range.

265. **Numenius hudsonicus** LATH.

Hudsonian Curlew.

*Numenius hudsonicus* LATH. Ind. Orn. II. 1790, 712.

[B 550, C 442, R 559, C 645.]

GEOG. DIST.—All of North and South America, including the West Indies; breeds in the high north, and winters chiefly south of the United States.

266. **Numenius borealis** (FORST.).

Eskimo Curlew.

*Scolopax borealis* FORST. Phil. Trans. LXII. 1772, 411, 431.

*Numenius borealis* LATH. Ind. Orn. II. 1790, 712.

[B 551, C 443, R 560, C 646.]

GEOG. DIST.—Eastern North America, breeding in the Arctic regions, and migrating south throughout South America.

[267.] **Numenius phaeopus** (LINN.).

Whimbrel.

*Scolopax phaeopus* LINN. S. N. ed. 10, I. 1758, 146.*Numenius phaeopus* LATH. Gen. Syn. Suppl. I. 1787, 291.

[B —, C —, R 561, C 644.]

GEOG. DIST.—Northern parts of the Old World; occasional in Greenland.

[268.] **Numenius tahitiensis** (GMEL.).

Bristle-thighed Curlew.

*Scolopax tahitiensis* GMEL. S. N. I. ii. 1788, 656.*Numenius tahitiensis* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880, 201.

[B —, C 442 bis, R 562, C 647.]

GEOG. DIST.—Islands of the Pacific Ocean. Occasional on the coast of Alaska and Lower California.

FAMILY **CHARADRIIDÆ**. PLOVERS.GENUS **VANELLUS** BRISSON.*Vanellus* BRISS. Orn. V. 1760, 94. Type, *Tringa vanellus* LINN.[269.] **Vanellus vanellus** (LINN.).

Lapwing.

*Tringa vanellus* LINN. S. N. ed. 10, I. 1758, 148.*Vanellus vanellus* LICHT. Nom. Mus. Berol. 1854, 95.

[B —, C —, R 512, C 593.]

GEOG. DIST.—Northern parts of eastern hemisphere. In North America, occasional in Greenland, the islands in Norton Sound, Alaska, and on Long Island.

GENUS **CHARADRIUS** LINNÆUS.SUBGENUS **SQUATAROLA** CUVIER.

*Squatarola* CUV. Règ. An. I. 1817, 467. Type, *Tringa squatarola* LINN.

270. **Charadrius squatarola** (LINN.).

**Black-bellied Plover.**

*Tringa squatarola* LINN. S. N. ed. 10, I. 1758, 149.

*Charadrius squatarola* NAUM. Vög. Deutschl. VII. 1834, 250.

[B 510, C 395, R 513, C 580.]

GEOG. DIST.—Nearly cosmopolitan, but chiefly in the northern hemisphere, breeding far north, and migrating south in winter; in America, to the West Indies, Brazil, and Colombia.

SUBGENUS **CHARADRIUS** LINNÆUS.

*Charadrius* LINN. S. N. ed. 10, I. 1758, 150. Type, *C. apricarius* LINN.

[271.] **Charadrius apricarius** LINN.

**Golden Plover.**

*Charadrius apricarius* LINN. S. N. ed. 10, I. 1758, 150.

[B —, C —, R 514, C 583.]

GEOG. DIST.—Europe, south to Africa in winter; eastern Greenland.

272. **Charadrius dominicus** MÜLL.

**American Golden Plover.**

*Charadrius dominicus* MÜLL. S. N. Suppl. 1776, 116.

[B 503, C 396, R 515, C 581.]

GEOG. DIST.—Arctic America except coast of Bering Sea, migrating southward throughout North and South America to Patagonia.

272 a. **Charadrius dominicus fulvus** (GMEL.).

Pacific Golden Plover.

*Charadrius fulvus* GMEL. S. N. I. ii. 1788, 687.

*Charadrius dominicus fulvus* RIDGW. Pr. U. S. Nat. Mus. III.  
1880, 198.

[B —, C —, R 515a, C 582.]

GEOG. DIST.—Breeding from northern Asia to the Pribilof Islands and coast of Alaska, south in winter through China and India to Australia and Polynesia.

GENUS **ÆGIALITIS** BOIE.

SUBGENUS **OXYECHUS** REICHENBACH.

*Oxyechus* REICH. Syst. Av. 1853, p. xviii. Type, *Charadrius vociferus* LINN.

273. **Ægialitis vocifera** (LINN.).

Killdeer.

*Charadrius vociferus* LINN. S. N. ed. 10, I. 1758, 150.

*Ægialites vociferus* BONAP. Geog. & Comp. List, 1838, 45.

[B 504, C 397, R 516, C 584.]

GEOG. DIST.—Temperate North America, breeding north to Newfoundland and Manitoba, migrating to the West Indies, and Central and northern South America. Bermuda.

SUBGENUS **ÆGIALITIS** BOIE.

*Ægialitis* BOIE, Isis, 1822, 558. Type, by elimination, *Charadrius hiaticula* LINN.

274. **Ægialitis semipalmata** BONAP.

Semipalmated Plover.

*Charadrius semipalmatus* BONAP. Journ. Acad. Nat. Sci. Phila.  
V. 1825, 98.

*Ægialites semipalmatus* BONAP. Geog. & Comp. List, 1838, 45.



[B 507, C 399, R 517, C 586.]

GEOG. DIST.—Arctic and subarctic America, migrating south throughout tropical America, as far as Brazil, Peru, and the Galapagos.

275. **Ægialitis hiaticula** (LINN.).

Ring Plover.

*Charadrius hiaticula* LINN. S. N. ed. 10, I. 1758, 150.

*Ægialitis hiaticula* BOIE, Isis, 1822, 558.

[B —, C —, R 518, C 589.]

GEOG. DIST.—Northern parts of the Old World and portions of Arctic America, breeding on the west shore of Cumberland Gulf.

[276.] **Ægialitis dubia** (SCOP.).

Little Ring Plover.

*Charadrius dubius* SCOPOLI, Delic. F. et Fl. Insubr. II. 1786, 93.

*Ægialites dubius* SWINH. P. Z. S. 1871, 404.

[B —, C 400 *bis*, R 519, C 590.]

GEOG. DIST.—Most of the eastern hemisphere, breeding northward. Accidental on the coast of California and in Alaska.

277. **Ægialitis meloda** (ORD).

Piping Plover.

*Charadrius melodus* ORD, ed. WILS. VII. 1824, 71.

*Ægialites melodus* BONAP. Geog. & Comp. List, 1838, 45.

[B 508, C 400, R 520, C 587.]

GEOG. DIST.—Eastern North America, breeding from the coast of Virginia northward to Newfoundland; in winter, West Indies.

277 a. **Ægialitis meloda circumcincta** RIDGW.

Belted Piping Plover.

*Ægialitis melodus* var. *circumcinctus* RIDGW. Am. Nat. VIII. Feb. 1874, 109.

[B —, C 400a, R 520a, C 588.]

GEOG. DIST.—Mississippi Valley, breeding from northern Illinois north to Lake Winnipeg; more or less frequent eastward to the Atlantic coast.

278. *Ægialitis nivosa* CASS.

Snowy Plover.

*Ægialitis nivosa* CASS. in BAIRD, B. N. Am. 1858, 696.

[B 509, C 401, R 521, C 591.]

GEOG. DIST.—Western United States, from California east to Kansas and western Gulf States; in winter, both coasts of Central America, and western South America to Chili. Western Cuba.

[279.] *Ægialitis mongola* (PALL.).

Mongolian Plover.

*Charadrius mongolus* PALL. Reise Russ. Reich. III. 1776, 700.*Ægialites mongolus* SWINH. P. Z. S. 1870, 140.

[B —, C —, R —, C —.]

GEOG. DIST.—Northern Asia, south in winter to India, Malay Archipelago, Philippines, and Australia. Choris Peninsula, Alaska; accidental.

SUBGENUS **OCHTHODROMUS** REICHENBACH.

*Ochthodromus* REICH. Syst. Av. 1852, p. xviii. Type, *Charadrius wilsonius* ORD.

280. *Ægialitis wilsonia* (ORD).

Wilson's Plover.

*Charadrius wilsonia* ORD, WILS. Orn. IX. 1814, 77, pl. 73, fig. 5.*Ægialites wilsonius* BONAP. Geog. & Comp. List, 1838, 45.

[B 506, C 398, R 522, C 585.]

GEOG. DIST.—Coasts of North and South America, from Long Island and Lower California southward to Brazil and Peru, including the West Indies. Casual north to Nova Scotia.

SUBGENUS **PODASOCYS** COUES.

*Podasocys* COUES, Pr. Ac. Nat. Sci. Phila. 1866, 96. Type, *Charadrius montanus* TOWNS.

281. **Ægialitis montana** (TOWNS.).

Mountain Plover.

*Charadrius montanus* TOWNS. Jour. Ac. Nat. Sci. Phila. VII.  
1837, 192.*Ægialitis montanus* CASS. in BAIRD, B. N. Am. 1858, 693.

[B 505, C 402, R 523, C 592.]

GEOG. DIST.—Chiefly the Plains, from central Kansas to the Rocky Mountains, north to the British boundary, breeding from Kansas northward; westward, especially in winter, to central and southern California, and south to Lower California and San Luis Potosi, Mexico. Accidental in Florida.

FAMILY **APHRIZIDÆ**. SURF BIRDS AND TURNSTONES.SUBFAMILY **APHRIZINÆ**. SURF BIRDS.GENUS **APHRIZA** AUDUBON.

*Aphriza* AUD. Orn. Biog. V. 1839, 249. Type, *A. townsendi* AUD.  
= *Tringa virgata* GMEL.

282. **Aphriza virgata** (GMEL.).

Surf Bird.

*Tringa virgata* GMEL. S. N. I. ii. 1788, 674.*Aphriza virgata* GRAY, Gen. B. III. 1847, pl. cxlvii.

[B 511, C 403, R 511, C 594.]

GEOG. DIST.—Pacific coast of America, from Alaska to Chili.

SUBFAMILY **ARENARIINÆ**. TURNSTONES.GENUS **ARENARIA** BRISSON.*Arenaria* BRISS. Orn. V. 1760, 132. Type, *Tringa interpres* LINN.283. **Arenaria interpres** (LINN.).

Turnstone.

*Tringa interpres* LINN. S. N. ed. 10, I. 1758, 148.*Arenaria interpres* VIEILL. Nouv. Dict. d'Hist. Nat. XXIV, 1819,

345.

[B 515, C 406, R 509, C 598.]

GEOG. DIST.— Nearly cosmopolitan. In America from Greenland and Alaska to the Straits of Magellan ; more or less common in the interior of North America, on the shores of the Great Lakes and the larger rivers. Breeds in high northern latitudes.

284. **Arenaria melanocephala** (VIG.).

**Black Turnstone.**

*Streptilas melanocephalus* VIG. Zool. Jour. IV. Jan. 1829, 356.

*Arenaria melanocephala* STEJN. Auk, I. July, 1884, 229.

[B 516, C 406a, R 510, C 599.]

GEOG. DIST.— Pacific coast of North America, from Point Barrow, Alaska, to Santa Margarita Island, Lower California ; breeding from Alaska south to British Columbia. Accidental in India.

FAMILY **HÆMATOPODIDÆ**. OYSTER-CATCHERS.

GENUS **HÆMATOPUS** LINNÆUS.

*Hæmatopus* LINN. S. N. ed. 10, I. 1758, 152. Type, *H. ostralegus* LINN.

[285.] **Hæmatopus ostralegus** LINN.

**Oyster-catcher.**

*Hæmatopus ostralegus* LINN. S. N. ed. 10, I. 1758, 152.

[B —, C —, R 506, C 595.]

GEOG. DIST.— Sea-coasts of Europe and part of Asia and Africa ; occasional in Greenland.

286. **Hæmatopus palliatus** TEMM.

**American Oyster-catcher.**

*Hæmatopus palliatus* TEMM. Man. II. 1820, 532.

[B 512, C 404, R 507, C 596.]

GEOG. DIST.— Sea-coasts of temperate and tropical America, from New Jersey and western Mexico to Patagonia ; occasional or accidental on the Atlantic coast north to Massachusetts and Grand Menan.

286.1. **Hæmatopus frazari** BREWST.**Frazar's Oyster-catcher.***Hæmatopus frazari* BREWST. Auk, V. Jan. 1888, 84.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California (both coasts), north to Los Coronados Islands.

287. **Hæmatopus bachmani** AUD.**Black Oyster-catcher.***Hæmatopus bachmani* AUD. Orn. Biog. V. 1839, 245, pl. 427.

[B 513, C 405, R 508, C 597.]

GEOG. DIST.—Pacific coast of North America, from the Aleutian Islands to La Paz, Lower California.

FAMILY **JACANIDÆ**. JACANAS.GENUS **JACANA** BRISSON.*Jacana* BRISS. Orn. V. 1760, 121. Type, by elimination, *Fulica spinosa* LINN.[288.] **Jacana spinosa** (LINN.).**Mexican Jacana.***Fulica spinosa* LINN. S. N. ed. 10, I. 1758, 152.*Jacana spinosa* ELLIOT, Auk, V. July, 1888, 297.

[B —, C —, R 568, C 672.]

GEOG. DIST.—Lower Rio Grande Valley, Texas, south to Panama. Cuba ; Haiti.

## ORDER GALLINÆ. GALLINACEOUS BIRDS.

## SUBORDER PHASIANI. PHEASANTS, GROUSE, PARTRIDGES, QUAILS, ETC.

FAMILY **TETRAONIDÆ**. GROUSE, PARTRIDGES, ETC.SUBFAMILY **PERDICINÆ**. PARTRIDGES.GENUS **COLINUS** LESSON.

*Colinus* LESS. Man. d'Orn. II. 1828, 190. Type, *Tetrao virginianus* LINN.

289. **Colinus virginianus** (LINN.).**Bob-white.**

*Tetrao virginianus* LINN. S. N. ed. 10, I. 1758, 161.

*Colinus virginianus* STEJN. Auk, II. Jan. 1885, 45.

[B 471, C 389, R 480, C 571.]

GEOG. DIST.— Eastern United States and southern Ontario, from southern Maine to the South Atlantic and Gulf States ; west to central South Dakota, Nebraska, Kansas, Oklahoma and eastern Texas. Of late years has gradually extended its range westward along lines of railroad and settlements ; also, introduced at various points in Colorado, New Mexico, Utah, Idaho, California, Oregon, and Washington. Breeds throughout its range.

289 a. **Colinus virginianus floridanus** (COUES).**Florida Bob-white.**

*Ortyx virginianus* var. *floridanus* COUES, Key, 1872, 237.

*Colinus virginianus floridanus* STEJN. Auk, II. Jan. 1885, 45.



[B —, C 389*a*, R 480*a*, C 572.]

GEOG. DIST.—Florida.

289 *b*. **Colinus virginianus texanus** (LAWR.).

**Texan Bob-white.**

*Ortyx texanus* LAWR. Ann. Lyc. N. Y. VI. April, 1853, 1.

*Colinus virginianus texanus* STEJN. Auk, II. Jan. 1885, 45.

[B 472, C 389*b*, R 480*b*, C 573.]

GEOG. DIST.—Southern and western Texas, south to central Tamaulipas and southern Nuevo Leon, Mexico. Western Mexico, near Guadalajara.

291. **Colinus ridgwayi** BREWST.

**Masked Bob-white.**

*Colinus ridgwayi* BREWST. Auk, II. April, 1885, 199.

[B —, C —, R —, C —.]

GEOG. DIST.—Sonora to southern Arizona.

GENUS **OREORTYX** BAIRD.

*Oreortyx* BAIRD, B. N. Am. 1858, 642. Type, *Ortyx picta* DOUGL.

292. **Oreortyx pictus** (DOUGL.).

**Mountain Partridge.**

*Ortyx picta* DOUGL. Trans. Linn. Soc. XVI. 1829, 143.

*Oreortyx pictus* BAIRD, B. N. Am. 1858, 642.

[B 473, C 390, R 481, C 574.]

GEOG. DIST.—Pacific coast region, from San Francisco Bay north to Washington. Introduced on Vancouver Island.

292 *a*. **Oreortyx pictus plumiferus** (GOULD).

**Plumed Partridge.**

*Ortyx plumifera* GOULD, P. Z. S. 1837, 42.

*Oreortyx pictus* var. *plumiferus* RIDGW. in Hist. N. Am. B. III.

1874, 476.

[B—, C—, R 481a, C—.]

GEOG. DIST.—Sierra Nevada (both slopes), east to Panamint Mountains, and to Mount Magruder, Nevada; south in the coast ranges from San Francisco Bay to Lower California (Campos).

292 *b.* **Oreortyx pictus confinis** ANTHONY.

San Pedro Partridge.

*Oreortyx picta confinis* ANTHONY, Proc. Cal. Ac. Sci. 2d. ser. II.  
Oct. 11, 1889, 74.

[B—, C—, R—, C—.]

GEOG. DIST.—San Pedro Mountains, Lower California.

## GENUS **CALLIPEPLA** WAGLER.

### SUBGENUS **CALLIPEPLA**.

*Callipepla* WAGLER, Isis, 1832, 277. Type, *C. strenua* WAGL.  
= *Ortyx squamatus* VIG.

293. **Callipepla squamata** (VIG.).

Scaled Partridge.

*Ortyx squamatus* VIG. Zool. Jour. V. 1830, 275.

*Callipepla squamata* GRAY, List Gen. B. ed. 1, 1840, 61.

[B 476, C 393, R 484, C 577.]

GEOG. DIST.—Tablelands of Mexico, from the valley of Mexico, north to central and western Texas, Santa Fé, New Mexico, and southern Arizona.

293 *a.* **Callipepla squamata castanogastris** BREWST.

Chestnut-bellied Scaled Partridge.

*Callipepla squamata castanogastris* BREWST. Bull. Nutt. Orn.  
Club, VIII. Jan. 1883, 34.

[B—, C—, R—, C—.]

GEOG. DIST.—Northeastern Mexico and Lower Rio Grande Valley in Texas.

SUBGENUS **LOPHORTYX** BONAPARTE.

*Lophortyx* BONAP. Geog. & Comp. List, 1838, 42. Type, *Tetrao californicus* SHAW.

294. **Callipepla californica** (SHAW).

California Partridge.

*Tetrao californicus* SHAW, Nat. Misc. IX. 1797 (?), pl. cccxlv.

*Callipepla californica* GOULD, Mon. Odont. "pt. i. 1844," pl. xvi.

[B 474, C 391, R 482, C 575.]

GEOG. DIST.—Coast region of California, south to Monterey. Introduced in Oregon, Washington, and British Columbia.

294 a. **Callipepla californica vallicola** RIDGWAY.

Valley Partridge.

*Callipepla californica vallicola* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B—, C—, R—, C—.]

GEOG. DIST.—Interior valleys of California and foothills of the Sierra Nevada, east to the Panamint Mountains, south to Cape St. Lucas.

295. **Callipepla gambelii** (GAMBEL).

Gambel's Partridge.

*Lophortyx gambelii* "NUTTALL," GAMB. Pr. Ac. Nat. Sci. Phila. 1843, 260.

*Callipepla gambeli* GAMBEL, Journ. Ac. Nat. Sci. Phila. n. s. I, 1849, 219.

[B 475, C 392, R 483, C 576.]

GEOG. DIST.—Western Texas, New Mexico, Arizona, southern Utah, southern Nevada, southern California in the Colorado Valley, and southward into northwestern Mexico.

GENUS **CYRTONYX** GOULD.

*Cyrtonyx* GOULD, Mon. Odont. "pt. i, 1844, pl. vii." Type, *Ortyx massena* LESS. = *O. montezumæ* VIG.

296. **Cyrtonyx montezumæ** (VIG.).

Massena Partridge.

*Ortyx montezumæ* VIGORS, Zool. Journ. V. 1830, 275.

*Cyrtonyx montezumæ* STEJN. Auk, II. Jan. 1885, 46.

[B 477, C 394, R 485, C 578.]

GEOG. DIST.—Tablelands of Mexico, from the city of Mexico north to western Texas, New Mexico, and Arizona.

SUBFAMILY **TETRAONINÆ**. GROUSE.

GENUS **DENDRAGAPUS** ELLIOT.

SUBGENUS **DENDRAGAPUS**.

*Dendragapus* ELLIOT, Pr. Ac. Nat. Sci. Phila. 1864, 23. Type,  
*Tetrao obscurus* SAY.

297. **Dendragapus obscurus** (SAY).

Dusky Grouse.

*Tetrao obscurus* SAY, LONG'S Exp. II. 1823, 14.

*Dendragapus obscurus* ELLIOT, Pr. Ac. Nat. Sci. Phila. 1864, 23.

[B 459, C 381, R 471, C 557.]

GEOG. DIST.—Rocky Mountains, from central Montana and southeastern Idaho to New Mexico and Arizona, eastward to the Black Hills, South Dakota, and westward to East Humboldt Mountains, Nevada.

297 *a*. **Dendragapus obscurus fuliginosus** RIDGW.

Sooty Grouse.

*Canace obscura* var. *fuliginosa* RIDGW. Bull. Essex Inst. V. Dec. 1873, 199.

*Dendragapus obscurus fuliginosus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B—, C 381*b*, R 471*a*, C 559.]

GEOG. DIST.—Northwest coast mountains, from California to Sitka, east to Nevada, western Idaho, and portions of British Columbia.

297 *b. Dendragapus obscurus richardsonii* (DOUGL.).

Richardson's Grouse.

*Tetrao richardsonii* "SAB. MS." DOUGL. Linn. Trans. XVI. iii. 1829, 141.*Dendragapus obscurus richardsoni* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B—, C 381a, R 471b, C 558.]

GEOG. DIST.—Rocky Mountains, especially on the eastern slopes, from central Montana, northern Wyoming, and southeastern Idaho into British America to Liard River.

SUBGENUS **CANACHITES** STEJNEGER.

*Canachites* STEJN. Pr. U. S. Nat. Mus. VIII. 1885, 410. Type, *Tetrao canadensis* LINN.

298. *Dendragapus canadensis* (LINN.).

Canada Grouse.

*Tetrao canadensis* LINN. S. N. ed. 10, I. 1758, 159.*Dendragapus canadensis* TURNER, Pr. U. S. Nat. Mus. VIII. 1885, 245.

[B 460, C 380, R 472, C 555.]

GEOG. DIST.—British America, east of the Rocky Mountains, and west in Alaska to the Pacific coast at Kadiak and St. Michaels, southeastward to northern Minnesota, northern Michigan, northern New York, and northern New England.

299. *Dendragapus franklinii* (DOUGL.).

Franklin's Grouse.

*Tetrao franklinii* DOUGL. Trans. Linn. Soc. XVI. iii. 1829, 139.*Dendragapus franklinii* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 461, C 380a, R 472a, C 556.]

GEOG. DIST.—Northern Rocky Mountains, from northwestern Montana to the coast ranges of Oregon and Washington, and northward in British America, reaching the Pacific coast of southern Alaska (lat. 60° N.).

GENUS **BONASA** STEPHENS.

*Bonasa* STEPH. Gen. Zool. XI. 1819, 298. Type, *Tetrao umbellus* LINN.

300. **Bonasa umbellus** (LINN.).

Ruffed Grouse.

*Tetrao umbellus* LINN. S. N. ed. 12, I. 1766, 275.

*Bonasa umbellus* STEPH. Gen. Zool. XI. 1819, 300.

[B 465, C 385, R 473, C 565.]

GEOG. DIST.—Eastern United States and southern Canada, west to Minnesota, south in the mountains to northern Georgia, Mississippi, and Arkansas.

300 *a.* **Bonasa umbellus togata** (LINN.).

Canadian Ruffed Grouse.

*Tetrao togatus* LINN. S. N. ed. 12, 1766, 275.

*Bonasa umbellus togata* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B —, C —, R —, C —.]

GEOG. DIST.—The spruce forests of northern New England, northern New York, and the British Provinces, west to Oregon, Washington, and British Columbia, north to James Bay.

300 *b.* **Bonasa umbellus umbelloides** (DOUGL.).

Gray Ruffed Grouse.

*Tetrao umbelloides* DOUGL. Trans. Linn. Soc. XVI. iii. 1829, 148.

*Bonasa umbellus* var. *umbelloides* BAIRD, B. N. Am. 1858, 925.

[B 465\*, C 385*a*, R 473*a*, C 566.]

GEOG. DIST.—Rocky Mountain region of the United States and British America, north to Alaska, east to Manitoba.

300 *c.* **Bonasa umbellus sabini** (DOUGL.).

Oregon Ruffed Grouse.

*Tetrao sabini* DOUGL. Trans. Linn. Soc. XVI. iii. 1829, 137.

*Bonasa umbellus* var. *sabinei* COUES, Key, 1872, 235.



[B 466, C 385*b*, R 473*b*, C 567.]

GEOG. DIST.—Coast ranges of northern California, Oregon, Washington, and British Columbia.

GENUS **LAGOPUS** BRISSON.

*Lagopus* BRISS. Ornith. I. 1760, 181. Type, *Tetrao lagopus* LINN.

301. **Lagopus lagopus** (LINN.).

**Willow Ptarmigan.**

*Tetrao lagopus* LINN. S. N. ed. 10, I. 1758, 159.

*Lagopus lagopus* STEJN. Pr. U. S. Nat. Mus. VIII. 1885, 20.

[B 467, 470, C 386, R 474, C 568.]

GEOG. DIST.—Arctic regions; in America south to Sitka and the British Provinces. Breeding range restricted to the arctic and sub-arctic regions, mainly north of 55° N. lat. Accidental in New England (Bangor, Me., and Essex Co., Mass.).

301 *a*. **Lagopus lagopus alleni** STEJN.

**Allen's Ptarmigan.**

*Lagopus alba alleni* STEJN. Auk, I. Oct. 1884, 369.

*Lagopus lagopus alleni* STEJN. Pr. U. S. Nat. Mus. VIII. 1885, 20.

[B —, C —, R —, C —.]

GEOG. DIST.—Newfoundland.

302. **Lagopus rupestris** (GMEL.).

**Rock Ptarmigan.**

*Tetrao rupestris* GMEL. S. N. I. ii. 1788, 751.

*Lagopus rupestris* LEACH, Zool. Misc. II. 1817, 290.

[B 468, C 387, R 475, C 569.]

GEOG. DIST.—Arctic America (except the northern extremity), from Alaska to Labrador and the Gulf of St. Lawrence; portions of Greenland; Aleutian Islands.

302 a. **Lagopus rupestris reinhardi** (BREHM).

Reinhardt's Ptarmigan.

*Tetrao reinhardi* BREHM, Lehrb. Eur. Vög. 1823, 440.

*Lagopus rupestris reinhardti* BLASIUS, List Eur. B. 1862, 16.

[B —, C —, R —, C —.]

GEOG. DIST.—Greenland, western shores of Cumberland Gulf, and northern extremity of Labrador.

302 b. **Lagopus rupestris nelsoni** STEJN.

Nelson's Ptarmigan.

*Lagopus rupestris nelsoni* STEJN. Auk, I. July, 1884, 226.

[B —, C —, R —, C —.]

GEOG. DIST.—Unalaska, and some adjacent Aleutian Islands.

302 c. **Lagopus rupestris atkhensis** (TURNER).

Turner's Ptarmigan.

*Lagopus mutus atkhensis* TURNER, Pr. U. S. Nat. Mus. V. July 29, 1882, 227, 230.

*Lagopus rupestris atkhensis* NELSON, Cruise Corwin, 1883, 56 e + 82.

[B —, C —, R —, C —.]

GEOG. DIST.—Atka, one of the Aleutian Islands.

303. **Lagopus welchi** BREWST.

Welch's Ptarmigan.

*Lagopus welchi* BREWST. Auk, II. April, 1885, 194.

[B —, C —, R —, C —.]

GEOG. DIST.—Newfoundland.

304. **Lagopus leucurus** SWAINS. & RICH.

White-tailed Ptarmigan.

*Lagopus leucurus* SWAINS. & RICH. Fauna Bor. Amer. II. 1831, pl. 63.

[B 469, C 388, R 476, C 570.]

GEOG. DIST.—Alpine summits of the mountains of western North America, from New Mexico to Liard River, British America, west on the highest ranges of Oregon, Washington, and British Columbia.

GENUS **TYMPANUCHUS** GLOGER.

*Tympanuchus* GLOGER, Gemeinnutzig. Hand. und Hilfsbuch Naturg. 1842, 396. Type, *Tetrao cupido* LINN.

305. **Tympanuchus americanus** (REICH.).

Prairie Hen.

*Cupidonia americanus* REICH. Syst. Av. 1852, p. xxix; based on Vollst. Naturg. Hühnen. pl. 217, figs. 1896-1898.

*Tympanuchus americanus* RIDGW. Auk, Jan. III. 1886, 133.

[B 464, C 384, R 477, C 563.]

GEOG. DIST.—Prairies of the Mississippi Valley; south to Louisiana and Texas, east to Kentucky, Indiana, Ohio, Michigan, and Ontario; west through eastern portions of North Dakota, South Dakota, Nebraska, Kansas, and the Indian Territory; north to Manitoba; general tendency to extension of range westward and contraction eastward; migration north and south in Minnesota, Iowa, and Missouri.

305 a. **Tympanuchus americanus attwateri** (BENDIRE).

Attwater's Prairie Hen.

*Tympanuchus attwateri* BENDIRE, Forest and Stream, XL. No. 20, May 18, 1893, 425.

*Tympanuchus americanus attwateri* A. O. U. Check-List, 6th Suppl. Auk, XI. April, 1894, 130.

[B 464, *part*, C 384, *part*, R 477, *part*, C 563, *part*.]

GEOG. DIST.—Coast region of Louisiana and Texas.

306. **Tympanuchus cupido** (LINN.).

Heath Hen.

*Tetrao cupido* LINN. S. N. ed. 10, I. 1758, 160.

*Tympanuchus cupido* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 464, *part*, C 384, *part*, R 477, *part*, C 563 *part*.]

GEOG. DIST.—Island of Martha's Vineyard, Mass. (Formerly southern New England and parts of the Middle States.)

307. **Tympanuchus pallidicinctus** RIDGW.

Lesser Prairie Hen.

*Cupidonia cupido* var. *pallidicincta* RIDGW. Bull. Essex Inst. V.  
Dec. 1873, 199.*Tympanuchus pallidicinctus* RIDGW. Pr. U. S. Nat. Mus. VIII.  
1885, 355.

[B—, C 384a, R 477a, C 564.]

GEOG. DIST.—Eastern edge of the Great Plains from western and probably southern Texas northward through Indian Territory to Kansas.

GENUS **PEDIOCÆTES** BAIRD.*Pediocætes* BAIRD, B. N. Am. 1858, 625. Type, *Tetrao phasianellus* LINN.308. **Pediocætes phasianellus** (LINN.).

Sharp-tailed Grouse.

*Tetrao phasianellus* LINN. S. N. ed. 10, I. 1758, 160.*Pediocætes phasianellus* ELLIOT, Pr. Ac. Nat. Sci. Phila. 1862,  
403 (nec BAIRD, 1858, qui subsp. *columbianus*).

[B—, C 383, R 478, C 561.]

GEOG. DIST.—Interior of British America, from Lake Superior and Hudson Bay to Fort Simpson.

308 a. **Pediocætes phasianellus columbianus** (ORD).

Columbian Sharp-tailed Grouse.

*Phasianus columbianus* ORD, GUTHRIE'S Geog. 2d Am. ed. II.  
1815, 317.*Pediocætes phasianellus* var. *columbianus* COUES, Key, 1872, 234.

[B 463, C 383a, R 478a, C 562.]

GEOG. DIST.—Plains of the northwestern United States and British Columbia to central portions of Alaska; northward chiefly west of the main Rocky Mountains; eastward in Montana and Wyoming; southward to Utah, northern Nevada, and northeastern California.

- 308 b. **Pediocætes phasianellus campestris** RIDGW.  
**Prairie Sharp-tailed Grouse.**

*Pedieetes phasianellus campestris* RIDGW. Proc. Biol. Soc. Wash.  
 II. April 10, 1884, 93.

[B —, C —, R —, C —.]

GEOG. DIST.— Plains and prairies of the United States east of the Rocky Mountains; north to Manitoba; east to Wisconsin and Illinois; south to New Mexico.

GENUS **CENTROCERCUS** SWAINSON.

*Centrocercus* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831,  
 358, 496. Type, *Tetrao urophasianus* BONAP.

309. **Centrocercus urophasianus** (BONAP.).  
**Sage Grouse.**

*Tetrao urophasianus* BONAP. Zool. Journ. III. 1827, 213.

*Centrocercus urophasianus* SWAINS. in SW. & RICH. Fauna Bor.  
 Am. II. 1831, 497, pl. 58.

[B 462, C 382, R 479, C 560.]

GEOG. DIST.— Sage regions of the Rocky Mountain plateau and westward, chiefly within the United States, but north to Assiniboia and the dry interior of British Columbia; east to North Dakota, South Dakota, Nebraska, and Colorado; south to northern New Mexico, Utah, and Nevada; west in California, Oregon and Washington to the Sierra Nevada and Cascade Range.

FAMILY **PHASIANIDÆ**. PHEASANTS, ETC.

SUBFAMILY **MELEAGRINÆ**. TURKEYS.

GENUS **MELEAGRIS** LINNÆUS.

*Meleagris* LINN. S. N. ed. 10, I. 1758, 156. Type, *M. gallopavo* LINN.

310. **Meleagris gallopavo** LINN.

Wild Turkey.

*Meleagris gallopavo* LINN. S. N. ed. 10, I. 1758, 156.

[B 457, C 379a, R 470a, C 554.]

GEOG. DIST.—United States, from Chesapeake Bay to the Gulf coast, and west to the Plains, along wooded river valleys; formerly north to southern Maine, southern Ontario, and up the Missouri River to North Dakota.

310 a. **Meleagris gallopavo mexicana** (GOULD).

Mexican Turkey.

*Meleagris mexicana* GOULD, P. Z. S. 1856, 61.

*Meleagris gallopavo* var. *mexicana* BAIRD, Hist. N. Am. B. III.  
1874, 410.

[B 458, C 379, R 470, C 553.]

GEOG. DIST.—Southwestern United States, from western Texas to Arizona; south over the tablelands of Mexico.

310 b. **Meleagris gallopavo osceola** SCOTT.

Florida Wild Turkey.

*Meleagris gallopavo osceola* SCOTT, Auk, VII. Oct. 1890, 376.[B 457, *part*, C 379a, *part*, R 470a, *part*, C 554, *part*.]

GEOG. DIST.—Southern Florida.

310 c. **Meleagris gallopavo ellioti** SENNETT.

Rio Grande Turkey.

*Meleagris gallopavo ellioti* SENNETT, Auk, IX. April, 1892, 167,  
pl. iii.

[B 458, *part*, C 379, *part*, R 470, *part*, C 553, *part*.]

GEOG. DIST.—Lowlands of southern Texas and northeastern Mexico.



## SUBORDER PENELOPES. CURASSOWS AND GUANS.

FAMILY **CRACIDÆ**. CURASSOWS AND GUANS.SUBFAMILY **PENELOPINÆ**. GUANS.GENUS **ORTALIS** MERREM.

*Ortalis* MERR. AV. RAR. Icones et Desc. II. 1786, 40. Type,  
*Phasianus motmot* LINN.

311. **Ortalis vetula maccalli** BAIRD.**Chachalaca.***Ortalida maccalli* BAIRD, B. N. Am. 1858, 611.*Ortalida vetula* var. *maccalli* BAIRD, Hist. N. Am. B. III. 1874,  
398.

[B 456, C 378, R 469, C 552.]

GEOG. DIST.—Valley of the Lower Rio Grande, and southward in  
 Mexico to Vera Cruz.

## ORDER COLUMBÆ. PIGEONS.

FAMILY **COLUMBIDÆ**. PIGEONS.GENUS **COLUMBA** LINNÆUS.

*Columba* LINN. S. N. ed. 10, I. 1758, 162. Type, by elimination,  
*C. anas* LINN.

312. **Columba fasciata** SAY.**Band-tailed Pigeon.***Columba fasciata* SAY, LONG'S Exp. II. 1823, 10.

[B 445, C 367, R 456, C 539.]

GEOG. DIST.—Western United States, from the Rocky Mountains to the Pacific; north to Washington and British Columbia; south to Mexico and the highlands of Guatemala; distribution irregular, chiefly in wooded mountain regions.

312 *a.* ***Columba fasciata vioscæ*** BREWST.

**Viosca's Pigeon.**

*Columba fasciata vioscæ* BREWST. Auk, V. Jan. 1888, 86.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California.

313. ***Columba flavirostris*** WAGL.

**Red-billed Pigeon.**

*Columba flavirostris* WAGL. Isis, 1831, 519.

[B 446, C 368, R 457, C 540.]

GEOG. DIST.—Southern border of the United States, from Arizona and the Rio Grande Valley south to Costa Rica, breeding throughout its United States range.

314. ***Columba leucocephala*** LINN.

**White-crowned Pigeon.**

*Columba leucocephala* LINN. S. N. ed. 10, I. 1758, 164.

[B 447, C 369, R 458, C 541.]

GEOG. DIST.—Southern Keys of Florida, the Bahamas, the Greater Antilles and some of the Lesser Antilles, and the coast of Honduras.

GENUS **ECTOPISTES** SWAINSON.

*Ectopistes* SWAINS. Zool. Journ. III. 1827, 362. Type, *Columba migratoria* LINN.

315. ***Ectopistes migratorius*** (LINN.).

**Passenger Pigeon.**

*Columba migratoria* LINN. S. N. ed. 12, I. 1766, 285 (♂).

*Ectopistes migratoria* SWAINS. Zool. Journ. III. 1827, 362.

[B 448, C 370, R 459, C 543.]

GEOG. DIST.—Eastern North America, from Hudson Bay southward, and west to the Great Plains, straggling thence to Nevada and Washington. Breeding range now mainly restricted to portions of the Canadas and the northern border of the United States, as far west as Manitoba and the Dakotas.

GENUS **ZENAIDURA** BONAPARTE.

*Zenaidura* BONAP. Consp. Av. II. Dec. 1854, 84. Type, *Columba carolinensis* LINN. = *C. macroura* LINN.

316. **Zenaidura macroura** (LINN.).

Mourning Dove.

*Columba macroura* LINN. S. N. ed. 10, 1758, 164 (part).

*Zenaidura macroura* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 451, C 371, R 460, C 544.]

GEOG. DIST.—Temperate North America; from southern Maine, southern Canada, and British Columbia, south to Panama and the West Indies, breeding throughout its North American range.

GENUS **ZENAIDA** BONAPARTE.

*Zenaida* BONAP. Geog. & Comp. List, 1838, 41. Type, *Columba zenaida* BONAP.

317. **Zenaida zenaida** (BONAP.).

Zenaida Dove.

*Columba zenaida* BONAP. Jour. Ac. Nat. Sci. Phila. V. 1825, 30.

*Zenaida zenaida* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 449, C 372, R 462, C 545.]

GEOG. DIST.—Florida Keys, Bahamas, Cuba, Jamaica, Porto Rico, Santa Cruz, Sombrero, and coast of Yucatan.

GENUS **LEPTOTILA** SWAINSON.

*Leptotila* SWAINS. Class. Bds. II. 1837, 349. Type, *P[eristera]. rufaxilla* Selby, Nat. Libr. V. pl. 24.

318. **Leptotila fulviventris brachyptera** (SALVADORI).

White-fronted Dove.

*Leptotila brachyptera* SALVADORI, Cat. Bds. Brit. Mus. XXI.  
1893, p. 545.*Leptotila fulviventris brachyptera* A. O. U. Check-List, 7th  
Suppl. Auk, XII, April, 1895, 167.

[B—, C—, R 463, C 542.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, and  
southward to Nicaragua.GENUS **MELOPELIA** BONAPARTE.*Melopelia* BONAP. Consp. Av. II. Dec. 1854, 81. Type, *Columba  
leucoptera* LINN.319. **Melopelia leucoptera** (LINN.).

White-winged Dove.

*Columba leucoptera* LINN. S. N. ed. 10, I. 1758, 164.*Melopelia leucoptera* BONAP. Consp. Av. II. Dec. 1854, 81.

[B 450, C 373, R 464, C 546.]

GEOG. DIST.—Southern border of the United States, from Florida,  
Texas, New Mexico, Arizona, and Lower California, southward to  
Costa Rica and the West Indies; Cuba; Jamaica; straggles north-  
ward to Colorado. Breeding range, in the United States, southern  
Texas, New Mexico, and Arizona.GENUS **COLUMBIGALLINA** BOIE.*Columbigallina* BOIE, Isis, 1826, 977. Type, *Columba passerina*  
LINN.320. **Columbigallina passerina terrestris** CHAPM.

Ground Dove.

*Columba passerina* LINN. Syst. Nat. ed. 10, I. 1758, 165.*Columbigallina passerina terrestris* CHAPMAN, Bull. Am. Mus.  
Nat. Hist., IV, 1892, 292.

[B 453, C 374, R 465, C 547.]

GEOG. DIST.—South Atlantic and Gulf States; West Indies and  
northern South America. Breeding, in the United States, from  
South Carolina to Louisiana, chiefly coastwise.

320 a. **Columbigallina passerina pallescens** (BAIRD).

Mexican Ground Dove.

*Chamæpelis passerina* ? var. *pallescens* BAIRD, Proc. Acad. Nat. Sci. Phila. 1859, 305.*Columbigallina passerina pallescens* FERRARI-PEREZ, Pr. U. S. Nat. Mus. IX, 1886, 175.

[B 453, part, C 374, part, R 465, part, C 547, part.]

GEOG. DIST.—Mexico, and contiguous territory of United States from Texas to Lower California; south on both Mexican coasts to Central America.

GENUS **SCARDAFELLA** BONAPARTE.*Scardafella* BONAP. Consp. Av. II. Dec. 1854, 85. Type, *Columba squamosa* TEMM.321. **Scardafella inca** (LESS.).

Inca Dove.

*Chamæpelis inca* LESSON, Descr. Mam. et Ois. 1847, 211.*Scardafella inca* BONAP. Consp. Av. II. Dec. 1854, 85.

[B 452, C 375, R 466, C 549.]

GEOG. DIST.—Rio Grande Valley, southern Arizona, and Lower California, south to Nicaragua.

GENUS **GEOTRYGON** GOSSE.*Geotrygon* GOSSE, B. Jam. 1847, 316, foot-note. Type, *G. sylvatica* GOSSE = *Columba cristata* TEMM.[322.] **Geotrygon martinica** (LINN.).

Key West Quail-Dove.

*Columba martinica* LINN. S. N. ed. 12, I. 1766, 283.*Geotrygon martinica* BONAP. Consp. Av. II. Dec. 1854, 74.

[B 454, C 376, R 467, C 550.]

GEOG. DIST.—Key West, and some of the other Florida Keys; Bahamas; Cuba; Hayti.

[322.1.] **Geotrygon montana** (LINN.).

Ruddy Quail-Dove.

*Columba montana* LINN. S. N. ed. 10, I. 1758, 163.*Geotrygon montana* BONAP. Consp. II, Dec. 1854, 72.

[B—, C—, R—, C—.]

GEOG. DIST.—Tropical America in general, including West Indies; north to Cuba and eastern Mexico; accidental at Key West, Florida.

GENUS **STARNÆNAS** BONAPARTE.

*Starnænas* BONAP. Geog. & Comp. List, 1838, 41. Type,  
*Columba cyanocephala* LINN.

[323.] **Starnænas cyanocephala** (LINN.).

Blue-headed Quail-dove.

*Columba cyanocephala* LINN. S. N. ed. 10, I. 1758, 163.*Starnænas cyanocephala* BONAP. Geog. & Comp. List, 1838, 41.

[B 455, C 377, R 468, C 551.]

GEOG. DIST.—Florida Keys and Cuba.

## ORDER RAPTORES. BIRDS OF PREY.

SUBORDER SARCORHAMPHI. AMERICAN  
VULTURES.FAMILY **CATHARTIDÆ**. AMERICAN VULTURES.GENUS **PSEUDOGRYPHUS** RIDGWAY.*Pseudogryphus* RIDGW. Hist. N. Am. B. III. Jan. 1874, 337, 338.Type, *Vultur californianus* SHAW.



324. **Pseudogryphus californianus** (SHAW).**California Vulture.***Vultur californianus* SHAW, Nat. Misc. IX. 1797, pl. ccci.*Pseudogryphus californianus* RIDGW. Hist. N. Am. B. III. 1874, 338.

[B 2, C 364, R 453, C 536.]

GEOG. DIST.—Coast ranges of southern California from Monterey Bay southward into Lower California; formerly north to Frazer River.

GENUS **CATHARTES** ILLIGER.

*Cathartes* ILLIG. Prodr. 1811, 236. Type, by elimination, *Vultur aura* LINN.

325. **Cathartes aura** (LINN.).**Turkey Vulture.***Vultur aura* LINN. S. N. ed. 10, I. 1758, 86.*Cathartes aura* SPIX, Aves Bras. I. 1825, 2.

[B 1, C 365, R 454, C 537.]

GEOG. DIST.—Temperate North America, from New Jersey, Ohio Valley, Saskatchewan region, and British Columbia southward to Patagonia and the Falkland Islands. Casual in New England.

GENUS **CATHARISTA** VIEILLOT.

*Catharista* VIEILL. Analyse, 1816, 21. Type, by elimination, *Vultur atratus* BARTR.

326. **Catharista atrata** (BARTR.).**Black Vulture.***Vultur atratus* BARTR. Trav. Car. 1792, 285.*Catharista atrata* GRAY, Handl. I. 1869, 3.

[B 3, C 366, R 455, C 538.]

GEOG. DIST.—South Atlantic and Gulf States, north regularly to North Carolina and the lower Ohio Valley, west to the Great Plains, and south through Mexico and Central America, the West Indies, and most of South America. Straggling north to New York, New England, and South Dakota. Breeds in the United States from North Carolina coastwise to Texas, and in the interior to Indiana, Illinois, and Kansas.

SUBORDER FALCONES. VULTURES, FALCONS,  
HAWKS, BUZZARDS, EAGLES, KITES,  
HARRIERS, ETC.

FAMILY **FALCONIDÆ**. VULTURES, FALCONS, HAWKS, EAGLES,  
ETC.

SUBFAMILY **ACCIPITRINÆ**. KITES, BUZZARDS, HAWKS, GOSHAWKS,  
EAGLES, ETC.

GENUS **ELANOIDES** VIEILLOT.

*Elanoides* VIEILL. Nouv. Dict. d'Hist. Nat. XXIV. 1818, 101.

Type, *Falco furcatus* = *F. forficatus* LINN.

327. **Elanoides forficatus** (LINN.).

Swallow-tailed Kite.

*Falco forficatus* LINN. S. N. ed. 10, I. 1758, 89.

*Elanoides forficatus* COUES, Pr. Ac. Nat. Sci. Phila. 1875, 345.

[B 34, C 337, R 426, C 493.]

GEOG. DIST.—United States, especially in the interior, from the Carolinas and Minnesota southward, throughout Central and South America; westward to the Great Plains. Casual eastward to southern New England, and northward to Manitoba and Assiniboia. Breeding range irregularly coincident with general distribution in the United States. Accidental in England.

GENUS **ELANUS** SAVIGNY.

*Elanus* SAVIG. Descr. de l'Égypte, 1809, 97. Type, *E. casius*  
= *Falco melanopterus* DAUD.

328. **Elanus leucurus** (VIEILL.).

White-tailed Kite.

*Milvus leucurus* VIEILL. Nouv. Dict. d'Hist. Nat. XX. 1818, 563  
(errore 556).

*Elanus leucurus* BONAP. Geog. & Comp. List, 1838, 4.

[B 35, C 336, R 427, C 492.]

GEOG. DIST.—Southern United States, from South Carolina and southern Illinois to Texas and California, southward to Chili and Argentine Republic; casual in Michigan. Breeds irregularly throughout its general distribution in the United States.

GENUS **ICTINIA** VIEILLOT.

*Ictinia* VIEILL. Analyse, 1816, 24. Type, *Falco mississippiensis* WILS.

329. **Ictinia mississippiensis** (WILS.).

Mississippi Kite.

*Falco mississippiensis* WILS. Am. Orn. III. 1811, 80, pl. 25, fig. 1.

*Ictinia mississippiensis* GRAY, Gen. B. I. 1845, 26.

[B 36, C 335, R 428, C 491.]

GEOG. DIST.—Southern United States, east of the Rocky Mountains; southward regularly from South Carolina on the coast, and casually from Pennsylvania, Wisconsin and Iowa in the interior, to Guatemala.

GENUS **ROSTRHAMUS** LESSON.

*Rostrhamus* LESS. Traité, 1831, 55. Type, *Falco hamatus* ILLIG.

330. **Rostrhamus sociabilis** (VIEILL.).

Everglade Kite.

*Herpetotheres sociabilis* VIEILL. Nouv. Dict. d'Hist. Nat. XVIII. 1817, 318.

*Rostrhamus sociabilis* D'ORB. Voy. Ois. II. 1847, 73.

[B 37, C 334, R 429, C 490.]

GEOG. DIST.—Florida, Cuba, eastern Mexico, Central America, and eastern South America, to the Argentine Republic.

GENUS **CIRCUS** LACÉPÈDE.

*Circus* LACÉP. Mem. de l'Inst. III. 1801, 506. Type, *Falco cyaneus* LINN.

331. **Circus hudsonius** (LINN.).

Marsh Hawk.

*Falco hudsonius* LINN. S. N. ed. 12, I. 1766, 128.*Circus hudsonius* VIEILL. Ois. Am. Sept. I. 1807, pl. 9.

[B 38, C 333, R 430, C 489.]

GEOG. DIST.—North America in general; south to Panama and Cuba. Breeds throughout its North American range.

GENUS **ACCIPITER** BRISSON.SUBGENUS **ACCIPITER**.

*Accipiter* BRISS. Orn. I. 1760, 310. Type, by elimination, *Falco nisus* LINN.

332. **Accipiter velox** (WILS.).

Sharp-shinned Hawk.

*Falco velox* WILS. Am. Orn. V. 1812, 116, pl. 45, fig. 1.*Accipiter velox* VIGORS, Zool. Journ. I. 1824, 338.

[B 17, C 338, R 432, C 494.]

GEOG. DIST.—North America in general; south to Panama. Breeds throughout its North American range.

333. **Accipiter cooperii** (BONAP.).

Cooper's Hawk.

*Falco cooperii* BONAP. Am. Orn. II. 1828, 1, pl. x. fig. 1.*Accipiter cooperi* GRAY, List B. Brit. Mus. Accipitres, 1844, 38.

[B 15, 16, C 339, R 431, C 495.]

GEOG. DIST.—North America, from southern British America south to southern Mexico. Breeds throughout its range.

SUBGENUS **ASTUR** LACÉPÈDE.

*Astur* LACÉP. Mem. de l'Inst. III. 1801, 505. Type, *Falco palumbarius* LINN.

334. **Accipiter atricapillus** (WILS.).

American Goshawk.

*Falco atricapillus* WILS. Am. Orn. VI. 1812, 80, pl. 52, fig. 3.*Accipiter atricapillus* SEEBOHM, Brit. Birds, I. 1883, iv.

[B 14, C 340, R 433, C 496.]

GEOG. DIST.—Northern and eastern North America, south in winter to the Middle States and southern Rocky Mountain region; casually west to Oregon. Accidental in England. Breeding range restricted to the Canadian Fauna of the United States and northward.

334 a. **Accipiter atricapillus striatulus** RIDGW.

Western Goshawk.

*Astur atricapillus* var. *striatulus* RIDGW. in Hist. N. Am. B. III. 1874, 240.*Accipiter atricapillus striatulus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B —, C —, R 433 a, C 497.]

GEOG. DIST.—Western North America; north to Sitka, Alaska; south to California; east to Idaho. Breeds in the Sierra Nevada south to Lat. 38°.

GENUS **PARABUTEO** RIDGWAY.*Parabuteo* RIDGW. in Hist. N. Am. B. III. Jan. 1874, 250. Type, *Falco harrisi* AUD.335. **Parabuteo unicinctus harrisi** (AUD.).

Harris's Hawk.

*Falco harrisi* AUD. B. Am. V. 1839, 30, pl. 392.*Parabuteo unicinctus* var. *harrisi* RIDGW. in Hist. N. Am. B. III. Jan. 1874, 250.

[B 46, C 348, R 434, C 512.]

GEOG. DIST.—Mississippi, Louisiana, Texas, southern New Mexico, southern Arizona, and Lower California; southward to Panama. Breeds from southern Texas westward to California, and southward.

GENUS **BUTEO** CUVIER.

*Buteo* CUV. Leç. Anat. Comp. I. tabl. ii. Ois. 1779-1800. Type,  
*Falco buteo* LINN.

[336.] **Buteo buteo** (LINN.).

European Buzzard.

*Falco buteo* LINN. S. N. ed. 10, I. 1758, 90.

*Buteo buteo* LICHT. Nomencl. Mus. Berol. 1854, 3.

[B—, C—, R 435, C—.]

GEOG. DIST.—Europe and Western Asia. Accidental in North America (Michigan?).

337. **Buteo borealis** (GMEL.).

Red-tailed Hawk.

*Falco borealis* GMEL. S. N. I. ii. 1788, 266.

*Buteo borealis* VIEILL. Nouv. Dict. d'Hist. Nat. IV. 1816, 478.

[B 23, C 351, R 436, C 516.]

GEOG. DIST.—Eastern North America, west to the Great Plains, north to about Lat. 60°, south to eastern Mexico. Breeds throughout its range, except possibly the extreme southern portion.

337 a. **Buteo borealis kriderii** HOOPES.

Krider's Hawk.

*Buteo borealis* var. *kriderii* HOOPES, Pr. Ac. Nat. Sci. Phila.

1873, 238, pl. 5.

[B—, C 351c, R 436a, C 519.]

GEOG. DIST.—Plains of the United States, from Wyoming and the Dakotas to Minnesota, and south to Texas; casual in Iowa and Illinois.

337 b. **Buteo borealis calurus** (CASS.).

Western Red-tail.

*Buteo calurus* CASS. Pr. Ac. Nat. Sci. Phila. VII. 1855, 281.

*Buteo borealis* var. *calurus* RIDGW. Bull. Essex Inst. V. Nov.

1873, 186.



[B 20, 24, C 351*a*, R 436*b*, C 517.]

GEOG. DIST.—Western North America, from the Rocky Mountains to the Pacific, south into Mexico; casual east to Illinois.

337 *c*. **Buteo borealis lucasanus** RIDGW.

Saint Lucas Red-tail.

*Buteo borealis* var. *lucasanus* RIDGW. in COUES'S Key, 1872, 216 (under *B. borealis*).

[B —, C 351*b*, R 436*c*, C 518.]

GEOG. DIST.—Peninsula of Lower California.

337 *d*. **Buteo borealis harlani** (AUD.).

Harlan's Hawk.

*Falco harlani* AUD. Orn. Biog. I. 1830, 441, pl. 86.

*Buteo borealis harlani* RIDGW. Auk, VII. April, 1890, p. 205.

[B 22, C 350, R 438, C 515.]

GEOG. DIST.—Gulf States and lower Mississippi Valley, north casually to Pennsylvania, Iowa, and Kansas; south to Central America.

339. **Buteo lineatus** (GMEL.).

Red-shouldered Hawk.

*Falco lineatus* GMEL. S. N. I. ii. 1788, 268.

*Buteo lineatus* VIEILL. Nouv. Dict. d'Hist. Nat. IV. 1816, 478.

[B 25, C 352, R 439, C 520.]

GEOG. DIST.—Eastern North America to Manitoba and Nova Scotia; west to Texas and the Plains; south to the Gulf States and Mexico. Breeds throughout its range.

339 *a*. **Buteo lineatus alleni** RIDGW.

Florida Red-shouldered Hawk.

*Buteo lineatus alleni* RIDGW. Pr. U. S. Nat. Mus. VII. Jan. 19, 1885, 514.

[B —, C —, R —, C —.]

GEOG. DIST.—South Atlantic and Gulf States, chiefly coastwise from South Carolina to Texas. Breeds throughout its range.

339 *b. Buteo lineatus elegans* (CASS.).

Red-bellied Hawk.

*Buteo elegans* CASS. Pr. Ac. Nat. Sci. Phila. 1855, 281.*Buteo lineatus* var. *elegans* RIDGW. in Hist. N. Am. B. III. Jan. 1874, 257, 277.

[B 26, C 352a, R 439a, C 521.]

GEOG. DIST.—Western United States, from western Texas to California and Oregon, south into Lower California, Sonora, and Chihuahua. Breeds along the Pacific coast of central Oregon, south to northern Lower California.

340. *Buteo abbreviatus* CAB.

Zone-tailed Hawk.

*Buteo abbreviatus* CAB. in SCHOMB. Reise Brit. Guian. III. 1848, 739.

[B —, C 353, R 440, C 522.]

GEOG. DIST.—Texas, New Mexico, Arizona, and southern California, south to northern South America.

SUBGENUS **TACHYTRIORCHIS** KAUP.

*Tachytriorchis* KAUP, Class. Säug. u. Vög. 1844, 123. Type, *Falco pterocles* TEMM. = *Buteo albicaudatus* VIEILL.

341. *Buteo albicaudatus sennetti* ALLEN.

Sennett's White-tailed Hawk.

*Buteo albicaudatus sennetti* ALLEN, Bull. Am. Mus. Nat. Hist. V. 1893, 144.

[B —, C —, R 441, C 513.]

GEOG. DIST.—Lower Rio Grande Valley, Texas, and southward into Mexico.

342. *Buteo swainsoni* BONAP.

Swainson's Hawk.

*Buteo swainsoni* BONAP. Geog. & Comp. List, 1838, 3.

[B 18, 19, 21, 28, C 354, R 442, C 523.]

GEOG. DIST.—Western North America, from Wisconsin, Illinois, Arkansas, and Texas to the Pacific coast; north to the arctic regions, and south to Argentine Republic. Casual east to Maine and Massachusetts. Breeds nearly throughout its North American range.

343. **Buteo latissimus** (WILS.).

Broad-winged Hawk.

*Falco latissimus* WILS. Am. Orn. VI. 1812, 92, pl. 54, fig. 1.

*Buteo latissimus* SHARPE, Cat. B. Brit. Mus. I. Accip. 1874, 193.

[B 27, C 355, R 443, C 524.]

GEOG. DIST.—Eastern North America, from New Brunswick and the Saskatchewan region to Texas and Mexico, and thence southward to northern South America and the West Indies. Breeds throughout its United States range.

SUBGENUS **BUTEOLA** BONAPARTE.

*Buteola* BONAP. Compt. Rend. XLI. 1855, 651. Type, *Buteo brachyurus* VIEILL.

344. **Buteo brachyurus** VIEILL.

Short-tailed Hawk.

*Buteo brachyurus* VIEILL. Nouv. Dict. d'Hist. Nat. IV. 1816, 477.

[B —, C —, R —, C —.]

GEOG. DIST.—Mexico, Central America, and most of South America, north to Florida, where it regularly breeds.

GENUS **URUBITINGA** LESSON.

*Urubitinga* LESS. Rev. Zool. 1839, 132. Type, *Falco urubitinga* GMEL.

345. **Urubitinga anthracina** (LICHT.).

Mexican Black Hawk.

*Falco anthracinus* LICHT. Preis-Verz. 1830, 3.

*Urubitinga anthracina* LAFR. Rev. Zool. 1848, 241.

[B—, C—, R 444, C 528.]

GEOG. DIST.—Lower Rio Grande Valley and Arizona, southward to northern South America.

GENUS **ASTURINA** VIEILLOT.

*Asturina* VIEILL. Analyse, 1816, 24. Type, *Falco nitidus* GMEL.

346. ***Asturina plagiata*** SCHLEGEL.

Mexican Goshawk.

*Asturina plagiata* "LICHT." SCHLEGEL, Mus. P. B. Asturinæ, 1862, 1.

[B 33, C 358, R 445, C 527.]

GEOG. DIST.—Southwestern border of the United States, southward to Panama.

GENUS **ARCHIBUTEO** BREHM.

*Archibuteo* BREHM, Isis, 1828, 1269. Type, *Falco lagopus* GMEL.

[347.] ***Archibuteo lagopus*** (BRÜNN.).

Rough-legged Hawk.

*Falco lagopus* BRÜNN. Orn. Bor. 1764, 4.

*Archibuteo lagopus* GRAY, List Gen. B. ed. 2, 1841, 3.

[B—, C—, R—, C—.]

GEOG. DIST.—Northern parts of the Old World; Alaska (?).

347a. ***Archibuteo lagopus sancti-johannis*** (GMEL.).

American Rough-legged Hawk.

*Falco sancti-johannis* GMEL. S. N. I. ii. 1788, 273.

*Archibuteo lagopus* var. *sancti-johannis* COUES, Key, 1872, 218.

[B 30, 31, C 356, R 447, C 525.]

GEOG. DIST.—North America north of Mexico, breeding north of the United States (excepting in Alaska).

348. **Archibuteo ferrugineus** (LICHT.).

Ferruginous Rough-Leg.

*Falco ferrugineus* LICHT. Abh. K. Akad. Berl. 1838, 428.*Archibuteo ferrugineus* GRAY, Gen. B. fol. ed. 1849, 12.

[B 32, C 357, R 448, C 526.]

GEOG. DIST.—Western North America, from the Plains (eastern North Dakota to Texas) westward to the Pacific, and from the Saskatchewan region south into Mexico; casually east to Illinois. Breeds from Utah, Colorado, and Kansas northward to the Saskatchewan Plains.

GENUS **AQUILA** BRISSON.*Aquila* BRISS. Orn. I. 1760, 419. Type, *Falco chrysaëtos* LINN.349. **Aquila chrysaëtos** (LINN.).

Golden Eagle.

*Falco chrysaëtos* LINN. S. N. ed. 10, I. 1758, 88.*Aquila chrysaëtos* DUMONT, Dict. Sci. Nat. I. 1816, 339.

[B 39, C 361, R 449, C 532.]

GEOG. DIST.—North America, south to Mexico, and northern parts of the Old World. Breeding range in the United States, practically restricted to the mountainous parts of unsettled regions.

GENUS **THRASAËTOS** GRAY.

*Thrasaëtos* GRAY, P. Z. S. 1837 (June, 1838), 108. Type,  
*Vultur harpyia* LINN.

[350.] **Thrasaëtos harpyia** (LINN.).

Harpy Eagle.

*Vultur harpyia* LINN. S. N. ed. 10, I. 1758, 86.*Thrasaëtos harpyia* GRAY, P. Z. S. 1837 (June, 1838), 108.

[B —, C —, R 450, C 631.]

GEOG. DIST.—Lower Rio Grande Valley, casual; south to Paraguay. Louisiana (?).

GENUS **HALIÆTUS** SAVIGNY.

*Haliæetus* SAVIGNY, Descr. de l'Égypte, 1809, 35. Type, *Falco albicilla* LINN.

[351.] **Haliæetus albicilla** (LINN.).

Gray Sea Eagle.

*Falco albicilla* LINN. S. N. ed. 10, I. 1758, 89.

*Haliæetus albicilla* LEACH, Syst. Cat. M. B. Br. Mus. 1816, 9.

[B 42, C —, R 452, C 533.]

GEOG. DIST.—Northern Europe and Asia; Greenland.

352. **Haliæetus leucocephalus** (LINN.).

Bald Eagle.

*Falco leucocephalus* LINN. S. N. ed. 12, I. 1766, 124.

*Haliæetus leucocephalus* BOIE, Isis, 1822, 548.

[B 41, 43, C 362, R 451, C 534.]

GEOG. DIST.—North America at large, south to Mexico, north-west through the Aleutian Islands to Kamchatka. Breeds locally throughout its range.

SUBFAMILY **FALCONINÆ**. FALCONS.GENUS **FALCO** LINNÆUS.

*Falco* LINN. S. N. ed. 10, I. 1758, 88. Type, by elimination, *F. subbuteo* LINN.

SUBGENUS **HIEROFALCO** CUVIER.

*Hierofalco* CUV. Règ. An. I. 1817, 312. Type, *Falco candicans* GMEL. = *F. islandus* BRÜNN.

353. **Falco islandus** BRÜNN.

White Gyrfalcon.

*Falco islandus* BRÜNN. Orn. Bor. 1764, 2.



[B 11, C 341a, R 412, C 501.]

GEOG. DIST.—Arctic regions, including Arctic America and Greenland, wandering south in winter to Labrador and northern Maine.

354. **Falco rusticolus** LINN.

Gray Gyrfalcon.

*Falco rusticolus* LINN. S. N. ed. 10, I. 1758, 88.

[B 12, C —, R 412a, C 500.]

GEOG. DIST.—Arctic regions, including Iceland, southern Greenland, and Arctic America, straggling southward in winter.

354 a. **Falco rusticolus gyrfalco** (LINN.).

Gyrfalcon.

*Falco gyrfalco* LINN. S. N. ed. 10, I. 1758, 91.

*Falco rusticolus gyrfalco* STEJN. Auk, II. 1885, 187.

[B —, C 341, R 412b, C 498.]

GEOG. DIST.—Northern Europe, Greenland, and Arctic America, from northern Labrador and Hudson Bay to Alaska; rarely south in winter to New England (Massachusetts and Rhode Island).

354 b. **Falco rusticolus obsoletus** (GMEL.).

Black Gyrfalcon.

*Falco obsoletus* GMEL. S. N. I. i. 1788, 268.

*Falco rusticolus obsoletus* STEJN. Auk, II. 1885, 187.

[B —, C —, R 412c, C 499.]

GEOG. DIST.—Labrador, south in winter to Canada, Maine, and New York.

355. **Falco mexicanus** SCHLEG.

Prairie Falcon.

*Falco mexicanus* SCHLEG. Abh. Geb. Zool. 1841, 15.

[B 10, C 342, R 413, C 502.]

GEOG. DIST.—United States, from the eastern border of the Plains to the Pacific, and from the Dakotas south into Mexico; casual eastward to Illinois. Breeds throughout its United States range.

SUBGENUS **RHYNCHODON** NITZSCH.

*Rhynchodon* NITZSCH, Pterylog. 1840, 78. Type, by elimination, *Falco peregrinus* LATH.

356. **Falco peregrinus anatum** (BONAP.).

Duck Hawk.

*Falco anatum* BONAP. Geog. & Comp. List, 1838, 4.

*Falco peregrinus*  $\beta$ . *anatum* BLASIUS, List B. Eur. 1862, 3.

[B 5, 6, C 343, R 414, C 503.]

GEOG. DIST.—North America at large, and south to Chili. Breeds locally throughout most of its United States range.

356 a. **Falco peregrinus pealei** RIDGW.

Peale's Falcon.

*Falco communis* var. *pealei* RIDGW. Bull. Essex Inst. V. Dec. 1873, 201.

*Falco peregrinus pealei* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880, 192.

[B —, C 343a, R 414a, C 504.]

GEOG. DIST.—Pacific coast region of North America, from Oregon to the Aleutian and Commander Islands, breeding throughout its range.

SUBGENUS **ÆSALON** KAUP.

*Æsalon* KAUP, Skizz. Entw.-Gesch. Eur. Thierw. 1829, 40. Type, *Falco æsalon* GMEL. = *F. regulus* PALL.

357. **Falco columbarius** LINN.

Pigeon Hawk.

*Falco columbarius* LINN. S. N. ed. 10, I. 1758, 90.

[B 7, C 344, R 417, C 505.]

GEOG. DIST.—The whole of North America, south to the West Indies and northern South America. Breeds chiefly north of the United States.

357 a. **Falco columbarius suckleyi** RIDGW.

Black Merlin.

*Falco columbarius* var. *suckleyi* RIDGW. Bull. Essex Inst. V. Dec. 1873, 201.

[B —, C 344a, R 417a, C 506.]

GEOG. DIST.—Northwest coast region of North America, from California to Sitka; eastward in Oregon and Washington.

358. **Falco richardsonii** RIDGW.

Richardson's Merlin.

*Falco (Hypotriorchis) richardsonii* RIDGW. Pr. Ac. Nat. Sci. Phila. Dec. 1870, 145.

[B —, C 345, R 418, C 507.]

GEOG. DIST.—Interior and western plains of North America, from the Mississippi River to the Pacific coast, and from the Saskatchewan region to Texas, Arizona, and probably into Mexico.

[358.1.] **Falco regulus** PALL.

Merlin.

*Falco regulus* PALL. Reis. Russ. Reichs. II. 1773, Anhang, 707.

[B —, C —, R —, C —.]

GEOG. DIST.—Europe, Asia, and Africa; accidental in Greenland.

SUBGENUS **RHYNCHOFALCO** RIDGWAY.*Rhynchofalco* RIDGW. Pr. Boston Soc. Nat. Hist. 1873, 46. Type, *Falco femoralis* TEMM. = *F. fusco-cærulescens* VIEILL.359. **Falco fusco-cærulescens** VIEILL.

Aplomado Falcon.

*Falco fusco-cærulescens* VIEILL. Nouv. Dict. d'Hist. Nat. XI. 1817, 90.

[B 9, C 347, R 419, C 511.]

GEOG. DIST.—Southern Texas, New Mexico and Arizona, south to Patagonia.

SUBGENUS **TINNUNCULUS** VIEILLOT.*Tinnunculus* VIEILL. Ois. Am. Sept. I. 1807, 39.[359.1.] **Falco tinnunculus** LINN.

Kestrel.

*Falco tinnunculus* LINN. S. N. ed. 10, I. 1758, 90.

[B —, C —, R 422, C —.]

GEOG. DIST.—Europe and northern Asia; accidental in Massachusetts.

360. **Falco sparverius** LINN.

American Sparrow Hawk.

*Falco sparverius* LINN. S. N. ed. 10, I. 1758, 90.

[B 13, C 346, 346a, R 420, 420a, C 508, 509.]

GEOG. DIST.—North America, east of the Rocky Mountains, and from Great Slave Lake south to northern South America.

360 a. **Falco sparverius deserticolus** MEARNs.

Desert Sparrow Hawk.

*Falco sparverius deserticolus* MEARNs, Auk, IX. July, 1892, 263.[B 13, *part*, C 346, *part*, R 420, *part*, C 508, *part*.]

GEOG. DIST.—Western United States, north to eastern British Columbia and western Montana, south to Mazatlan in northwestern Mexico.

360 b. **Falco sparverius peninsularis** MEARNs.

St. Lucas Sparrow Hawk.

*Falco sparverius peninsularis* MEARNs, Auk, IX. July, 1892, 267.[B 13, *part*, C 346, *part*, R 420, *part*, C 508, *part*.]

GEOG. DIST.—Lower California.

[361.] **Falco dominicensis** GMEL.

Cuban Sparrow Hawk.

*Falco dominicensis* GMEL. Syst. Nat. I. i. 1788, 288.

[B —, C —, R 421, C 510.]

GEOG. DIST.—Cuba; casual in Florida.

GENUS **POLYBORUS** VIEILLOT.

*Polyborus* VIEILL. Analyse, 1816, 22. Type, *Falco tharus* MOL.

362. **Polyborus cheriway** (JACQ.).

Audubon's Caracara.

*Falco cheriway* JACQ. Beitr. 1784, 17, tab. 4.

*Polyborus cheriway* CAL. in SCHOMB. Reise Brit. Guiana, III. 1848, 741.

[B 45, C 363, R 423, C 535.]

GEOG. DIST.—Florida, Texas, Arizona, and Lower California, south to Ecuador and Guiana. Breeds throughout its United States range and southward.

363. **Polyborus lutosus** RIDGW.

Guadalupe Caracara.

*Polyborus lutosus* RIDGW. Bull. U. S. Geog. & Geol. Surv. Terr. No. 6, 2d ser. Feb. 8, 1876, 459.

[B —, C —, R 424, C —.]

GEOG. DIST.—Guadalupe Island, Lower California.

SUBFAMILY **PANDIONINÆ**. OSPREYS.GENUS **PANDION** SAVIGNY.

*Pandion* SAVIGN. Descr. de l'Égypte, Ois. 1809, 95. Type, *Falco haliaëtus* LINN.

364. **Pandion haliaëtus carolinensis** (GMEL.).

American Osprey.

*Falco carolinensis* GMEL. S. N. I. i. 1788, 263.

*Pandion haliaëtus carolinensis* RIDGW. Pr. Ac. Nat. Sci. Phila. Dec. 1870, 143.

[B 44, C 360, R 425, C 530.]

GEOG. DIST.—North America, from Hudson Bay and Alaska south to the West Indies and northern South America. Breeds throughout its North American range.

## SUBORDER STRIGES. OWLS.

FAMILY **STRIGIDÆ**. BARN OWLS.GENUS **STRIX** LINNÆUS.

*Strix* LINN. S. N. ed. 10, I. 1758, 92. Type, *S. aluco* LINN.  
ed. 10.

365. **Strix pratincola** BONAP.

American Barn Owl.

*Strix pratincola* BONAP. Geog. & Comp. List, 1838, 7.

[B 47, C 316, R 394, C 461.]

GEOG. DIST.—United States, rarely to the northern border, and Ontario, southward through Mexico; northern limit of breeding range about Lat. 41°.

FAMILY **BUBONIDÆ**. HORNED OWLS, ETC.GENUS **ASIO** BRISSON.

*Asio* BRISS. Orn. I. 1760, 28. Type, *Strix otus* LINN.

366. **Asio wilsonianus** (LESS.).

American Long-eared Owl.

*Otus wilsonianus* LESS. Traité, 1831, 110.

*Asio wilsonianus* COUES, Check List, ed. 2, 1882, 81, No. 472.

[B 51, C 320, R 395, C 472.]

GEOG. DIST.—Temperate North America; south to the tablelands of Mexico. Breeds throughout its range.

367. **Asio accipitrinus** (PALL.).

Short-eared Owl.

*Strix accipitrina* PALL. Reise Russ. Reichs. I. 1771, 455.

*Asio accipitrinus* NEWT. YARR. Brit. B. ed. 4, I. 1872, 163.



[B 52, C 321, R 396, C 473.]

GEOG. DIST.—Throughout North America; nearly cosmopolitan. Breeds, somewhat irregularly and locally, from about Lat. 39° northward.

GENUS **SYRNIUM** SAVIGNY.

*Syrnium* SAVIGN. Descr. de l'Égypte, Ois. 1809, 298. Type,  
*Strix stridula* LINN.

368. **Syrnium nebulosum** (FORST.).

Barred Owl.

*Strix nebulosa* FORST. Philos. Trans. XXII. 1772, 386.

*Syrnium nebulosum* BOIE, Isis, 1828, 315.

[B 54, C 323, R 397, C 476.]

GEOG. DIST.—Eastern United States, west to Minnesota, Nebraska, Kansas, and Texas, north to Nova Scotia and Quebec. Breeds throughout its range.

368 a. **Syrnium nebulosum alleni** RIDGW.

Florida Barred Owl.

*Strix nebulosa alleni* RIDGW. Pr. U. S. Nat. Mus. III. March, 1880, 8.

*Syrnium nebulosum alleni* RIDGW. Pr. U. S. Nat. Mus. VIII. Sept. 1885, 355.

[B —, C —, R 397a, C 477.]

GEOG. DIST.—South Atlantic and Gulf States, chiefly coastwise, from South Carolina to Texas.

369. **Syrnium occidentale** XANTUS.

Spotted Owl.

*Syrnium occidentale* XANTUS, Pr. Ac. Nat. Sci. Phila. 1859, 193.

[B —, C 324, R 398, C 478.]

GEOG. DIST.—Southern Colorado, New Mexico, Arizona, California, Lower California and Mexico.

GENUS **SCOTIAPTEX** SWAINSON.

*Scotiaptex* SWAINS. Classif. B. II. 1837, 217. Type, *Strix cinerea* GMEL.

370. **Scotiaptex cinerea** (GMEL.).

Great Gray Owl.

*Strix cinerea* GMEL. S. N. I. i. 1788, 291.

*Scotiaptex cinerea* SWAINS. Classif. B. II. 1837, 217.

[B 53, C 322, R 399, C 474.]

GEOG. DIST.—Arctic America, straggling southward, in winter, to southern New England, New York, New Jersey, Ohio, Illinois, Idaho and northern Montana.

[370 a.] **Scotiaptex cinerea lapponica** (RETZ.).

Lapp Owl.

*Strix lapponica* RETZ. Faun. Suec. 1800, 79.

*Scotiaptex cinerea lapponica* RIDGW. Man. N. Am. B. 1887, 260.

[B —, C —, R 399a, C 475.]

GEOG. DIST.—Arctic portions of the Old World; accidental in Alaska (Norton Sound).

GENUS **NYCTALA** BREHM.

*Nyctala* BREHM, Isis, 1828, 1271. Type, *Strix tengmalmi* GMEL.

371. **Nyctala tengmalmi richardsoni** (BONAP.).

Richardson's Owl.

*Nyctale richardsoni* BONAP. Geog. & Comp. List, 1838, 7.

*Nyctale tengmalmi* var. *richardsoni* RIDGW. Am. Nat. VI. May 1872, 285.

[B 55, C 327, R 400, C 482.]

GEOG. DIST.—Arctic America, south in winter into the northern United States. Breeds from the Gulf of St. Lawrence and Manitoba northward.

372. **Nyctala acadica** (GMEL.).

Saw-whet Owl.

*Strix acadica* GMEL. S. N. I. 1788, 296.*Nyctale acadica* BONAP. Geog. & Comp. List, 1838, 7.

[B 56, 57, C 328, R 401, C 483.]

GEOG. DIST.—North America at large, breeding from the Middle States northward, and in mountainous regions of the West southward into Mexico.

GENUS **MEGASCOPS** KAUP.*Megascops* KAUP, Isis, 1848, 765. Type, *Strix asio* LINN.373. **Megascops asio** (LINN.).

Screech Owl.

*Strix asio* LINN. Syst. Nat. ed. 10, I. 1758, 92.*Megascops asio* STEJN. Auk, II. April, 1885, 184.[B 49, *part*, C 318, R 402, C 465.]

GEOG. DIST.—Temperate eastern North America, south to Georgia, and west to the Plains. Accidental in England.

373 *a*. **Megascops asio floridanus** (RIDGW.).

Florida Screech Owl.

*Scops asio* var. *floridanus* RIDGW. Bull. Essex Inst. Dec. 1873, 200.*Megascops asio floridanus* STEJN. Auk, II. April, 1885, 184.[B —, C 318*c*, R 402*a*, C 469.]

GEOG. DIST.—South Atlantic and Gulf States, from South Carolina to Louisiana, chiefly coastwise.

373 *b*. **Megascops asio trichopsis** (WAGL.).

Texas Screech Owl.

*Scops trichopsis* WAGL. Isis, 1832, 276.*Megascops asio trichopsis* RIDGW. Auk, XII. Oct. 1895, —.

[B 50, C 318*b*, R 402*b*, C 468.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, south to Guatemala.

373 *c.* **Megascops asio bendirei** (BREWST.).

California Screech Owl.

*Scops asio bendirei* BREWST. Bull. Nutt. Orn. Club, VII. Jan. 1882, 31.

*Megascops asio bendirei* STEJN. Auk, II. April, 1885, 184.

[B —, C —, R —, C —.]

GEOG. DIST.—Coast region of California.

373 *d.* **Megascops asio kennicottii** (ELLIOT).

Kennicott's Screech Owl.

*Scops kennicottii* ELLIOT, Pr. Ac. Nat. Sci. Phila. 1867, 99.

*Megascops asio kennicottii* STEJN. Auk, II. April, 1885, 184.

[B —, C 318*a*, R 402*d*, C 466.]

GEOG. DIST.—Northwest coast region, from Oregon to Sitka.

373 *e.* **Megascops asio maxwelliæ** (RIDGW.).

Rocky Mountain Screech Owl.

*Scops asio* var. *maxwelliæ* RIDGW. Field & Forest, June, 1877, 210, 213.

*Megascops asio maxwelliæ* STEJN. Auk, II. April, 1885, 184.

[B —, C —, R 402*c*, C 467.]

GEOG. DIST.—Rocky Mountains, from Colorado to Montana.

373 *f.* **Megascops asio cineraceus** RIDGW.

Mexican Screech Owl.

*Megascops asio cineraceus* RIDGW. Auk, XII. Oct. 1895, —.

[B —, C —, R 403, C 470.]

GEOG. DIST.—New Mexico, Arizona, Lower California, and western Mexico.

373 g. **Megascops asio aikeni** BREWST.

Aiken's Screech Owl.

*Megascops asio aikeni* BREWST. Auk, VIII. April, 1891, 139.

[B —, C —, R —, C —.]

GEOG. DIST.—Plains, El Paso County, Colorado, south probably to central New Mexico and northeastern Arizona.

373 h. **Megascops asio macfarlanei** BREWST.

MacFarlane's Screech Owl.

*Megascops asio macfarlanei* BREWST. Auk, VIII. April, 1891, 140.[B 49, *part*, C 318, *part*, R 402, *part*, C 465, *part*.]

GEOG. DIST.—East of the Cascades in Washington, interior of southern British Columbia, southward to central Oregon and eastward into Montana.

374. **Megascops flammeola** (KAUP).

Flammulated Screech Owl.

*Scops flammeola* KAUP, Trans. Zool. Soc. Lond. IV. 1862, 226.*Megascops flammeolus* STEJN. Auk, II. April, 1885, 184.

[B —, C 319, R 404, C 471.]

GEOG. DIST.—Highlands of Guatemala and central Mexico, north to Colorado and northern California.

374 a. **Megascops flammeola idahoensis** MERRIAM.

Dwarf Screech Owl.

*Megascops flammeolus idahoensis* MERRIAM, North Am. Fauna,  
No. 5, July, 1891, 96, pl. i.

[B —, C —, R —, C —.]

GEOG. DIST.—Ketchum, Idaho. Known only from the type.

GENUS **BUBO** DUMÉRIL.

*Bubo* DUMÉR. Zool. Anal. 1806, 34. Type, by implication, *Strix*  
*bubo* LINN.

375. **Bubo virginianus** (GMEL.).

## Great Horned Owl.

*Strix virginiana* GMEL. S. N. I. i. 1788, 287.*Bubo virginianus* BONAP. Geog. & Comp. List, 1838, 6.

[B 48, C 317, R 405, C 462.]

GEOG. DIST.—Eastern North America, west to the Mississippi Valley, and from Labrador south to Costa Rica.

375 a. **Bubo virginianus subarcticus** (HOY).

## Western Horned Owl.

*Bubo subarcticus* HOY, Pr. Ac. Nat. Sci. Phila. VI. 1852, 211.*Bubo virginianus*  $\beta$  *subarcticus* RIDGW. Orn. 40th Par. 1877, 572.[B 48, *part*, C 317a, *part*, R 405a, *part*, C 463, *part*.]

GEOG. DIST.—Western United States, from the Great Plains westward; southward to the Mexican tablelands; east, casually, to Wisconsin and Illinois; north to Manitoba and British Columbia.

375 b. **Bubo virginianus arcticus** (SWAINS.).

## Arctic Horned Owl.

*Strix (Bubo) arctica* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831, 86, pl. 30.*Bubo virginianus* var. *arcticus* CASS. Illust. B. Cal. etc. 1854, 178.[B 48, *part*, C 317a, *part*, R 405b, C 463, *part*.]

GEOG. DIST.—Arctic America, chiefly in the interior, south, in winter, to Idaho, Montana, Wyoming, and South Dakota.

375 c. **Bubo virginianus saturatus** RIDGW.

## Dusky Horned Owl.

*Bubo virginianus saturatus* RIDGW. Orn. 40th Par. 1877, 572, foot-note.[B 48, *part*, C 317b, R 405c, C 464.]

GEOG. DIST.—Pacific coast region, from Monterey Bay, California, northward to Alaska and eastward to Hudson Bay and Labrador; southward through the mountainous regions of the West to Arizona.



GENUS **NYCTEA** STEPHENS.

*Nyctea* STEPH. Gen. Zool. XIII. ii. 1826, 63. Type, *Strix nyctea* LINN.

376. **Nyctea nyctea** (LINN.).  
Snowy Owl.

*Strix nyctea* LINN. S. N. ed. 10, I. 1758, 93.

*Nyctea nyctea* LICHT. Nomen. Mus. Berol. 1854, 7.

[B 61, C 325, R 406, C 479.]

GEOG. DIST.—Northern portions of the northern hemisphere. In North America breeding wholly north of the United States; in winter migrating south to the Middle States, straggling to South Carolina, Texas, California, and Bermuda.

GENUS **SURNIA** DUMÉRIL.

*Surnia* DUMÉR. Zool. Anal. 1806, 34. Type, *Strix ulula* LINN.

[377.] **Surnia ulula** (LINN.).  
Hawk Owl.

*Strix ulula* LINN. S. N. ed. 10, I. 1758, 93.

*Surnia ulula* BONAP. Cat. Met. Ucc. Eur. 1842, 22.

[B—, C—, R 407*a*, C 481.]

GEOG. DIST.—Arctic portions of the Old World. Casual in Alaska (St. Michaels).

377 *a*. **Surnia ulula caparoch** (MÜLL.).  
American Hawk Owl.

*Strix caparoch* MÜLL. S. N. Suppl. 1776, 69.

*Surnia ulula caparoch* STEJN. Auk, I. Oct. 1884, 363.

[B 62, C 326, R 407, C 480.]

GEOG. DIST.—Arctic America, breeding from Newfoundland northward, and migrating in winter to the northern border of the United States. Occasional in England.

GENUS **SPEOTYTO** GLOGER.

*Speotyto* GLOG. Handb. Naturg. 1842, 226. Type, *Strix cunicularia* MOL.

378. **Speotyto cunicularia hypogæa** (BONAP.).  
Burrowing Owl.

*Strix hypogæa* BONAP. Am. Orn. I. 1825, 72.

*Sphcotyto cunicularia* var. *hypogæa* COUES Key, 1872, 208.

[B 58, 59, C 332, R 408, C 487.]

GEOG. DIST.—Western United States, from the Pacific coast east throughout the Great Plains, north somewhat over the border of British America, south to Central America. Accidental in New York and Massachusetts.

378 a. **Speotyto cunicularia floridana** RIDGW. \  
Florida Burrowing Owl.

*Speotyto cunicularia* var. *floridana* RIDGW. Am. Sportsm. V. July 4, 1874, 216.

[B —, C —, R 408a, C 488.]

GEOG. DIST.—Southern Florida; Bahamas.

GENUS **GLAUCIDIUM** BOIE.

*Glaucidium* BOIE, Isis, 1826, 970. Type, *Strix nana* KING.

379. **Glaucidium gnoma** WAGL.  
Pygmy Owl.

*Glaucidium gnoma* WAGL. Isis, 1832, 275.

[B 60, C 329, R 409, C 484.]

GEOG. DIST.—Western North America in mountainous regions from British Columbia to eastern slopes of the Rocky Mountains and south to the tablelands of Mexico.

379 *a*. **Glaucidium gnoma californicum** (SCL.).

California Pygmy Owl.

*Glaucidium californicum* SCL. P. Z. S. 1857, 4.*Glaucidium gnoma californicum* A. O. U. Check-List, 1st Suppl. 1889, 9.[B 60, *part*, C 329, *part*, R 409, *part*, C 484, *part*.]

GEOG. DIST.—Pacific coast region, from California to British Columbia.

379.1. **Glaucidium hoskinsii** (BREWST.).

Hoskins's Pygmy Owl.

*Glaucidium gnoma hoskinsii* BREWST. Auk, V. April, 1888, 136.*Glaucidium hoskinsii* A. O. U. Check-List, 1st Suppl. 1889, 9.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California.

380. **Glaucidium phalænoides** (DAUD.).

Ferruginous Pygmy Owl.

*Strix phalænoides* DAUD. Traité d'Orn. II. 1800, 206.*Glaucidium phalænoides* CAB. J. f. O. 1869, 208.

[B —, C 330, R 410, C 485.]

GEOG. DIST.—Southern border of the United States, from Texas to Arizona, south to southern Brazil.

GENUS **MICROPALLAS** COUES.*Micropallas* COUES, Auk, VI. Jan. 1889, 71. Type, *Athene whitneyi* COOPER.381. **Micropallas whitneyi** (COOPER).

Elf Owl.

*Athene whitneyi* COOPER, Pr. Cal. Ac. Sci. 1861, 118.*Micropallas whitneyi* SENNETT, Auk, VI. July, 1889, 276.

[B —, C 331, R 411, C 486.]

GEOG. DIST.—Southern and Lower California, Arizona, New Mexico, and southern Texas, south to Puebla, Mexico.

ORDER PSITTACI. PARROTS, MACAWS, PAR-  
OQUETS, ETC.

FAMILY **PSITTACIDÆ**. PARROTS AND PAROQUETS.

GENUS **CONURUS** KUHL.

*Conurus* KUHL, Consp. Psitt. 1820, 4. Type, *Psittacus carolinensis* LINN.

382. **Conurus carolinensis** (LINN.).  
Carolina Paroquet.

*Psittacus carolinensis* LINN. S. N. ed. 10, I. 1758, 97.

*Conurus carolinensis* LESS. Traité, 1831, 211.

[B 63, C 315, R 392, C 460.]

GEOG. DIST.—Formerly Florida and the Gulf States north to Maryland, the Great Lakes, Iowa, and Nebraska, west to Colorado, the Indian Territory, and Texas, and straggling northeastward to Pennsylvania and New York. Now restricted to Florida, Arkansas, and Indian Territory, where it is only of local occurrence.

## ORDER COCCYGES. CUCKOOS, ETC.

## SUBORDER CUCULI. CUCKOOS, ETC.

FAMILY **CUCULIDÆ**. CUCKOOS, ANIS, ETC.SUBFAMILY **CROTOPHAGINÆ**. ANIS.GENUS **CROTOPHAGA** LINNÆUS.

*Crotophaga* LINN. S. N. ed. 10, I. 1758, 105. Type, *C. ani* LINN.

[383.] **Crotophaga ani** LINN.

Ani.

*Crotophaga ani* LINN. S. N. ed. 10, I. 1758, 105.

[B 66, 67, C 288, R 389, C 425.]

GEOG. DIST.—West Indies and eastern South America. Rare or casual in southern Florida and Louisiana, and accidental near Philadelphia.

384. **Crotophaga sulcirostris** SWAINS.

Groove-billed Ani.

*Crotophaga sulcirostris* SWAINS. Philos. Mag. I. 1827, 440.

[B—, C—, R 390, C 426.]

GEOG. DIST.—Mexico, north to the southern part of the peninsula of Lower California, and valley of the Lower Rio Grande in Texas, south to Peru.

SUBFAMILY **COCCYGINÆ**. AMERICAN CUCKOOS.GENUS **GEOCOCCYX** WAGLER.

*Geococcyx* WAGLER, Isis, 1831, 524. Type, *G. variegata* WAGL.  
= *Saurothera californiana* LESS.

385. **Geococcyx californianus** (LESS.).

Road-runner.

*Saurothera californiana* LESSON, Compl. Buff. VI. 1829, 420.

*Geococcyx californianus* BAIRD, B. N. Am. 1858, 73.

[B 68, C 289, R 385, C 427.]

GEOG. DIST.—Northern and central Mexico, north to head of Sacramento Valley in California, southern Utah, southeastern Colorado, southwestern Kansas, western Oklahoma, and western Texas.

GENUS **COCYZUS** VIEILLOT.

*Coccyzus* VIEILL. Analyse, 1816, 28. Type, *Cuculus americanus* LINN.

386. **Coccyzus minor** (GMEL.).

Mangrove Cuckoo.

*Cuculus minor* GMEL. S. N. I. i. 1788, 411.

*Coccyzus minor* CAB. J. f. O. 1856, 104.

[B 71, C 292, R 386, C 429.]

GEOG. DIST.—Key West, Florida, Louisiana, the West Indies, and Central America to northeastern South America.

[386 a.] **Coccyzus minor maynardi** (RIDGW.).

Maynard's Cuckoo.

*Coccyzus maynardi* RIDGW. Man. N. Am. B. 1887, 274.

*Coccyzus minor maynardi* A. O. U. Check-List, 1st Suppl. 1889, 9.

[B 71, *part*, C 292, *part*, R 386, *part*, C 429, *part*.]

GEOG. DIST.—Bahamas and Florida Keys. Cuba?

387. **Coccyzus americanus** (LINN.).

Yellow-billed Cuckoo.

*Cuculus americanus* LINN. S. N. ed. 10, I. 1758, 111.

*Coccyzus americanus* BONAP. Journ. Ac. Nat. Sci. Phila. III. ii. 1824, 367.



[B 69, C 291, R 387, C 429.]

GEOG. DIST.—Eastern temperate North America, breeding from Florida north to New Brunswick, Canada, and Minnesota, west to the eastern border of the Plains, and south, in winter, to Costa Rica and the West Indies.

387 *a.* **Coccyzus americanus occidentalis** RIDGW.

**California Cuckoo.**

*Coccyzus americanus occidentalis* RIDGW. Man. N. Am. B. 1887, 273.

[B—, C—, R 387, *part*, C 429, *part*.]

GEOG. DIST.—Western temperate North America, from northern Lower California north to southern British Columbia, east to New Mexico and western Texas, and south over tablelands of Mexico.

388. **Coccyzus erythrophthalmus** (WILS.).

**Black-billed Cuckoo.**

*Cuculus erythrophthalmus* WILS. Am. Orn. IV. 1811, 16, pl. 28.

*Coccyzus erythrophthalmus* BONAP. Journ. Ac. Nat. Sci. Phila. III. ii. 1824, 367.

[B 70, C 290, R 388, C 428.]

GEOG. DIST.—Eastern North America, west to the Rocky Mountains, breeding north to Labrador, Manitoba, and eastern Assiniboia; south, in winter, to the West Indies and the valley of the Amazon. Accidental in the British Islands and Italy.

GENUS **CUCULUS** LINNÆUS.

*Cuculus* LINN. S. N. ed. 10, I. 1758, 110. Type, *C. canorus* LINN.

[388.1.] **Cuculus canorus telephonus** (HEINE).

**Kamchatkan Cuckoo.**

*Cuculus telephonus* HEINE, J. f. O. 1863, 352.

*Cuculus canorus telephonus* STEJN. Bull. 29, U. S. Nat. Mus. 1885, 224.

[B—, C—, R—, C—.]

GEOG. DIST.—An Asiatic species, occurring accidentally on the Pribilof Islands.

## SUBORDER TROGONES. TROGONS.

FAMILY **TROGONIDÆ**. TROGONS.GENUS **TROGON** LINNÆUS.

*Trogon* LINN. S. N. ed. 12, I. 1766, 167. Type, *T. viridis* LINN.

389. **Trogon ambiguus** GOULD.

Coppery-tailed Trogon.

*Trogon ambiguus* GOULD, P. Z. S. 1835, 30.

[B 65, C 284, R 384, C 422.]

GEOG. DIST.—Southern and central Mexico, from Oaxaca and Guerrero, north to the valley of the Lower Rio Grande in Texas, and southern Arizona.

## SUBORDER ALCYONES. KINGFISHERS.

FAMILY **ALCEDINIDÆ**. KINGFISHERS.GENUS **CERYLE** BOIE.

*Ceryle* BOIE, Isis, 1828, 316. Type, *Alcedo rudis* LINN.

SUBGENUS **STREPTOCERYLE** BONAPARTE.

*Streptoceryle* BONAP. Consp. Vol. Anisod. 1854. 10. Type, *Alcedo torquata* LINN.

390. **Ceryle alcyon** (LINN.).

Belted Kingfisher.

*Alcedo alcyon* LINN. S. N. ed. 10, I. 1758, 115.

*Ceryle alcyon* BONAP. P. Z. S. for 1837 (1838), 108.

[B 117, C 286, R 382, C 423.]

GEOG. DIST.—North America, from the Arctic Ocean south to Panama and the West Indies. Breeds from the southern border of the United States northward.

[390.1.] **Ceryle torquata** (LINN.).**Ringed Kingfisher.***Alcedo torquata* LINN. Syst. Nat. ed. 12, I. 1766, 180.*Ceryle torquata* BOIE, Isis, 1828, 316.

[B—, C—, R—, C—.]

GEOG. DIST.—Mexico and southward to southern South America. Casual on the Lower Rio Grande, Texas.

SUBGENUS **CHLOROCERYLE** KAUP.

*Chloroceryle* KAUP, Fam. Eisv. 1848, 8. Type, *Alcedo superciliosa* LINN.

391. **Ceryle americana septentrionalis** SHARPE.**Texas Kingfisher.**

*Ceryle americana septentrionalis* SHARPE, Cat. Birds Brit. Mus. XVII, 1892, 134.

[B 118, C 287, R 383, C 424.]

GEOG. DIST.—Southern Texas, south to Panama.

## ORDER PICI. WOODPECKERS, WRYNECKS, ETC.

FAMILY **PICIDÆ**. WOODPECKERS.GENUS **CAMPEPHILUS** GRAY.

*Campephilus* GRAY, List Gen. B. 1840, 54. Type, *Picus principalis* LINN.

392. **Campephilus principalis** (LINN.).

Ivory-billed Woodpecker.

*Picus principalis* LINN. S. N. ed. 10, I. 1758, 113.*Campephilus principalis* GRAY, List Gen. B. 1840, 54.

[B 72, C 293, R 359, C 431.]

GEOG. DIST.—Formerly South Atlantic and Gulf States, from North Carolina to Texas, north in the Mississippi Valley to Missouri, southern Illinois, and southern Indiana. Now restricted to the Gulf States and the Lower Mississippi Valley, where only locally distributed.

GENUS **DRYOBATES** BOIE.*Dryobates* BOIE, Isis, 1826, 977. Type, *Picus pubescens* LINN.393. **Dryobates villosus** (LINN.).

Hairy Woodpecker.

*Picus villosus* LINN. S. N. ed. 12, I. 1766, 175.*Dryobates villosus* CABANIS, Mus. Hein. IV. ii. June 15, 1863, 66.[B 74, *part*, C 298, *part*, R 360, C 438, *part*.]

GEOG. DIST.—Northern and middle portions of the eastern United States, from the Atlantic coast to the Great Plains.

393 a. **Dryobates villosus leucomelas** (BODD.).

Northern Hairy Woodpecker.

*Picus leucomelas* BODDAERT, Tabl. Pl. Enl. 1783, 21.*Dryobates villosus leucomelas* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.[B 74, *part*, C 298, *part*, R 360a, C 438, *part*.]

GEOG. DIST.—Northern North America, south to about the northern border of the United States.

393 b. **Dryobates villosus audubonii** (SWAINS.).

Southern Hairy Woodpecker.

*Picus audubonii* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831, 306.*Dryobates villosus audubonii* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 74, *part*, C 298, *part*, R 360, *part*, C 438, *part*.]

GEOG. DIST.—South Atlantic and Gulf States, north to North Carolina and Tennessee, west to Louisiana and southeastern Texas.

393 *c.* **Dryobates villosus harrisii** (AUD.).

Harris's Woodpecker.

*Picus harrisii* AUD. Orn. Biog. V. 1839, 191.

*Dryobates villosus harrisii* RIDGW. Pr. U. S. Nat. Mus. VIII.  
1885, 355.

[B 75, C 298*a*, R 360*b*, C 439.]

GEOG. DIST.—Pacific coast of North America, from northern California to southern Alaska.

393 *d.* **Dryobates villosus hyloscopus** (CAB.).

Cabanis's Woodpecker.

*Dryobates hyloscopus* CAB. & HEINE, Mus. Hein. IV. ii. 1863, 69.

[*Dryobates villosus*] *hyloscopus* BREWST. Auk, V. July, 1888, 252  
(in text).

[B 75, *part*, C 298*a*, *part*, R 360*b*, *part*, C 439, *part*.]

GEOG. DIST.—Western United States, except northwest coast, and south into Mexico.

394. **Dryobates pubescens** (LINN.).

Downy Woodpecker.

*Picus pubescens* LINN. S. N. ed. 12, I. 1766, 175.

*Dryobates pubescens* CABANIS, Mus. Hein. IV. ii. June 15, 1863, 62.

[B 76, C 299, R 361, C 440.]

GEOG. DIST.—Northern and eastern North America, west to British Columbia and the eastern edge of the Plains; south to the Gulf of Mexico.

394 *a.* **Dryobates pubescens gairdnerii** (AUD.).

Gairdner's Woodpecker.

*Picus gairdnerii* AUD. Orn. Biog. V. 1839, 317.

*Dryobates pubescens gairdnerii* RIDGW. Pr. U. S. Nat. Mus. VIII.  
1885, 355.

[B 77, *part*, C 299*a*, *part*, R 361*a*, *part*, C 441, *part*.]

GEOG. DIST.—Pacific coast of the United States, north to British Columbia (lat. 55°).

394 *b*. **Dryobates pubescens oreæcus** BATCH.

Batchelder's Woodpecker.

*Dryobates pubescens oreæcus* BATCH. AUK, VI. July, 1889, 253.

[B 77, *part*, C 299*a*, *part*, R 361*a*, *part*, C 441, *part*.]

GEOG. DIST.—Rocky Mountain region of the United States.

395. **Dryobates borealis** (VIEILL.).

Red-cockaded Woodpecker.

*Picus borealis* VIEILL. Ois. Am. Sept. II. 1807, 66.

*Dryobates borealis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 80, C 296, R 362, C 433.]

GEOG. DIST.—Southeastern United States, from North Carolina, Tennessee and Indian Territory, south to eastern Texas and the Gulf coast.

396. **Dryobates scalaris bairdi** (MALHERBE).

Texan Woodpecker.

*Picus bairdi* (SCL. MSS.) MALHERBE, Mon. Pic. I. 1861, 118,  
pl. 27.

*Dryobates scalaris bairdi* RIDGW. Man. N. Am. B. 1887, 285.

[B 79, C 297, R 363, C 434.]

GEOG. DIST.—Southern border of the United States, from Texas to California, north from western Arizona and southern Nevada to southwestern Utah, south to the tablelands of Mexico.

396 *a*. **Dryobates scalaris lucasanus** (XANTUS).

Saint Lucas Woodpecker.

*Picus lucasanus* XANTUS, Pr. Ac. Nat. Sci. Phila. 1859, 298.

*Dryobates scalaris lucasanus* RIDGW. Pr. U. S. Nat. Mus. VIII.  
1885, 355.

[B—, C 297*b*, R 363*a*, C 436.]

GEOG. DIST.—Lower California, north to Lat. 34° in the Colorado Desert, California.



397. **Dryobates nuttallii** (GAMB.).

Nuttall's Woodpecker.

*Picus nuttallii* GAMBEL, Pr. Ac. Nat. Sci. Phila. April, 1843, 259.*Dryobates nuttallii* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 78, C 297a, R 364, C 435.]

GEOG. DIST.—Southern Oregon, California, and northern Lower California.

398. **Dryobates arizonæ** (HARGITT).

Arizona Woodpecker.

*Picus arizonæ* HARGITT, Ibis, April, 1886, 115.*Dryobates arizonæ* RIDGW. Man. N. Am. B. 1887, 286.

[B —, C —, R 365, C 437.]

GEOG. DIST.—Southern Arizona, south into northern Mexico.

GENUS **XENOPICUS** BAIRD.*Xenopicus* BAIRD, B. N. Am. 1858, 83. Type, *Leuconerpes albolarvatus* CASS.399. **Xenopicus albolarvatus** (CASS.).

White-headed Woodpecker.

*Leuconerpes albolarvatus* CASSIN, Pr. Ac. Nat. Sci. Phila. Oct. 1850, 106.*Xenopicus albolarvatus* MALHERBE, Mon. Pic. II. 1862, 221.

[B 81, C 295, R 366, C 452.]

GEOG. DIST.—Mountains of the Pacific coast, from southern British Columbia to southern California (including the eastern slope of the Sierra Nevada), and east to the Blue Mountains of Oregon and west-central Idaho.

GENUS **PICOIDES** LACÉPÈDE.*Picoides* LACÉPÈDE, Mém. de l'Inst. III. 1801, 509. Type *Picus tridactylus* LINN.

400. **Picoides arcticus** (SWAINS.).

Arctic Three-toed Woodpecker.

*Picus (Apternus) arcticus* SWAINS. in SW. & RICH. Fauna Bor.  
Am. II. 1831, 313.*Picoides arcticus* GRAY, Gen. B. I. 1845, 434.

[B 82, C 300, R 367, C 443.]

GEOG. DIST.—Northern North America, from the arctic regions south to the northern United States (New England, New York, Michigan, Minnesota, and Idaho), and in the Sierra Nevada to Lake Tahoe.

401. **Picoides americanus** BREHM.

American Three-toed Woodpecker.

*Picoides americanus* BREHM, Handb. Vög. Deutschl. 1831, 195.

[B 83, C 301, R 368, C 444.]

GEOG. DIST.—Northern North America, east of the Rocky Mountains, from the arctic regions southward to the northern United States (Maine, Mass., New York).

401 a. **Picoides americanus alascensis** (NELS.).

Alaskan Three-toed Woodpecker.

*Picoides tridactylus alascensis* NELSON, Auk, I. April, 1884, 165.*Picoides americanus alascensis* RIDGW. Pr. U. S. Nat. Mus. VIII.  
1885, 355.

[B —, C —, R —, C —.]

GEOG. DIST.—Alaska, south to northern Washington.

401 b. **Picoides americanus dorsalis** BAIRD.

Alpine Three-toed Woodpecker.

*Picoides dorsalis* BAIRD, B. N. Am. 1858, 100.*Picoides americanus dorsalis* BAIRD, Orn. Calif. I. 1870, 386.

[B 84, C 301a, R 368a, C 445.]

GEOG. DIST.—Rocky Mountain region, from British Columbia and Idaho south into New Mexico.

GENUS **SPHYRAPICUS** BAIRD.

*Sphyrapicus* BAIRD, B. N. Am. 1858, 80, 101. Type, *Picus varius* LINN.

402. **Sphyrapicus varius** (LINN.).

Yellow-bellied Sapsucker.

*Picus varius* LINN. S. N. ed. 12, I. 1766, 176.

*Sphyrapicus varius* BAIRD, B. N. Am. 1858, 103.

[B 85, C 302, R 369, C 446.]

GEOG. DIST.— Eastern North America north to about Lat. 63° 30' (north of Fort Simpson), breeding from Massachusetts northward; south, in winter, to the West Indies, Mexico and Costa Rica.

402 a. **Sphyrapicus varius nuchalis** BAIRD.

Red-naped Sapsucker.

*Sphyrapicus varius* var. *nuchalis* BAIRD, B. N. Am. 1858, 103.

[B 86, C 302a, R 369a, C 447.]

GEOG. DIST.— Rocky Mountain region, from British Columbia to the Sierra Bolaños, Jalisco, Mexico, and the Cape region of Lower California; and westward across the Great Basin to the eastern slope of the Sierra Nevada in northern California.

403. **Sphyrapicus ruber** (GMEL.).

Red-breasted Sapsucker.

*Picus ruber* GMEL. S. N. I. 1788, 429.

*Sphyrapicus ruber* BAIRD, B. N. Am. 1858, 104.

[B 87, C 302b, 303? R 369b, C 448.]

GEOG. DIST.— Pacific coast region, from British Columbia south in the mountains and foothills of California to northern Lower California.

404. **Sphyrapicus thyroideus** (CASS.).

Williamson's Sapsucker.

*Picus thyroideus* CASSIN, Pr. Ac. Nat. Sci. Phila. 1850-1851, 349.

*Sphyrapicus thyroideus* BAIRD, B. N. Am. 1858, 106.

[B 88, 89, C 304, 305, R 370, C 449.]

GEOG. DIST.—Rocky Mountain region of the United States, west to the Sierra Nevada, Cascades, and northern Coast Ranges; south on the tablelands of Mexico to the Sierra Bolaños, Jalisco.

GENUS **CEOPHLÆUS** CABANIS.

*Ceophlæus* CABANIS, J. f. O. 1862, 176. Type, *Picus pileatus* LINN.

405. **Ceophlæus pileatus** (LINN.).

**Pileated Woodpecker.**

*Picus pileatus* LINN. S. N. ed. 10, I. 1758, 113.

*C[ceophlæus] pileatus* CABANIS, J. f. O. 1862, 176.

[B 90, C 294, R 371, C 432.]

GEOG. DIST.—Formerly the heavily wooded region of North America south of about Lat. 63°, except in the southern Rocky Mountains; now rare or extirpated in the more thickly settled parts of the Eastern States.

GENUS **MELANERPES** SWAINSON.

SUBGENUS **MELANERPES.**

*Melanerpes* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831, 316.

Type, *Picus erythrocephalus* LINN.

406. **Melanerpes erythrocephalus** (LINN.).

**Red-headed Woodpecker.**

*Picus erythrocephalus* LINN. S. N. ed. 10, I. 1758, 113.

*Melanerpes erythrocephalus* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831, 316.

[B 94, C 309, R 375, C 453.]

GEOG. DIST.—United States, west to the Rocky Mountains, and north from Florida to about Lat. 50°, straggling westward to Salt Lake Valley and Arizona; rare or local east of the Hudson River.

407. **Melanerpes formicivorus bairdi** RIDGW.  
**Californian Woodpecker.**

*Melanerpes formicivorus bairdi* RIDGW. Bull. No. 21 U. S. Nat.  
 Mus. 1881, 34, 85.

[B 95, C 310, R, 377, C 454.]

GEOG. DIST.—Pacific coast region of the United States, from southern Oregon south to northern Lower California and Mexico, east through Arizona to southern New Mexico and western Texas.

- 407 a. **Melanerpes formicivorus angustifrons** BAIRD.  
**Narrow-fronted Woodpecker.**

*Melanerpes formicivorus* var. *angustifrons* BAIRD, Orn. Cal. I. 1870,  
 405.

[B —, C 310a, R 377a, C 455.]

GEOG. DIST.—Southern Lower California.

SUBGENUS **ASYNDESMUS** COUES.

*Asyndesmus* COUES, Pr. Ac. Nat. Sci. Phila. 1866, 55. Type,  
*Picus torquatus* WILS.

408. **Melanerpes torquatus** (WILS.).  
**Lewis's Woodpecker.**

*Picus torquatus* WILSON, Am. Orn. III. 1811, 31, pl. xx. fig. 3.

*Melanerpes torquatus* BONAP. Geog. & Comp. List, 1838, 40.

[B 96, C 311, R 376, C 456.]

GEOG. DIST.—Western United States, from the Black Hills and the Rocky Mountains to the Pacific, and from southern British Columbia and southern Alberta south to Arizona, and (in winter) western Texas. Casual in Kansas.

SUBGENUS **CENTURUS** SWAINSON.

*Centurus* SWAINS. Classif. B. II. 1837, 310. Type, *Picus carolinus* LINN.

409. **Melanerpes carolinus** (LINN.).

Red-bellied Woodpecker.

*Picus carolinus* LINN. S. N. ed. 10. I. 1758, 113.*Melanerpes carolinus* RIDGW. Ann. Lyc. N. Y. X. Jan. 1874, 378.

[B 91, C 306, R 372, C 450.]

GEOG. DIST.—Eastern and southern United States, north casually to Massachusetts, New York, Ontario, southern Michigan, and central Iowa, west to eastern Nebraska, eastern Kansas, Indian Territory and Texas.

410. **Melanerpes aurifrons** (WAGL.).

Golden-fronted Woodpecker.

*Picus aurifrons* WAGLER, Isis, 1829, 512.*Melanerpes aurifrons* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 92, C 307, R 373, C 451.]

GEOG. DIST.—Central and southern Texas, southward over the tablelands of Mexico to the City of Mexico, and from southern Tamaulipas west to Aguas Calientes and Jalisco.

411. **Melanerpes uropygialis** (BAIRD).

Gila Woodpecker.

*Centurus uropygialis* BAIRD, Pr. Ac. Nat. Sci. Phila. June, 1854, 120.*Melanerpes uropygialis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 93, C 308, R 374, C 452.]

GEOG. DIST.—Southwestern New Mexico, and southern and western Arizona, from Lat. 35° southward through Lower California to Cape St. Lucas, and through western Mexico to Aguas Calientes and Jalisco.

GENUS **COLAPTES** SWAINSON.*Colaptes* SWAINS. Zool. Journ. III. Dec. 1827, 353. Type,  
*Cuculus auratus* LINN.412. **Colaptes auratus** (LINN.).

Flicker.

*Cuculus auratus* LINN. S. N. ed. 10, I. 1758, 112.*Colaptes auratus* VIGORS, Zool. Journ. III. 1827, 444.



[B 97, C 312, R 378, C 457.]

GEOG. DIST.—Northern and eastern North America, west to the eastern slope of the Rocky Mountains and Alaska. Occasional on the Pacific slope, from California northward. Accidental in Europe.

413. **Colaptes cafer** (GMEL.).

**Red-shafted Flicker.**

*Picus cafer* GMEL. S. N. I. 1788, 431.

*Colaptes cafer* STEJN. Stand. Nat. Hist. IV. 1885, 428.

[B 98, C 314, R 378b, C 459.]

GEOG. DIST.—Rocky Mountain region, from British Columbia south to southern Mexico, and west to the Coast Ranges in Oregon and Washington, and to the Pacific coast from northern California southward.

413 a. **Colaptes cafer saturator** (RIDGW.).

**Northwestern Flicker.**

*Colaptes mexicanus saturator* RIDGW. Pr. Biol. Soc. Wash. II.

April 10, 1884, 90.

*Colaptes cafer saturator* A. O. U. Check-list, 1886, 218.

[B —, C —, R —, C —.]

GEOG. DIST.—Northwest coast, from northern California to Sitka.

414. **Colaptes chrysoides** (MALH.).

**Gilded Flicker.**

*Geopicus chrysoides* MALH. Rev. et Mag. Zool. IV. 1852, 553.

*Colaptes chrysoides* REICH. Handb. Spec. Ornith. Scansoriæ, 1854, 413.

[B 99, C 313, R 379, C 458.]

GEOG. DIST.—Central and southern Arizona, from Lat. 34° to southern Sonora, and Lower California south of Lat. 30°.

415. **Colaptes rufipileus** RIDGW.

**Guadalupe Flicker.**

*Colaptes mexicanus rufipileus* RIDGW. Bull. U. S. Geol. & Geog. Surv. Terr. II. No. 2, April 1, 1876, 191.

*Colaptes rufipileus* RIDGW. Bull. Nutt. Orn. Club, II. July, 1877, 60.

[B —, C —, R 380, C —.]

GEOG. DIST.—Guadalupe Island, Lower California.

ORDER MACROCHIRES. GOATSUCKERS,  
SWIFTS, ETC.

SUBORDER CAPRIMULGI. GOATSUCKERS, ETC.

FAMILY **CAPRIMULGIDÆ**. GOATSUCKERS, ETC.

GENUS **ANTROSTOMUS** GOULD.

*Antrostomus* GOULD, Icones Avium, 1838. Type, *Caprimulgus carolinensis* GMEL.

416. **Antrostomus carolinensis** (GMEL.).

Chuck-will's-widow.

*Caprimulgus carolinensis* GMEL. S. N. I. ii. 1788, 1028.

*Antrostomus carolinensis* GOULD, Icones Avium, 1838.

[B 111, C 264, R 353, C 396.]

GEOG. DIST.—South Atlantic and Gulf States, from Virginia south through eastern Mexico to Central America ; Cuba. North, in the interior, to southern Illinois and Kansas. Accidental in Massachusetts.

417. **Antrostomus vociferus** (WILS.).

Whip-poor-will.

*Caprimulgus vociferus* WILS. Am. Orn. V. 1812, 71, pl. 41, figs. 1-3.

*Antrostomus vociferus* BONAP. Geog. & Comp. List, 1838, 8.

[B 112, C 265, R 354, C 397.]

GEOG. DIST.—Eastern North America to the Plains, and from Lat. 50° south to Guatemala.

417 a. **Antrostomus vociferus macromystax** (WAGLER).

Stephens's Whip-poor-will.

*Caprimulgus macromystax* WAGLER, Ibis, 1831, 533.*Caprimulgus vociferus macromystax* HARTERT, Ibis, 1892, 286.*Antrostomus vociferus macromystax* A. O. U. Check-List, 6th Suppl. Auk, XI. Jan. 1894, 48.

[B —, C —, R —, C 881.]

GEOG. DIST.—Arizona, New Mexico, and tablelands of Mexico south to Guatemala.

GENUS **PHALÆNOPTILUS** RIDGWAY.*Phalænoptilus* RIDGW. Pr. U. S. Nat. Mus. III. 1880, 5. Type,*Caprimulgus nuttallii* AUD.418. **Phalænoptilus nuttallii** (AUD.).

Poor-will.

*Caprimulgus nuttallii* AUD. B. Am. VII. 1843, 350, pl. 495.*Phalænoptilus nuttalli* RIDGW. Pr. U. S. Nat. Mus. III. 1880, 5.

[B 113, C 266, R 355, C 398.]

GEOG. DIST.—Western United States, from the Sierra Nevada eastward to eastern Nebraska and eastern Kansas, north to central Idaho and Montana, and south to southern Mexico.

418 a. **Phalænoptilus nuttallii nitidus** BREWST.

Frosted Poor-will.

*Phalænoptilus nuttalli nitidus* BREWST. Auk, IV. April, 1877, 147.[B 113, *part*, C 266, *part*, R 355, *part*, C 398, *part*.]

GEOG. DIST.—Texas to Arizona, and north to western Kansas.

418 b. **Phalænoptilus nuttallii californicus** RIDGW.

Dusky Poor-will.

*Phalænoptilus nuttalli californicus* RIDGW. Man. N. Am. B. 1887, 588, foot-note.[B 113, *part*, C 266, *part*, R 355, *part*, C 398, *part*.]

GEOG. DIST.—Coast of California.

GENUS **NYCTIDROMUS** GOULD.

*Nyctidromus* GOULD, Icones Avium, II. 1838, pl. ii. Type,  
*N. derbyanus* GOULD = *Caprimulgus albicollis* GMEL.

419. **Nyctidromus albicollis merrilli** SENN.

Merrill's Parauque.

*Nyctidromus albicollis merrilli* SENN. Auk, V. Jan. 1888, 44.

[B —, C —, R 356, C 395.]

GEOG. DIST.—Valley of the Lower Rio Grande, north to the Nueces River, south into northeastern Mexico.

GENUS **CHORDEILES** SWAINSON.

*Chordeiles* SWAINS. in Sw. & RICH. Fauna Bor. Am. II. 1831, 496. Type, *Caprimulgus virginianus* GMEL.

420. **Chordeiles virginianus** (GMEL.).

Nighthawk.

*Caprimulgus virginianus* GMEL. S. N. I. ii. 1788, 1028.

*Chordeiles virginianus* SWAINS. in Sw. & RICH. Fauna Bor. Am. II. 1831, 496.

[B 114, C 267, R 357, C 399.]

GEOG. DIST.—Northern and eastern North America, west to the Great Plains and central British Columbia, and from Labrador south through tropical America to the Argentine Republic.

420 a. **Chordeiles virginianus henryi** (CASS.).

Western Nighthawk.

*Chordeiles henryi* CASS. Illustr. B. Cal. Tex. etc. I. 1855, 233.

*Chordeiles virginianus* var. *henryi* COUES, Key, 1872, 181.

[B 115, C 267a, R 357a, C 400.]

GEOG. DIST.—Western United States, from the Great Plains to the Pacific coast, and from British Columbia south to northern South America.

420 *b. Chordeiles virginianus chapmani* (COUES).

Florida Nighthawk.

[*Chordeiles popetue*] *chapmani* (SENNET, MS.) COUES, Auk, V.  
Jan. 1888, 37.*Chordeiles virginianus chapmani* SCOTT, Auk, V. April, 1888, 186.[B —, C —, R 357*b*, C 401.]

GEOG. DIST.— Florida and the Gulf coast of Texas, south in winter to South America.

421. *Chordeiles acutipennis texensis* (LAWR.).

Texan Nighthawk.

*Chordeiles texensis* LAWR. Ann. Lyc. N. Y. VI. Dec. 1856, 167.*Chordeiles acutipennis texensis* RIDGW. in BAIRD, BREWER & RIDGWAY, N. Am. Birds, II. 1874, 406.

[B 116, C 268, R 358, C 402.]

GEOG. DIST.— Southern border of the United States, from Texas to southern California, north into southern Utah; south to Cape St. Lucas and Veragua.

## SUBORDER CYPSELI. SWIFTS.

## FAMILY MICROPODIDÆ. SWIFTS.

## SUBFAMILY CHÆTURINÆ. SPINE-TAILED SWIFTS.

## GENUS CYPSELOIDES STREUBEL.

*Cypseloides* STREUBEL, Isis, 1848, 366. Type, *Hemiprocne fumigata* NATT. MS.422. *Cypseloides niger* (GMEL.).

Black Swift.

*Hirundo nigra* GMEL. S. N. I. ii. 1788, 1025.*Cypseloides niger* SCL. P. Z. S. 1865, 615.

[B 108, C 270, R 350, C 404.]

GEOG. DIST.—Rocky Mountain region (Colorado), west to the Pacific coast; north to British Columbia, and south to Lower California, Mexico, Costa Rica, and the West Indies.

GENUS **CHÆTURA** STEPHENS.

*Chætura* STEPH. Gen. Zool. XIII. pt. ii. 1825, 76. Type, *Hirundo pelagica* LINN.

423. **Chætura pelagica** (LINN.).

Chimney Swift.

*Hirundo pelagica* LINN. S. N. ed. 10, I. 1758, 192.

*Chætura pelagica* STEPH. Gen. Zool. XIII. pt. ii. 1825, 76.

[B 109, C 271, R 351, C 405.]

GEOG. DIST.—Eastern North America, north to Labrador and the Fur Countries, west to the Plains, and passing south of the United States in winter at least to Jalapa, Mexico, and Cozumel Island.

424. **Chætura vauxii** (TOWNS.).

Vaux's Swift.

*Cypselus vauxii* TOWNS. Journ. Ac. Nat. Sci. Phila. VIII. 1839, 148.

*Chætura vauxii* DEKAY, Zool. N. Y. II. 1844, 36.

[B 110, C 272, R 352, C 406.]

GEOG. DIST.—Pacific coast of the United States northward to British Columbia; migrating south in winter to Lower California and Mexico.

SUBFAMILY **MICROPODINÆ**. TYPICAL SWIFTS.

GENUS **ÆRONAUTES** HARTERT.

*Æronautes* HARTERT, Cat. Birds Brit. Mus. XVI, 1892, 459.

Type, *Cypselus melanoleucus* BAIRD.



425. **Aëronautes melanoleucus** (BAIRD).

White-throated Swift.

*Cypselus melanoleucus* BAIRD, Pr. Ac. Nat. Sci. Phila. June, 1854,  
118.*Aëronautes melanoleucus* HARTERT, Cat. Birds Brit. Mus. XVI,  
1892, 459.

[B 107, C 269, R 349, C 403.]

GEOG. DIST.—Western United States, from the Black Hills, northern Wyoming, and southern Montana to the Pacific; south, in winter, to Guatemala.

## SUBORDER TROCHILI. HUMMINGBIRDS.

FAMILY **TROCHILIDÆ**. HUMMINGBIRDS.GENUS **EUGENES** GOULD.*Eugenes* GOULD, Mon. Troch. pt. xii. 1856, pl. 59. Type, *Trochilus fulgens* SWAINS.426. **Eugenes fulgens** (SWAINS.).

Rivoli Hummingbird.

*Trochilus fulgens* SWAINS. Phil. Mag. I. 1827, 441.*Eugenes fulgens* GOULD, Mon. Troch. II. 1856, pl. 59.

[B —, C 274 bis, R 334, C 408.]

GEOG. DIST.—Southern Arizona and tablelands of Mexico to Nicaragua.

GENUS **CÆLIGENA** LESSON.*Cæligena* LESS. Ind. & Synop. Gen. Troch. 1832, p. xviii. Type, *Ornismya clemenciæ* LESS.

427. **Cœligena clemenciæ** LESS.

Blue-throated Hummingbird.

*Ornismya clemenciæ* LESS. Ois. Mouch. 1829, 216, pl. 80.*Cœligena clemenciæ* LESS. Ind. & Synop. Gen. Troch. 1832, p. xviii.

[B —, C —, R —, C —.]

GEOG. DIST.—Southern Arizona, and the tablelands of Mexico to Guerrero and Oaxaca.

GENUS **TROCHILUS** LINNÆUS.SUBGENUS **TROCHILUS**.*Trochilus* LINN. S. N. ed. 10, I. 1758, 119. Type, by elimination, *T. colubris* LINN.428. **Trochilus colubris** LINN.

Ruby-throated Hummingbird.

*Trochilus colubris* LINN. S. N. ed. 10, I. 1758, 120.

[B 101, C 275, R 335, C 409.]

GEOG. DIST.—Eastern North America to the Plains, north to the Fur Countries, breeding from Florida to Labrador, and south, in winter, to Cuba, Mexico, and Veragua.

429. **Trochilus alexandri** BOURC. & MULS.

Black-chinned Hummingbird.

*Trochilus alexandri* BOURC. & MULS. Ann. Soc. Agric. Lyons, IX. 1846, 330.

[B 102, C 276, R 336, C 410.]

GEOG. DIST.—Pacific coast region, from California to British Columbia, east to Utah and Arizona, and south to Lower California, the Valley of Mexico, and Guerrero.

429.1. **Trochilus violajugulum** JEFFRIES.

Violet-throated Hummingbird.

*Trochilus violajugulum* JEFFRIES, Auk, V. April, 1888, 168.

[B —, C —, R —, C —.]

GEOG. DIST.—Coast of southern California (Santa Barbara). Known only from the type specimen.

GENUS **CALYPTE** GOULD.

*Calypte* GOULD, Mon. Troch. III. 1856, pl. 134. Type, *Ornismya costæ* BOURC.

430. **Calypte costæ** (BOURC.).

Costa's Hummingbird.

*Ornismya costæ* BOURC. Rev. Zool. 1839, 294.

*Calypte costæ* GOULD, Mon. Troch. III. pt. xi, 1856, pl. 134.

[B 106, C 280, R 337, C 415.]

GEOG. DIST.—Southern California, southern Nevada, Arizona, Lower California, and western Mexico.

431. **Calypte anna** (LESS.).

Anna's Hummingbird.

*Ornismya anna* LESS. Suppl. Ois. Mouch. 1831, 115, pl. vii.

*Calypte annæ* GOULD, Mon. Troch. III. pt. xi, 1856, pl. 135.

[B 105, C 279, R 338, C 414.]

GEOG. DIST.—Central and southern California, southern Arizona, and Lower California, casually to Guadalupe Island.

GENUS **SELASPHORUS** SWAINSON.

*Selasphorus* SWAINS. in Sw. & RICH. Fauna Bor. Am. II. 1831, 324. Type, *Trochilus rufus* GMEL.

[431.1.] **Selasphorus floresii** GOULD.

Flores's Hummingbird.

*Selasphorus floresii* GOULD, Mon. Troch. pt. xxiii. Sept. 1, 1861 (Vol. III. pl. 139).

[B —, C —, R —, C —.]

GEOG. DIST.—Mexico (Bolaños, Jalisco); accidental at San Francisco, California.

432. **Selasphorus platycercus** (SWAINS.).

Broad-tailed Hummingbird.

*Trochilus platycercus* SWAINS. Phil. Mag. I. 1827, 441.

*Selasphorus platycercus* BON. Conspect. Av. I. 1850, 82.

[B 104, C 278, R 339, C 413.]

GEOG. DIST.—Rocky Mountain region (Idaho, Wyoming, and Colorado), west to the Sierra Nevada, and south through Arizona, New Mexico and Mexico to Guatemala.

433. **Selasphorus rufus** (GMEL.).

Rufous Hummingbird.

*Trochilus rufus* GMEL. S. N. I. i. 1788, 497.

*Trochilus (Selasphorus) rufus* SWAINS. in SW. & RICH. Faun. Bor. Am. II. 1831, 324.

[B 103, C 277, R 340, C 411.]

GEOG. DIST.—Western Texas to Montana, west to the Pacific, and north to Mount St. Elias, Alaska; south in winter on the tablelands of Mexico to Vera Cruz and Oaxaca.

434. **Selasphorus alleni** HENSH.

Allen's Hummingbird.

*Selasphorus alleni* HENSH. Bull. Nutt. Orn. Club, II. 1877, 54.

[B —, C —, R 341, C 412.]

GEOG. DIST.—Pacific coast, north to British Columbia, east to southern Arizona.

GENUS **STELLULA** GOULD.

*Stellula* GOULD, Introd. Troch. 1861, 90. — Type, *Trochilus calliope* GOULD.

436. **Stellula calliope** GOULD.

Calliope Hummingbird.

*Trochilus (Calothorax) calliope* GOULD, P. Z. S. 1847, 11.*Stellula calliope* GOULD, Introd. Troch. 1861, 90.

[B —, C 282, R 343, C 417.]

GEOG. DIST.—Mountains of the Pacific slope, from British Columbia to Montana, Nevada, and New Mexico; south to the Valley of Mexico and mountains of Guerrero.

GENUS **CALOTHORAX** GRAY.

*Calothorax* GRAY, Gen. B. 1840, 13. Type, *Cynanthus lucifer* SWAINS.

437. **Calothorax lucifer** (SWAINS.).

Lucifer Hummingbird.

*Cynanthus lucifer* SWAINS. Phil. Mag. I. 1827, 442.

*Calothorax lucifer* GRAY, Gen. B. I. 1844, 110.

[B —, C —, R 344, C 418.]

GEOG. DIST.—Tablelands of Mexico, from Puebla and the Valley of Mexico north to southern Arizona.

GENUS **AMAZILIA** REICHENBACH.

*Amazilia* REICH. Syst. Av. 1849, pl. 39. Type, *Orthorhynchus amazili* LESS.

438. **Amazilia fuscicaudata** (FRASER).

Rieffer's Hummingbird.

*Trochilus fuscicaudatus* FRAS. P. Z. S. 1840, 17.

*Amazilia fuscicaudata* RIDGW. Pr. U. S. Nat. Mus. I. 1878, 118, 147.

[B —, C —, R 345, C 419.]

GEOG. DIST.—Lower Rio Grande Valley in Texas, south through eastern Mexico to Central America and northern South America.

439. **Amazilia cerviniventris** GOULD.

Buff-bellied Hummingbird.

*Amazilius cerviniventris* GOULD, P. Z. S. 1856, 150.

[B —, C —, R 346, C 420.]

GEOG. DIST.—Valley of the Lower Rio Grande and south in eastern Mexico to southern Vera Cruz.

GENUS **BASILINNA** BOIE.

*Basilinna* BOIE, Isis, 1831, 546. Type, *Trochilus leucotis* VIEILL

440. **Basilinna xantusi** (LAWR.).

**Xantus's Hummingbird.**

*Amazilia xantusi* LAWR. Ann. Lyc. N. Y. 1860, 109.

*Basilinna xanthusi* ELLIOT, Class. & Synop. Troch. March, 1879, 227.

[B—, C 273, R 347, C 407.]

GEOG. DIST.—Lower California north to Lat. 29°.

440.1. **Basilinna leucotis** (VIEILL.).

**White-eared Hummingbird.**

*Trochilus leucotis* VIEILL. Nouv. Dict. d'Hist. Nat. XXIII. 1818, 428.

*Basilinna leucotis* BOIE, Isis, 1831, 546.

[B—, C —, R —, C —.]

GEOG. DIST.—Mexico and Nicaragua, north to the Chiricahua Mountains, Arizona.

GENUS **IACHE** ELLIOT.

*Iache* ELLIOT, Class. & Synop. Troch. March, 1879, 234. Type, *Cyananthus latirostris* SWAINS.

441. **Iache latirostris** (SWAINS.).

**Broad-billed Hummingbird.**

*Cyananthus latirostris* SWAINS. Phil. Mag. I. 1827, 441.

*Iache latirostris* ELLIOT, Class. & Synop. Troch. March, 1879, 235.

[B—, C —, R 348, C 421.]

GEOG. DIST.—Southern Arizona and south to the Valley of Mexico.



## ORDER PASSERES. PERCHING BIRDS.

## SUBORDER CLAMATORES. SONGLESS PERCHING BIRDS.

FAMILY **COTINGIDÆ**. THE COTINGAS.GENUS **Platypsaris** SCLATER.

*Platypsaris* SCL. P. Z. S. 1857, 72 (ex BONAP., 1854, = nomen nudum). Type, *Pachyrhamphus latirostris* BONAP.

441.1. **Platypsaris albiventris** (LAWR.).

Xantus's Becard.

*Hadrostromus albiventris* LAWR. Ann. Lyc. N. Y. VIII. 1867, 475.

*Platypsaris albiventris* RIDGW. Man. N. Am. B. 1887, 325.

[B—, C—, R—, C—.]

GEOG. DIST.—Western and southern Mexico, north to southern Arizona (Huachuca Mts.).

FAMILY **TYRANNIDÆ**. TYRANT FLYCATCHERS.GENUS **MILVULUS** SWAINSON.

*Milvulus* SWAINSON, Zool. Journ. III. July, 1827, 165. Type, *Tyrannus savanna* VIEILL. = *Muscicapa tyrannus* LINN.

[442.] **Milvulus tyrannus** (LINN.).

Fork-tailed Flycatcher.

*Muscicapa tyrannus* LINN. S. N. ed. 12, I. 1766, 325.

*Milvulus tyrannus* BONAP. Geogr. & Comp. List, 1838, 25.

[B 122, C 240, R 302, C 366.]

GEOG. DIST.—Southern Mexico and southward throughout Central and most of South America. Accidental in the United States (Mississippi, Kentucky, New Jersey, southern California).

443. **Milvulus forficatus** (GMEL.).

Scissor-tailed Flycatcher.

*Muscicapa forficata* GMEL. S. N. I. ii. 1788, 931.*Milvulus forficatus* SWAINS. Classif. B. II. 1827, 225.

[B 123, C 241, R 301, C 367.]

GEOG. DIST.—Texas, Oklahoma, Indian Territory, southern Kansas and southwestern Missouri, south through eastern Mexico to Costa Rica. Accidental in southern Florida (Key West), New Jersey, New England, and at York Factory, Hudson Bay.

GENUS **TYRANNUS** CUVIER.

*Tyrannus* CUVIER, Leç. d'An. Comp. I. 1799, tabl. ii. (Cf. Tabl. Elem. 1797, p. 201.) Type, *Lanius tyrannus* LINN.

444. **Tyrannus tyrannus** (LINN.).

Kingbird.

*Lanius tyrannus* LINN. S. N. ed. 10, I. 1758, 94.*Tyrannus tyrannus* JORDAN, Man. Vert. ed. 4, 1884, 96.

[B 124, C 242, R 304, C 368.]

GEOG. DIST.—North America, from the British Provinces south, in winter, through eastern Mexico, Central and South America. Less common west of the Rocky Mountains. Not recorded from New Mexico and Arizona.

445. **Tyrannus dominicensis** (GMEL.).

Gray Kingbird.

*Lanius tyrannus*  $\beta$ . *dominicensis* GMEL. S. N. I. 1788, 302.

*Tyrannus dominicensis* RICHARDSON, Rep. Sixth Meet. Brit. Ass. V. 1837, 170.

[B 125, C 243, R 303, C 369.]

GEOG. DIST.—South Atlantic States (South Carolina, Georgia, Florida), West Indies, Atlantic coast of Central America, and northern South America. Accidental in Massachusetts.

446. **Tyrannus melancholicus couchii** (BAIRD).

Couch's Kingbird.

*Tyrannus couchii* BAIRD, B. N. Am. 1858, 175.*Tyrannus melancholicus* var. *couchii* COUES, Check-list, ed. 1, Dec. 1873, 51.

[B 128, 129, C 246, R 305, C 372.]

GEOG. DIST.—Southern border of the United States (Texas), south to Guatemala.

447. **Tyrannus verticalis** SAY.

Arkansas Kingbird.

*Tyrannus verticalis* SAY, LONG'S Exp. II. 1823, 60.

[B 126, C 244, R 306, C 370.]

GEOG. DIST.—Western United States, from the Plains to the Pacific, and from British Columbia south through Lower California and western Mexico to Guatemala. Accidental in Maryland, New Jersey, New York, and Maine.

448. **Tyrannus vociferans** SWAINS.

Cassin's Kingbird.

*Tyrannus vociferans* SWAINS. Quart. Jour. Sci. XX. 1826, 273.

[B 127, C 245, R 307, C 371.]

GEOG. DIST.—Western United States from southern Wyoming, eastern Colorado, New Mexico, Arizona, western Texas and southern California, south to Guatemala.

GENUS **PITANGUS** SWAINSON.

*Pitangus* SWAINSON, Zool. Journ. III. July, 1827, 165. Type,  
*Tyrannus sulphuratus* VIEILL.

449. **Pitangus derbianus** (KAUP).

Derby Flycatcher.

*Saurophagus derbianus* KAUP, P. Z. S. 1851 (Oct. 1852), 44, pl. xxxvi.*Pitangus derbianus* SCLATER, P. Z. S. 1856 (Jan. 1857), 297.

[B—, C—, R 308, C 364.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, south to northern South America.

GENUS **MYIOZETETES** SCLATER.*Myiozetetes* SCL. P. Z. S. 1859, 46. Type, *Muscicapa cayennensis* LINN.[450.] **Myiozetetes texensis** (GIRAUD).

Giraud's Flycatcher.

*Muscicapa texensis* GIRAUD, Sixteen Sp. Texas B. 1841, pl. 1.*Myiozetetes texensis* SCL. P. Z. S. 1859, 56.

[B—, C—, R 309, C—.]

GEOG. DIST.—"Texas" (GIRAUD), south to Central America and northern South America.

GENUS **MYIODYNASTES** BONAPARTE.*Myiodynastes* BONAP. Bull. Soc. Linn. Normandée, II. 1857, 35 (ex BONAP. Comptes Rend. XXXVIII. 1854, 657, nomen nudum). Type, *Muscicapa audax* GMEL.451. **Myiodynastes luteiventris** SCL.

Sulphur-bellied Flycatcher.

*Myiodynastes luteiventris* SCL. P. Z. S. 1859, 42 (ex BONAP. Comptes Rend. XXXVIII. 1854, 657, nomen nudum).

[B—, C—, R 310, C 365.]

GEOG. DIST.—Southern Arizona, south to Panama.

GENUS **MYIARCHUS** CABANIS.*Myiarchus* CAB. Arch. f. Naturg. 1844, i, 272. Type, *Muscicapa ferox* GMEL.

452. **Myiarchus crinitus** (LINN.).

Crested Flycatcher.

*Muscicapa crinita* LINN. S. N. ed. 12, I. 1766, 325.*Myiarchus crinitus* LICHT. Nomencl. Mus. Berol. 1854, 16.

[B 130, C 247, R 312, C 373.]

GEOG. DIST.—Eastern United States and southern Canada, west to Manitoba and the Plains, south through eastern Mexico to Costa Rica, Panama and Colombia. Breeds from Florida northward.

453. **Myiarchus mexicanus** (KAUP).

Mexican Crested Flycatcher.

*Tyrannula mexicana* KAUP, P. Z. S. 1851, 51.*Myiarchus mexicanus* BAIRD, B. N. Am. 1858, 179.

[B 132, C —, R 311, C 374.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, southward to Guatemala.

453 a. **Myiarchus mexicanus magister** RIDGW.

Arizona Crested Flycatcher.

*Myiarchus mexicanus magister* RIDGW. Pr. Biol. Soc. Wash. II.

April 10, 1884, 90.

[B —, C —, R —, C —.]

GEOG. DIST.—Southern Arizona, south into western Mexico.

454. **Myiarchus cinerascens** (LAWR.).

Ash-throated Flycatcher.

*Tyrannula cinerascens* LAWR. Ann. Lyc. N. Y. V. 1851, 121.*Myiarchus cinerascens* SCL. & SALV. Ibis, I. Apr. 1859, 121.

[B 131, C 248, R 313, C 375.]

GEOG. DIST.—Western United States, north to Oregon, Nevada, Utah, and Colorado, south to Guatemala.

454 a. **Myiarchus cinerascens nuttingi** (RIDGW.).

Nutting's Flycatcher.

*Myiarchus nuttingi* RIDGW. Pr. U. S. Nat. Mus. V. 1882, 394.*Myiarchus cinerascens nuttingi* ALLEN, Bull. Am. Mus. Nat. Hist.

IV. Dec. 1892, 346.

[B 131, *part*, C 248, *part*, R 313, *part*, C 375, *part*.]

GEOG. DIST.—Southern Arizona, southward through western Mexico to Costa Rica.

[455.] **Myiarchus lawrenceii** (GIRAUD).

Lawrence's Flycatcher.

*Muscicapa lawrenceii* GIRAUD, Sixteen Sp. Texas B. 1841, 9 (by actual counting, the text not being paged).

*Myiarchus lawrencii* BAIRD, B. N. Am. 1858, 181.

[B 133, C 248, R 314, C 376.]

GEOG. DIST.—"Texas" (GIRAUD) and eastern Mexico.

455 a. **Myiarchus lawrencei olivascens** RIDGW.

Olivaceous Flycatcher.

*Myiarchus lawrencei olivascens* RIDGW. Pr. Biol. Soc. Wash. II.

April 10, 1884, 91.

[B —, C —, R —, C —.]

GEOG. DIST.—Arizona and western Mexico. Casual at Fort Lyon, Colorado.

GENUS **SAYORNIS** BONAPARTE.

*Sayornis* BONAP. Coll. Delattre, 1854, 87. Type, *Tyrannula nigricans* SWAINS.

456. **Sayornis phœbe** (LATH.).

Phœbe.

*Muscicapa phæbe* LATHAM, Ind. Orn. II. 1790, 489.

*Sayornis phæbe* STEJN. Auk, II. Jan. 1885, 51.

[B 135, C 252, R 315, C 379.]

GEOG. DIST.—Eastern North America, west to eastern Colorado and western Texas, and from the British Provinces south to eastern Mexico and Cuba, wintering from the South Atlantic and Gulf States southward. Breeds from South Carolina northward.



457. **Sayornis saya** (BONAP.).

Say's Phoebe.

*Muscicapa saya* BONAP. Am. Orn. I. 1825, 20.*Sayornis sayus* BAIRD, B. N. Am. 1858, 185.

[B 136, C 250, R 316, C 377.]

GEOG. DIST.—Western United States, from the Plains (central North and South Dakota, Nebraska and Kansas) to the Pacific; north along the Yukon River to the Arctic Circle; south to Cape St. Lucas and over the Mexican Plateau to Puebla and central Vera Cruz. Accidental in Massachusetts.

458. **Sayornis nigricans** (SWAINS.).

Black Phoebe.

*Tyrannula nigricans* SWAINS. Phil. Mag. I. May, 1827, 367.*Sayornis nigricans* BONAP. Coll. Delattre, 1854, 87.

[B 134, C 251, R 317, C 378.]

GEOG. DIST.—Southwestern United States, from Texas through southern New Mexico and Arizona to California, and northward along the Pacific coast to Washington; south to Cape St. Lucas, Lower California, and in Mexico to Oaxaca.

GENUS **CONTOPUS** CABANIS.

*Contopus* CAB. J. f. O. III. Nov. 1855, 479. Type, *Muscicapa virens* LINN.

SUBGENUS **NUTTALLORNIS** RIDGW.

*Nuttallornis* RIDGW. Man. N. Am. B. 1887, 337. Type, *Tyrannus borealis* SWAINS.

459. **Contopus borealis** (SWAINS.).

Olive-sided Flycatcher.

*Tyrannus borealis* SWAINS. in SW. & RICH. Faun. B. A. II. 1831, 141, pl. 35.

*Contopus borealis* BAIRD, B. N. Am. 1858, 188.

[B 137, C 253, R 318, C 380.]

GEOG. DIST.—North America, breeding from the northern and the higher mountainous parts of the United States northward to British Columbia and the Saskatchewan River. Accidental on the Lower Yukon and in Greenland. In winter, south to Central America, Colombia, and northern Peru.

SUBGENUS **CONTOPUS** CABANIS.

460. **Contopus pertinax** CAB.

Coues's Flycatcher.

*Myiarchus pertinax* LICHT. Nomen. Mus. Berol. 1854, 16 (nomen nudum).

*Contopus pertinax* CAB. Mus. Hein. II. Sept. 30, 1859, 72.

[B —, C 254, R 319, C 381.]

GEOG. DIST.—Mountains of southern and central Arizona, south through Mexico to Guatemala.

461. **Contopus virens** (LINN.).

Wood Pewee.

*Muscicapa virens* LINN. S. N. ed. 12, I. 1766, 327.

*Contopus virens* CAB. J. f. O. 1855, 479.

[B 139, C 255, R 320, C 382.]

GEOG. DIST.—Eastern North America, west to the Plains, and from southern Canada southward, migrating through eastern Mexico and Honduras to Colombia and Ecuador. Breeds from Florida to Newfoundland.

462. **Contopus richardsonii** (SWAINS.).

Western Wood Pewee.

*Tyrannula richardsonii* SWAINS. in SW. & RICH. F. B. A. II. 1831, 146, pl. 46, lower fig.

*Contopus richardsonii* BAIRD, B. N. Am. 1858, 189.

[B 138, C 255a, R 321, C 383.]

GEOG. DIST.—Western United States, from the Plains to the Pacific, north to British Columbia and the interior of British America, south through Central America to Colombia.

- 462 *α*. **Contopus richardsonii peninsulæ** BREWST.  
Large-billed Wood Pewee.

*Contopus richardsonii peninsulæ* BREWST. Auk, VIII. April, 1891,  
144.

[B —, C —, R —, C —.]

GEOG. DIST.—Sierra de la Laguna, Lower California.

GENUS **EMPIDONAX** CABANIS.

*Empidonax* CAB. J. f. O. 1855, 480. Type, *Tyrannula pusilla*  
SWAINS.

463. **Empidonax flaviventris** BAIRD.  
Yellow-bellied Flycatcher.

*Tyrannula flaviventris* BAIRD (W. M. & S. F.), Pr. Ac. Nat. Sci.  
Phila. July, 1843, 283.

*Empidonax flaviventris* BAIRD, B. N. Am. 1858, 198.

[B 144, C 259, R 322, C 388.]

GEOG. DIST.—Eastern North America west to the Plains, and from  
southern Labrador south through eastern Mexico to Panama, breed-  
ing from the Northern States northward. Casual in Greenland.

464. **Empidonax difficilis** BAIRD.  
Western Flycatcher.

*Empidonax difficilis* BAIRD, B. N. Am. 1858, 198 (in text).

[B 144a, C 259, *part*, R 323, C 389.]

GEOG. DIST.—Western United States, from the Plains to the  
Pacific; south to Cape St. Lucas and through western Mexico to  
Costa Rica; north to southern Alaska.

- 464.1. **Empidonax cineritius** BREWST.  
St. Lucas Flycatcher.

*Empidonax cineritius* BREWST. Auk, V. Jan. 1888, 90.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California.

465. **Empidonax virescens** (VIEILL.).

Green-crested Flycatcher.

*Platyrhynchus virescens* VIEILL. Nouv. Dict. d'Hist. Nat. XXVII.  
1818, 22.*Empidonax virescens* BREWST. Auk, XII. April, 1895, 157.

[B 143, C 256, R 324, C 384.]

GEOG. DIST.—Eastern United States, north to southern New York and southern Michigan, west to the Plains, south to Cuba and Costa Rica. Rare or casual in southern New England (Hyde Park, Mass.).

466. **Empidonax traillii** (AUD.).

Traill's Flycatcher.

*Muscicapa traillii* AUD. Orn. Biog. I. 1832, 236.*Empidonax traillii* BAIRD, Bds. N. Am. 1858, 193.

[B 141, C 257a, R 325, C 386.]

GEOG. DIST.—Western North America, from the Mississippi Valley (Ohio, Illinois, and Michigan) to the Pacific, and from the Fur Countries south into Mexico.

466 a. **Empidonax traillii alnorum** BREWST.

Alder Flycatcher.

*Empidonax traillii alnorum* BREWST. Auk, XII. April, 1895, 161.

[B 140, C 257, R 325a, C 385.]

GEOG. DIST.—Eastern North America, from the Maritime Provinces and New England westward at least to northern Michigan, etc., breeding from the southern edge of the Canadian Fauna northward; in winter south to Central America.

467. **Empidonax minimus** BAIRD.

Least Flycatcher.

*Tyrannula minima* BAIRD (W. M. & S. F.), Pr. Ac. Nat. Sci.  
Phila. July, 1843, 284.*Empidonax minimus* BAIRD, B. N. Am. 1858, 195.

[B 142, C 258, R 326, C 387.]

GEOG. DIST.—Chiefly eastern North America, west to eastern Colorado and central Montana, south in winter to Central America. Breeds from the Northern States northward.

468. **Empidonax hammondi** (XANTUS).

Hammond's Flycatcher.

*Tyrannula hammondi* XANTUS, Pr. Ac. Nat. Sci. Phila. May, 1858, 117.*Empidonax hammondi* BAIRD, B. N. Am. 1858, 199.

[B 145, C 260, R 327, C 390.]

GEOG. DIST.—Western North America, from the western border of the Plains westward, north to Lesser Slave Lake and Alaska, and south to southern Mexico.

469. **Empidonax wrightii** BAIRD.

Wright's Flycatcher.

*Empidonax wrightii* BAIRD, B. N. Am. 1858, 200 (in text).

[B 146, C 261, R 328, C 391.]

GEOG. DIST.—Western United States, north to Oregon and Montana, and south to southern Mexico.

469.1. **Empidonax griseus** BREWST.

Gray Flycatcher.

*Empidonax griseus* BREWST. Auk, VI. April, 1889, 87.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California and portions of Sonora. (Arizona?)

[470.] **Empidonax fulvifrons** (GIRAUD).

Fulvous Flycatcher.

*Muscicapa fulvifrons* GIRAUD, Sixteen Sp. Tex. B. 1841, pl. ii.*Empidonax fulvifrons* SCL. P. Z. S. 1858, 301.

[B —, C —, R 329, C —.]

GEOG. DIST.—“Texas” (GIRAUD), and eastern Mexico.

470 a. **Empidonax fulvifrons pygmæus** (COUES).

Buff-breasted Flycatcher.

*Empidonax pygmæus* COUES, Ibis, 1865, 537.*Empidonax fulvifrons pygmæus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 356.

[B —, C 262, R 329a, C 392.]

GEOG. DIST.—Western New Mexico and southern Arizona, south into western Mexico.

GENUS **PYROCEPHALUS** GOULD.

*Pyrocephalus* GOULD, Zool. Voy. Beag. 1841, 44. Types, "*Pyrocephalus parvirostris* (GOULD), and *Muscicapa coronata* (AUCT.)."

471. **Pyrocephalus rubineus mexicanus** (SCL.).

Vermilion Flycatcher.

*Pyrocephalus mexicanus* SCL. P. Z. S. 1859, 45.

*Pyrocephalus rubineus* var. *mexicanus* COUES, Key, 1872, 177.

[B 147, C 263, R 330, C 394.]

GEOG. DIST.—Southern and central Arizona, southwestern Utah, and the valley of the Lower Rio Grande in Texas, south to Lower California and Guatemala.

GENUS **ORNITHION** HARTLAUB.

*Ornithion* HARTLAUB, J. f. O. 1853, 35. Type, *O. inermis* HARTL.

472. **Ornithion imberbe** (SCL.).

Beardless Flycatcher.

*Camptostoma imberbe* SCL. P. Z. S. 1857, 203.

*Ornithion imberbe* LAWR. Ibis, 1876, 497.

[B —, C —, R 331, C 393.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, south into eastern Mexico, Guatemala, and Nicaragua.

472 a. **Ornithion imberbe ridgwayi** BREWST.

Ridgway's Flycatcher.

*Ornithium imberbe ridgwayi* BREWST. Bull. Nutt. Orn. Cl. VII.  
Oct. 1882, 208.

[B —, C —, R —, C —.]

GEOG. DIST.—Southern Arizona, south in Mexico to Puebla and Jalisco.



## SUBORDER OSCINES. SONG BIRDS.

FAMILY **ALAUDIDÆ**. LARKS.GENUS **ALAUDA** LINNÆUS.

*Alauda* LINN. S. N. ed. 10, I. 1758, 165. Type, by elimination,  
*A. arvensis* LINN.

[473.] **Alauda arvensis** LINN.  
 Skylark.

*Alauda arvensis* LINN. S. N. ed. 10, I. 1758, 165.

[B —, C 55 *bis*, R 299, C 88.]

GEOG. DIST.—Europe and Asia. Accidental in Greenland and Bermuda.

GENUS **OTOCORIS** BONAPARTE.

*Otocoris* BONAP. Nouvi Ann. Sci. Nat. Bologna, II. 1838, 407.  
 Type, *Alauda alpestris* LINN.

474. **Otocoris alpestris** (LINN.).  
 Horned Lark.

*Alauda alpestris* LINN. S. N. ed. 10, 1758, 166.

*Otocoris alpestris* BONAP. Nouvi Ann. Sci. Nat. Bologna, II.  
 1838. 407.

[B 302, C 53, R 300, C 82.]

GEOG. DIST.—Northeastern North America, Greenland, and northern parts of the Old World; in winter south in eastern United States to the Carolinas, Illinois, etc.

474 a. **Otocoris alpestris leucolæma** (COUES).**Pallid Horned Lark.***Eremophila alpestris* b. *leucolæma* COUES, B. N. W. 1875, 38 (part).*Otocoris alpestris leucolæma* STEJN. Pr. U. S. Nat. Mus. V.

June 5, 1882, 34.

[B —, C 53b, R 300a, C 83.]

GEOG. DIST.—Interior of British America, and Alaska, south in winter into western United States.

474 b. **Otocoris alpestris praticola** HENSH.**Prairie Horned Lark.***O[tocorys] alpestris praticola* HENSH. Auk, I. July, 1884, 264.

[B —, C —, R —, C —.]

GEOG. DIST.—Upper Mississippi Valley and the region of the Great Lakes to New England, breeding eastward to northeastern New York and western Massachusetts, New Hampshire and Vermont, and migrating south to South Carolina, Texas, etc.

474 c. **Otocoris alpestris arenicola** HENSH.**Desert Horned Lark.***O[tocorys] alpestris arenicola* HENSH. Auk, I. July, 1884, 265.

[B —, C —, R —, C —.]

GEOG. DIST.—Great Plains and Great Basin of the United States, south in winter to northern Mexico.

474 d. **Otocoris alpestris giraudi** HENSH.**Texan Horned Lark.***Otocorys alpestris giraudi* HENSH. Auk, I. July, 1884, 266.

[B —, C —, R —, C —.]

GEOG. DIST.—Eastern and southeastern Texas.

474 e. **Otocoris alpestris chrysolæma** (WAGL.).**Mexican Horned Lark.***Alauda chrysolæma* WAGL. Isis, 1831, 530.*Otocoris alpestris chrysolæma* STEJN. Pr. U. S. Nat. Mus. V.

June 5, 1882, 34.

[B—, C 53a, R 300b, C 84.]

GEOG. DIST.—Coast district of California (north to Nicasio), northern Lower California, and parts of Mexico (Mirador, Vera Cruz, Valley of Mexico, etc.).

474f. **Otocoris alpestris rubea** HENSH.

Ruddy Horned Lark.

*O[tocorys] alpestris rubeus* HENSH. Auk, I. July, 1884, 267.

[B—, C—, R—, C—.]

GEOG. DIST.—Sacramento and San Joaquin Valleys, California.

474g. **Otocoris alpestris strigata** HENSH.

Streaked Horned Lark.

*O[tocorys] alpestris strigata* HENSH. Auk, I. July, 1884, 267.

[B—, C—, R—, C—.]

GEOG. DIST.—Coast region of Washington, Oregon, and British Columbia. Islands off coast of southern California.

474h. **Otocoris alpestris adusta** DWIGHT.

Scorched Horned Lark.

*Otocoris alpestris adusta* DWIGHT, Auk, VII. April, 1890, 148.

[B—, C—, R—, C—.]

GEOG. DIST.—Southern Arizona and New Mexico, western Texas, and southward into Mexico.

474i. **Otocoris alpestris merrilli** DWIGHT.

Dusky Horned Lark.

*Otocoris alpestris merrilli* DWIGHT, Auk, VII. April, 1890, 153.

[B—, C—, R—, C—.]

GEOG. DIST.—Eastern Oregon, Washington, and British Columbia, between the Cascade and Rocky Mountains; southward in winter into Nevada and California.

474j. **Otocoris alpestris pallida** TOWNSEND.

Sonoran Horned Lark.

*Otocoris alpestris pallida* TOWNSEND, Proc. U. S. Nat. Mus., XIII, 1890, 138.

[B—, C—, R—, C—.]

GEOG. DIST.—Lower California and Sonora.

FAMILY **CORVIDÆ**. CROWS, JAYS, MAGPIES, ETC.SUBFAMILY **GARRULINÆ**. MAGPIES AND JAYS.GENUS **PICA** BRISSON.

*Pica* BRISS. Orn. II. 1760, 35. Type, *Corvus pica* LINN.

475. **Pica pica hudsonica** (SAB.).

American Magpie.

*Corvus hudsonicus* SAB. App. Frankl. Journ. 1823, 25, 671.

*Pica pica hudsonica* JORDAN, Man. Vert. ed. 4, 1884, 94.

[B 432, C 233, R 286, C 347.]

GEOG. DIST.—Northern and western North America, from the Plains to the Cascade Mountains and north to Alaska; casually east and south to Michigan (accidental in northern Illinois in winter), and in the Rocky Mountains to New Mexico and Arizona. Replaced in California, west of the Sierra Nevada, by the next species.

476. **Pica nuttalli** AUD.

Yellow-billed Magpie.

*Pica nuttalli* AUD. Orn. Biog. IV. 1838, 450, pl. 362.

[B 433, C 233a, R 287, C 348.]

GEOG. DIST.—California, west of the Sierra Nevada, from Sacramento Valley south to about latitude 34°, locally distributed.

GENUS **CYANOCITTA** STRICKLAND.

*Cyanocitta* STRICKL. Ann. Nat. Hist. XV. 1845, 261. Type,

*Corvus cristatus* LINN.

477. **Cyanocitta cristata** (LINN.).

Blue Jay.

*Corvus cristatus* LINN. S. N. ed. 10, I. 1758, 106.

*Cyanocitta cristata* STRICKL. Ann. Nat. Hist. XV. 1845, 261.

[B 434, C 234, R 289, C 349.]

GEOG. DIST.—Eastern North America to the Plains, and from the Fur Countries south to Florida and eastern Texas.

477 *a.* **Cyanocitta cristata florincola** COUES.  
Florida Blue Jay.

*Cyanocitta cristata florincola* COUES, Key, ed. 2, 1884, 421.

[B —, C —, R —, C —.]

GEOG. DIST.—Florida and the Gulf coast to southeastern Texas, casually along the coast to southwestern Texas.

478. **Cyanocitta stelleri** (GMEL.).  
Steller's Jay.

*Corvus stelleri* GMEL. S. N. I. 1788, 370.

*Cyanocitta stelleri* STRICKL. Ann. Nat. Hist. XV. 1845, 261.

[B 435, C 235, R 290, C 350.]

GEOG. DIST.—Pacific coast of North America, from Sitka south through the Cascade and Coast Ranges to Monterey Bay.

478 *a.* **Cyanocitta stelleri frontalis** (RIDGW.).  
Blue-fronted Jay.

*Cyanura stelleri* var. *frontalis* RIDGW. Am. Journ. Sc. & Arts, 3d ser. V. Jan. 1873, 41.

*Cyanocitta stelleri* var. *frontalis* BOUCARD, Cat. Av. 1876, 279.

[B —, C 235*b*, R 290*a*, C 353.]

GEOG. DIST.—Southern coast ranges and Sierra Nevada of California and western Nevada, from Fort Crook south to northern Lower California.

478 *b.* **Cyanocitta stelleri macrolopha** (BAIRD).  
Long-crested Jay.

*Cyanocitta macrolopha* BAIRD, Pr. Ac. Nat. Sci. Phila. 1854, 118.

*Cyanocitta stelleri macrolopha* COUES, Bull. Nutt. Orn. Cl. V. April, 1880, 98.

[B 436, C 235*a*, R 290*b*, 290*c*, C 352.]

GEOG. DIST.—Southern Rocky Mountains, southern Arizona and northwestern Mexico.

478 *c*. **Cyanocitta stelleri annectens** (BAIRD).

**Black-headed Jay.**

[*Cyanura stelleri*] var. *annectens* BAIRD, Hist. N. Am. B. II. 1874, 281, in text.

*Cyanocitta stelleri annectens* RIDGW. Proc. U. S. Nat. Mus. III. 1880, 184.

[B 436, *part*, C 255*a*, *part*, R 290*b*, C 352, *part*.]

GEOG. DIST.—Northern Rocky Mountains, south to Wahsatch range, west to eastern Oregon and Washington. .

GENUS **APHELOCOMA** CABANIS.

*Aphelocoma* CABANIS, Mus. Hein. I. Oct. 15, 1851, 221. Type, *Garrulus californicus* VIG.

479. **Aphelocoma floridana** (BARTR.)

**Florida Jay.**

*Corvus floridanus* BARTR. Trav. Carol. 1791, 291.

*Aphelocoma floridana* CAB. Mus. Hein. I. 1851, 221.

[B 439, C 236, R 291, C 354.]

GEOG. DIST.—Florida, of local distribution.

480. **Aphelocoma woodhouseii** (BAIRD).

**Woodhouse's Jay.**

*Cyanocitta woodhouseii* BAIRD, B. N. Am. 1858, 585, pl. 59.

*Aphelocoma woodhouseii* RIDGW. Field and Forest, June, 1877, 208.

[B 438, C 236*a*, R 292, C 355.]

GEOG. DIST.—Western United States, from the desert ranges of southern California north to eastern Oregon, east to Montana, Wyoming, Colorado and New Mexico, and south to northern Mexico.



480.1. **Aphelocoma cyanotis** RIDGW.

Blue-eared Jay.

*Aphelocoma cyanotis* RIDGW. Man. N. Am. Bds. 1887, 357.

[B —, C —, R —, C —.]

GEOG. DIST.—Northern Mexico, ranging northward into western Texas.

481. **Aphelocoma californica** (VIG.).

California Jay.

*Garrulus californicus* VIG. Zool. Beech. Voy. 1839, 21, pl. v.*Aphelocoma californica* CAB. Mus. Hein. I. Oct. 15, 1851, 221.[B 437, C 236*b*, R 293, C 356.]

GEOG. DIST.—Pacific coast region, including both slopes of the Sierra Nevada, from the Columbia River to northern Lower California.

481 *a*. **Aphelocoma californica hypoleuca** RIDGW.

Xantus's Jay.

*Aphelocoma californica hypoleuca* RIDGW. Man. N. Am. B. 1887, 356.[B 437, *part*, C 236*b*, *part*, R 293, *part*, C 356, *part*.]

GEOG. DIST.—Lower California, from Cape St. Lucas north to Lat. 28°.

481 *b*. **Aphelocoma californica obscura** ANTHONY.

Belding's Jay.

*Aphelocoma californica obscura* ANTHONY, Proc. Cal. Ac. Sci. 2d ser. II. Oct. 11, 1889, 75.

[B —, C —, R —, C —.]

GEOG. DIST.—San Pedro Martir Mountains, Lower California.

481.1. **Aphelocoma insularis** HENSH.

Santa Cruz Jay.

*Aphelocoma insularis* HENSH. Auk, III. Oct. 1886, 452.

[B —, C —, R —, C —.]

GEOG. DIST.—Santa Cruz Island, southern California.

482. **Aphelocoma sieberii arizonæ** RIDGW.

Arizona Jay.

*Cyanocitta ultramarina* var. *arizonæ* RIDGW. Bull. Essex Inst. V.  
Dec. 1873, 199.*Aphelocoma sieberii arizonæ* RIDGW. Pr. U. S. Nat. Mus. VIII.  
1885, 355.

[B 440, C 237, R 295, C 357.]

GEOG. DIST.—Southern New Mexico and Arizona, and southward  
into Sonora and Chihuahua.GENUS **XANTHOURA** BONAPARTE.*Xanthoura* BONAP. Consp. Av. I. May 6, 1850, 380. Type,  
*Corvus yncas* BODD.483. **Xanthoura luxuosa** (LESS.).

Green Jay.

*Garrulus luxuosus* LESS. Rev. Zool. 1839, 100.*Xanthoura luxuosa* BONAP. Consp. Av. I. 1850, 380.

[B 442, C 238, R 296, C 358.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, and  
southward in eastern Mexico to Vera Cruz and Puebla.GENUS **PERISOREUS** BONAPARTE.*Perisoreus* BONAP. Saggio, 1831, 43. Type, *Corvus infaustus*  
LINN.484. **Perisoreus canadensis** (LINN.).

Canada Jay.

*Corvus canadensis* LINN. S. N. ed. 12, I. 1766, 158.*Perisoreus canadensis* BONAP. Geog. & Comp. List, 1838, 27.

[B 443, C 239, R 297, C 359.]

GEOG. DIST.—Northern New York, northern New England, and  
northern Michigan, northward to Arctic America.

484 *a.* **Perisoreus canadensis capitalis** RIDGW.

Rocky Mountain Jay.

*Perisoreus canadensis* var. *capitalis* "BAIRD MS." RIDGW. Bull. Essex Inst. V. Nov. 1873, 193.[B —, C 239*b*, R 297*a*, C 362.]

GEOG. DIST.—Rocky Mountain region of the United States, south to New Mexico and Arizona.

484 *b.* **Perisoreus canadensis fumifrons** RIDGW.

Alaskan Jay.

*Perisoreus canadensis fumifrons* RIDGW. Pr. U. S. Nat. Mus. III. March 27, 1880, 5.[B —, C —, R 297*b*, C 360.]

GEOG. DIST.—Alaska, except southern coast district.

484 *c.* **Perisoreus canadensis nigricapillus** RIDGW.

Labrador Jay.

*Perisoreus canadensis nigricapillus* RIDGW. Pr. U. S. Nat. Mus. V. June 5, 1882, 15.

[B —, C —, R —, C —.]

GEOG. DIST.—Coast district of Labrador, north to Ungava Bay.

485. **Perisoreus obscurus** (RIDGW.).

Oregon Jay.

*Perisoreus canadensis* var. *obscurus* RIDGW. Bull. Essex Inst. Nov. 1873, 194.*Perisoreus obscurus* SHARPE, Brit. Mus. Cat. B. III. 1877, 105.[B —, C 239*a*, R 298, C 361.]

GEOG. DIST.—Northwest coast, from the northern Sierra Nevada and Humboldt Bay, in California, to British Columbia.

SUBFAMILY **CORVINÆ**. CROWS.GENUS **CORVUS** LINNÆUS.*Corvus* LINN. S. N. ed. 10, I. 1758, 105. Type, by elimination, *C. corax* LINN.

486. **Corvus corax sinuatus** (WAGL.).

American Raven.

*Corvus sinuatus* WAGLER, Isis, 1829, 748.*Corvus corax sinuatus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 423, 424, C 226, R 280, C 338.]

GEOG. DIST.—Western United States, from the Rocky Mountains westward, and south to Guatemala.

486 a. **Corvus corax principalis** RIDGW.

Northern Raven.

*Corvus corax principalis* RIDGW. Man. N. Am. B. 1887, 361.[B 423, *part*, C 226, *part*, R 280, *part*, C 338, *part*.]

GEOG. DIST.—Northern North America, south to British Columbia, northern Michigan, New Brunswick, Maine, New Jersey, North Carolina, etc.

487. **Corvus cryptoleucus** COUCH.

White-necked Raven.

*Corvus cryptoleucus* COUCH, Pr. Ac. Nat. Sci. Phila. April, 1854, 66.

[B 425, C 227, R 281, C 339.]

GEOG. DIST.—Southern border of the United States, from Texas to southern California, north to Colorado and western Kansas, south into northern Mexico.

488. **Corvus americanus** AUD.

American Crow.

*Corvus americanus* AUD. Orn. Biog. II. 1834, 317.

[B 426, C 228, R 282, C 340.]

GEOG. DIST.—North America, from the Fur Countries to the southern border of the United States. Locally distributed in the West.

488 a. **Corvus americanus floridanus** BAIRD.

Florida Crow.

*Corvus americanus* var. *floridanus* BAIRD, B. N. Am. 1858, 568.

[B 427, C 228a, R 282a, C 341.]

GEOG. DIST.—Florida.

489. **Corvus caurinus** BAIRD.

Northwest Crow.

*Corvus caurinus* BAIRD, B. N. Am. 1858, 569.[B 428, C 228*b*, R 282*b*, C 342.]

GEOG. DIST.—Northwest coast, from California to Sitka.

490. **Corvus ossifragus** WILS.

Fish Crow.

*Corvus ossifragus* WILS. Am. Orn. V. 1812, 27, pl. 37, fig. 2.

[B 429, C 229, R 283, C 343.]

GEOG. DIST.—Atlantic and Gulf coasts, from southern Connecticut to Louisiana. Common in the lower Hudson Valley; casual in Massachusetts.

GENUS **NUCIFRAGA** BRISSON.*Nucifraga* BRISSON, Orn. II. 1760, 58. Type, *N. caryocatactes*.SUBGENUS **PICICORVUS** BONAPARTE.*Picicorvus* BONAP. Consp. Av. I. 1850, 384. Type, *Corvus columbianus* WILS.491. **Nucifraga columbiana** (WILS.).

Clarke's Nutcracker.

*Corvus columbianus* WILS. Am. Orn. III. 1811, 29, pl. 20, fig. 3.*Nucifraga columbiana* AUD. Orn. Biog. IV. 1834, 459.

[B 430, C 230, R 284, C 344.]

GEOG. DIST.—Higher coniferous forests of western North America, from the eastern slope of the Rocky Mountains to the Pacific, and from Putnam River, Alaska, south to Arizona and northern Lower California. Accidental in Kansas, Missouri and Arkansas.

GENUS **CYANOCEPHALUS** BONAPARTE.*Cyanocephalus* BONAP. Oss. Stat. Zool. Eur. Vertebr. 1842, 17.Type, *Gymnorhinus cyanocephalus* WIED.

492. **Cyanocephalus cyanocephalus** (WIED).

Piñon Jay.

*Gymnorhinus cyanocephalus* WIED, Reise N. Amer. II. 1841, 21.*Cyanocephalus cyanocephalus* STEJN. Auk, I. 1884, 230.

[B 431, C 231, R 285, C 345.]

GEOG. DIST.—Rocky Mountains, west to the Cascade Range and the Sierra Nevada, and from British America south to northern Lower California. Accidental in eastern Kansas and eastern Nebraska.

FAMILY **STURNIDÆ**. STARLINGS.GENUS **STURNUS** LINNÆUS.

*Sturnus* LINN. S. N. ed. 10, I. 1758, 167. Type, by elimination,  
*S. vulgaris* LINN.

[493.] **Sturnus vulgaris** LINN.

Starling.

*Sturnus vulgaris* LINN. S. N. ed. 10, I. 1758, 167.

[B—, C—, R 279, C 363.]

GEOG. DIST.—Europe and northern Asia; accidental in Greenland. Introduced and apparently well-established in the vicinity of New York city.

FAMILY **ICTERIDÆ**. BLACKBIRDS, ORIOLES, ETC.GENUS **DOLICHONYX** SWAINSON.

*Dolichonyx* SWAINS. Phil. Mag. I. June, 1827, 435. Type,  
*Fringilla oryzivora* LINN.

494. **Dolichonyx oryzivorus** (LINN.).

Bobolink.

*Fringilla oryzivora* LINN. S. N. ed. 10, I. 1758, 179.*Dolichonyx oryzivorus* SWAINS. Zool. Jour. III. 1827, 351.



[B 399, C 210, R 257, C 312.]

GEOG. DIST.—Eastern North America, west to Montana, eastern Nevada, Utah, and Idaho; north to Ontario, and the southern parts of Manitoba, Assiniboia, and Alberta; south, in winter, to the West Indies and South America. Breeds from the Middle States northward, and winters south of the United States.

GENUS **MOLOTHRUS** SWAINSON.

*Molothrus* SWAINS. F. B. A. II. 1831, 277. Type, *Fringilla pecoris* GMEL. = *Oriolus ater* BODD.

495. **Molothrus ater** (BODD.).

Cowbird.

*Oriolus ater* BODD. Tabl. Pl. Enlum. 1783, 37.

*Molothrus ater* GRAY, Handl. B. II. 1870, 36.

[B 400, C 211, R 258, C 313.]

GEOG. DIST.—United States, from the Atlantic to the Pacific, north into southern British America, south, in winter, into Mexico.

495 a. **Molothrus ater obscurus** (GMEL.).

Dwarf Cowbird.

*Sturnus obscurus* GMEL. S. N. I. ii. 1788, 804.

*M[olothrus]* *ater* var. *obscurus* COUES, B. N. W. 1874, 180, in text.

[B —, C 211a, R 258a, C 314.]

GEOG. DIST.—Southern United States, from Texas to southern Arizona and Lower California, south into Mexico.

GENUS **CALLOTHRUS** CASSIN.

*Callothrus* CASS. Proc. Ac. Nat. Sci. Phil. 1866, 18. Type, *Psarocolius æneus* WAGL.

496. **Callothrus robustus** (CAB.).

Red-eyed Cowbird.

*Molothrus robustus* CAB. Mus. Hein. I. Sept. 1851, 193, footnote.*Callothrus robustus* RIDGW. Man. N. Am. B. 1887, 589.

[B—, C—, R 259, C 315.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, and southward to Panama.

GENUS **XANTHOCEPHALUS** BONAPARTE.*Xanthocephalus* BONAP. Consp. Av. I. 1850, 431. Type, *Icterus icterocephalus* BONAP. = *I. xanthocephalus* BONAP.497. **Xanthocephalus xanthocephalus** (BONAP.).

Yellow-headed Blackbird.

*Icterus xanthocephalus* BONAP. Journ. Ac. Nat. Sci. Phila. V. 1826, 223.*Xanthocephalus xanthocephalus* JORDAN, Man. Vert. ed. 4, 1884, 92.

[B 404, C 213, R 260, C 319.]

GEOG. DIST.—Western North America, from Wisconsin, Illinois and Texas to the Pacific coast, and from British Columbia and the Saskatchewan River southward to the Valley of Mexico. Accidental in Ontario and the Atlantic States (Massachusetts, District of Columbia, South Carolina, Florida).

GENUS **AGELAIUS** VIEILLOT.*Agelaius* VIEILL. Analyse, 1816, 33. Type, *Oriolus phoeniceus* LINN.498. **Agelaius phoeniceus** (LINN.).

Red-winged Blackbird.

*Oriolus phoeniceus* LINN. S. N. ed. 12, I. 1766, 161.*Agelaius phoeniceus* VIEILL. Nouv. Dict. d'Hist. Nat. XXXIV. 1819, 539.

[B 401, C 212, R 261, C 316.]

GEOG. DIST.—North America in general, from Great Slave Lake south to Costa Rica, excepting western Mexico and the lower Colorado Valley, southern Florida, the Gulf coast of Louisiana, and the lower Rio Grande Valley in Texas during the breeding season.

498 a. **Agelaius phoeniceus sonoriensis** RIDGW.  
Sonoran Red-wing.

*Agelaius phoeniceus sonoriensis* RIDGW. Man. N. Am. B. 1887, 370.

[B 401, *part*, C 212, *part*, R 261, *part*, C 316, *part*.]

GEOG. DIST.—Northern Mexico and contiguous borders of the United States, from the Lower Rio Grande Valley and southern Arizona north to the lower Colorado Valley, California, and Chilliwack, British Columbia.

498 b. **Agelaius phoeniceus bryanti** RIDGW.  
Bahaman Red-wing.

*Agelaius phoeniceus bryanti* RIDGW. Man. N. Am. B. 1887, 370.

[B 401, *part*, C 212, *part*, R 261, *part*, C 316, *part*.]

GEOG. DIST.—Bahamas and southern Florida, west to the Gulf coast of Louisiana (Lake Borgne), south to Yucatan and Nicaragua.

499. **Agelaius gubernator** (WAGL.).  
Bicolored Blackbird.

*Psarocolius gubernator* WAGL. Isis, IV. 1832, 281.

*Agelaius gubernator* BONAP. P.Z.S. 1837 (June, 1838), 110.

[B 402, C 212a, R 261a, C 317.]

GEOG. DIST.—Pacific coast districts, from western Washington, south to Lower California, west of the Cascades and the Sierra Nevada. Casually to western Nevada and southeastern California (Inyo Co.).

500. **Agelaius tricolor** (AUD.).  
Tricolored Blackbird.

*Icterus tricolor* "NUTT." AUD. Orn. Biog. V. 1839, pl. 388, fig. 1.

*Agelaius tricolor* BONAP. Geog. & Comp. List, 1838, 29.

[B 403, C 212b, R 262, C 318.]

GEOG. DIST.—Southwestern Oregon, south through California, west of the Sierra Nevada, to northern Lower California.

GENUS **STURNELLA** VIEILLOT.

*Sturnella* VIEILL. Analyse, 1816, 34. Type, *Alauda magna* LINN.

501. **Sturnella magna** LINN.).

Meadowlark.

*Alauda magna* LINN. S. N. ed. 10, I. 1758, 167.

*Sturnella magna* SWAINS. Phil. Mag. I. 1827, 436.

[B 406, C 214, R 263, C 320.]

GEOG. DIST.—Eastern United States and southern Canada to the Plains. Breeds from the Gulf of Mexico northward.

501 a. **Sturnella magna mexicana** (SCL.).

Mexican Meadowlark.

*Sturnella mexicana* SCL. Ibis, 1861, 179.

*Sturnella magna* var. *mexicana* B. B. & R. Hist. N. Am. B. II. 1874, 172.

[B —, C —, R 263a, C 321.]

GEOG. DIST.—Valley of the Lower Rio Grande and southern Arizona, and south through eastern and central Mexico to Panama.

501 b. **Sturnella magna neglecta** (AUD.).

Western Meadowlark.

*Sturnella neglecta* AUD. B. Am. VII. 1843, 339, pl. 487.

*Sturnella magna* var. *neglecta* ALLEN, Bull. M. C. Z. III. No. 2, July, 1872, 178.

[B 407, C 214a, R 264, C 322.]

GEOG. DIST.—Western United States, from Wisconsin, Illinois, Iowa, Texas, etc., west to the Pacific coast and north to British Columbia and Manitoba, south through central and western Mexico to Guanajuato and Jalisco.

GENUS **ICTERUS** BRISSON.SUBGENUS **ICTERUS**.

*Icterus* BRISS. Orn. II. 1760, 85. Type, by elimination, *Oriolus icterus* LINN.

[502.] **Icterus icterus** (LINN.).

Troupial.

*Oriolus icterus* LINN. S. N. ed. 12, I. 1766, 161.*Icterus icterus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 408, C —, R 265, C 323.]

GEOG. DIST.—West Indies (introduced) and northern South America. Accidental at Charleston, S. C. (AUDUBON).

503. **Icterus audubonii** GIRAUD.

Audubon's Oriole.

*Icterus audubonii* GIRAUD, Sixteen Sp. Texas B. 1841, 3.

[B 409, C 220, R 266, C 330.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, and southward in Mexico to Oaxaca.

504. **Icterus parisorum** BONAP.

Scott's Oriole.

*Icterus parisorum* BONAP. P. Z. S. 1837 (June, 1838), 110.

[B 411, C 219, R 268, C 329.]

GEOG. DIST.—Southwestern United States, from western Texas to southern California; north to northern New Mexico, southwestern Utah, southern Nevada, and California to Lat. 38° east of the Sierra Nevada; south to Lower California and on the tablelands of Mexico to Puebla and Vera Cruz.

SUBGENUS **PENDULINUS** VIEILLOT.

*Pendulinus* VIEILL. Analyse, 1816, 33. Type, *Oriolus spurius* LINN.

505. **Icterus cucullatus** SWAINS.

Hooded Oriole.

*Icterus cucullatus* SWAINS. Phil. Mag. I. 1827, 436.

[B 413, C 218, R 269, C 328.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, and southward through eastern and southern Mexico to British Honduras.

505 *a. Icterus cucullatus nelsoni* RIDGW.

Arizona Hooded Oriole.

*Icterus cucullatus nelsoni* RIDGW. Pr. U. S. Nat. Mus. Vol. VIII.

April 20, 1885, 19.

[B—, C—, R—, C—.]

GEOG. DIST.—Southwestern New Mexico and southern Arizona, west to Santa Barbara, California, and south to Mazatlan and Cape St. Lucas.

506. *Icterus spurius* (LINN.).

Orchard Oriole.

*Oriolus spurius* LINN. S. N. ed. 12, I. 1766, 162.*Icterus spurius* BONAP. Journ. Ac. Nat. Sci. Phila. III. 1823, 363.

[B 414, C 215, R 270, C 324.]

GEOG. DIST.—Eastern United States, north to the southern portions of New England, New York, Ontario, Michigan, and North Dakota, west to the Plains, south, in winter, to northern Colombia. Breeds throughout its United States range.

SUBGENUS **YPHANTES** VIEILLOT.

*Yphantes* VIEILL. Analyse, 1816, 33. Type, *Coracias galbula* LINN.

507. *Icterus galbula* (LINN.).

Baltimore Oriole.

*Coracias galbula* LINN. S. N. ed. 10, 1758, 108.*Icterus galbula* COUES, Bull. Nutt. Orn. Cl. V. Apr. 1880, 98.

[B 415, C 216, R 271, C 326.]

GEOG. DIST.—Eastern United States, north to Ontario and Manitoba, west nearly to the Rocky Mountains, south, in winter, through Mexico to Colombia.

508. *Icterus bullocki* (SWAINS.).

Bullock's Oriole.

*Xanthornus bullocki* SWAINS. Phil. Mag. I. 1827, 436.*Icterus bullocki* BONAP. Geog. & Comp. List, 1838, 29.



[B 416, C 217, R 272, C 327.]

GEOG. DIST.—Tablelands of Mexico, from Puebla and Valley of Mexico north through the western United States to Manitoba and British Columbia east of the Cascades, and from the western portion of the Plains to the Pacific.

GENUS **SCOLECOPHAGUS** SWAINSON.

*Scolecophagus* SWAINS. F. B. A. II. 1831, 286. Type, *Oriolus ferrugineus* GMEL. = *Turdus carolinus* MÜLL.

509. **Scolecophagus carolinus** (MÜLL.).

**Rusty Blackbird.**

*Turdus carolinus* MÜLLER, Syst. Nat. Suppl. 1776, 140.

*Scolecophagus carolinus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 356.

[B 417, C 221, R 273, C 331.]

GEOG. DIST.—Eastern North America, west to Alaska and the Plains. Breeds from northern New England, northern New York, and northern Michigan northward. Accidental in Lower California.

510. **Scolecophagus cyanocephalus** (WAGL.).

**Brewer's Blackbird.**

*Psarocolius cyanocephalus* WAGL. Isis, 1829, 758.

*Scolecophagus cyanocephalus* CAB. Mus. Hein. I. 1851, 193.

[B 418, C 222, R 274, C 332.]

GEOG. DIST.—Western North America, from the Plains to the Pacific, and from the Saskatchewan region south on the highlands of Mexico to Oaxaca.

GENUS **QUISCALUS** VIEILLLOT.SUBGENUS **QUISCALUS**.

*Quiscalus* VIEILL. Anal. 1816, 36. Type, *Gracula quiscula* LINN.

511. **Quiscalus quiscula** (LINN.).

Purple Grackle.

*Gracula quiscula* LINN. S. N. ed. 10, 1758, 109.*Quiscalus quiscula* JORDAN, Man. Vert. ed. 4, 1884, 93.

[B 421, C 225, R 278, C 335.]

GEOG. DIST.—Northern Alabama, eastern Tennessee, and east of the Alleghanies from Georgia to Massachusetts.

511 *a.* **Quiscalus quiscula aglæus** (BAIRD).

Florida Grackle.

*Quiscalus aglæus* BAIRD, Am. Jour. Sci. & Arts, XLI. Jan. 1866, 84.*Quiscalus quiscula aglæus* STEJN. Auk, II. Jan. 1885, 43, foot-note.[B 422, C —, R 278*a*, C 336.]

GEOG. DIST.—Florida, and the southern part of the Gulf States to Texas ; north along the Atlantic coast to Virginia.

511 *b.* **Quiscalus quiscula æneus** (RIDGW.).

Bronzed Grackle.

*Quiscalus æneus* RIDGW. Pr. Ac. Nat. Sci. Phila. June, 1869, 134.*Quiscalus quiscula æneus* STEJN. Auk, II. Jan. 1885, 43, foot-note.[B —, C 225*a*, R 278*b*, C 337.]

GEOG. DIST.—From the Alleghanies and southern New England north to Newfoundland and Great Slave Lake, west to the eastern base of the Rocky Mountains, and south to Louisiana and Texas. In migrations, the southeastern States, except Florida and the Atlantic coast district south of Virginia.

SUBGENUS **MEGAQUISCALUS** CASSIN.*Megaquiscalus* CASS. Pr. Ac. Nat. Sci. Phila. 1866, 409. Type,*Quiscalus major* VIEILL.512. **Quiscalus macrourus** SWAINS.

Great-tailed Grackle.

*Quiscalus macrourus* SWAINS. Anim. in Menag. 1838, 299.

[B 419, C 223, R 275, C 333.]

GEOG. DIST.—Eastern Texas, south to Central America.

513. **Quiscalus major** VIEILL.

Boat-tailed Grackle.

*Quiscalus major* VIEILL. Nouv. Dict. d'Hist. Nat. XXVIII. 1819, 487.

[B 420, C 224, R 277, C 334.]

GEOG. DIST.—Coast region of the South Atlantic and Gulf States, from Virginia to Texas.

FAMILY **FRINGILLIDÆ**. FINCHES, SPARROWS, ETC.GENUS **COCCOTHAUSTES** BRISSON.*Coccothraustes* BRISS. Orn. III. 1760, 218. Type, *Loxia coccothraustes* LINN.SUBGENUS **HESPERIPHONA** BONAPARTE.*Hesperiphona* BONAP. Compt. Rend. XXXI. Sept. 1850, 424. Type, *Fringilla vespertina* COOPER.514. **Coccothraustes vespertinus** (COOP.).

Evening Grosbeak.

*Fringilla vespertina* COOP. Ann. Lyc. N. Y. I. ii. 1825, 220.*Coccothraustes vespertina* SW. & RICH. Fauna Bor. Am. II. 1831, 269, pl. 68.

[B 303, C 136, R 165, C 189.]

GEOG. DIST.—Western British Provinces, east to Lake Superior, and casually to Michigan, Ohio, Ontario, New York, and New England.

514a. **Coccothraustes vespertinus montanus** (RIDGW.).

Western Evening Grosbeak.

*Hesperiphona vespertina* var. *montana* RIDGW. in Hist. N. Am. Bds., Land Bds., I. 1874, 449.*Coccothraustes vespertina montana* MEARNs, Auk, VII. July, 1890, 246.

[B 303, *part*, C 136, *part*, R 165, *part*, C 189, *part*.]

GEOG. DIST.—Western North America, from the Pacific coast eastward to the Rocky Mountains; southward over the tablelands of Mexico to Orizaba.

GENUS **PINICOLA** VIEILLLOT.

*Pinicola* VIEILL. Ois. Am. Sept. I. 1807, p. iv. Type, *P. rubra*  
VIEILL. = *Loxia enucleator* LINN.

515. **Pinicola enucleator** (LINN.).

Pine Grosbeak.

*Loxia enucleator* LINN. S. N. ed. 10, I. 1758, 171.

*Pinicola enucleator* CAB. in ERSCH & GRUBER, Encycl. 1st Sect.  
I. 1849, 219.

[B 304, C 137, R 166, C 190.]

GEOG. DIST.—Northern parts of the northern hemisphere, breeding in North America from northern New England, Quebec, the Rocky Mountains in Colorado, and about Lat. 37° in the Sierra Nevada, northward nearly to the limit of trees; south in winter irregularly into northeastern United States.

GENUS **PYRRHULA** BRISSON.

*Pyrrhula* BRISS. Orn. III. 1760, 308. Type, *Loxia pyrrhula*  
LINN.

[516.] **Pyrrhula cassini** (BAIRD).

Cassin's Bullfinch.

*Pyrrhula coccinea* var. *cassini* BAIRD, Trans. Chicago Ac. Sci. I.  
1869, 316.

*Pyrrhula cassini* TRISTRAM, Ibis, Apr. 1871, 231.

[B—, C 138, R 167, C 191.]

GEOG. DIST.—Eastern Siberia. Accidental at Nulato, Alaska.

GENUS **CARPODACUS** KAUP.

*Carpodacus* KAUP, Skizz. Entw.-Gesch. Eur. Thierw. 1829, 161.  
Type, *Loxia erythrina* PALL.

517. **Carpodacus purpureus** (GMEL.).

Purple Finch.

*Fringilla purpurea* GMEL. S. N. I. ii. 1788, 923.*Carpodacus purpureus* GRAY, Gen. B. II. 1844, 384.

[B 305, C 139, R 168, C 194.]

GEOG. DIST.— Eastern North America, from the Atlantic coast to the Plains. Breeds from the Middle States northward.

517 a. **Carpodacus purpureus californicus** BAIRD.

California Purple Finch.

*Carpodacus californicus* BAIRD, B. N. Am. 1858, 413.

*Carpodacus purpureus* var. *californicus* B. B. & R. Hist. N. Am.  
B. I. 1874, 465.

[B 306, C —, R 168a, C —.]

GEOG. DIST.— Pacific coast region, from British Columbia south in winter to southern California.

518. **Carpodacus cassini** BAIRD.

Cassin's Purple Finch.

*Carpodacus cassini* BAIRD, Pr. Ac. Nat. Sci. Phila. June, 1854,  
119.

[B 307, C 140, R 169, C 195.]

GEOG. DIST.— Western United States, from the eastern base of the Rocky Mountains to the Pacific coast, and south over the plateau region of Mexico to Mt. Orizaba.

SUBGENUS **BURRICA** RIDGW.

*Burrica* RIDGW. Man. N. Am. B. 1887, 390. Type, *Fringilla mexicana* MÜLL.

519. **Carpodacus mexicanus frontalis** (SAY).

House Finch.

*Fringilla frontalis* SAY, LONG's Exp. II. 1823, 40.

*Carpodacus mexicanus frontalis* RIDGW. Man. N. Am. B. 1887,  
391.

[B 308, C 141, R 170, C 196.]

GEOG. DIST.—Colorado and western Texas, westward to Oregon and California, south into Lower California and western Mexico.

519 *b*. **Carpodacus mexicanus ruberrimus** RIDGW.

St. Lucas House Finch.

*Carpodacus frontalis ruberrimus* RIDGW. Man. N. Am. B. 1887, 391, foot-note.

*Carpodacus mexicanus ruberrimus* RIDGW. Man. N. Am. B. 1887, 594.

[B —, C 141a, R 170a, C 197.]

GEOG. DIST.—Lower California, and probably adjacent parts of Sonora.

520. **Carpodacus amplus** RIDGW.

Guadalupe House Finch.

*Carpodacus amplus* RIDGW. Bull. U. S. Geol. & Geog. Surv. Terr. II. No. 2, April 1, 1876, 187.

[B —, C —, R 171, C —.]

GEOG. DIST.—Guadalupe Island, Lower California.

GENUS **LOXIA** LINNÆUS.

*Loxia* LINN. S. N. ed. 10, l. 1758, 171. Type, by elimination,  
*Loxia curvirostra* LINN.

521. **Loxia curvirostra minor** (BREHM).

American Crossbill.

*Crucirostra minor* BREHM, Naumannia, 1853, 193.

*Loxia curvirostra minor* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 318, C 143, R 172, C 199.]

GEOG. DIST.—Northern North America, resident sparingly south in the eastern United States to Maryland and Tennessee, and in the Alleghanies; irregularly abundant in winter. Casual at Charleston, S. C., and New Orleans, La.



521 *a.* ***Loxia curvirostra stricklandi*** RIDGW.

Mexican Crossbill.

*Loxia curvirostra stricklandi* RIDGW. Pr. U. S. Nat. Mus. VIII.  
1885, 354.[B 318*a*, C 143*a*, R 172*a*, C 200.]

GEOG. DIST.—Mountains of Wyoming and Colorado, west to the Sierra Nevada, and south through New Mexico, Arizona, and the tablelands of Mexico to Guatemala.

522. ***Loxia leucoptera*** GMEL.

White-winged Crossbill.

*Loxia leucoptera* GMEL. S. N. I. ii. 1788, 540.

[B 319, C 142, R 173, C 198.]

GEOG. DIST.—Northern parts of North America, south into the United States in winter. Breeds from northern New England northward.

GENUS **LEUCOSTICTE** SWAINSON.*Leucosticte* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831,  
265. Type, *Linaria tephrocotis* SWAINS.523. ***Leucosticte griseonucha*** (BRANDT).

Aleutian Leucosticte.

*Fringilla (Linaria) griseonucha* BRANDT, Bull. Ac. St. Pétersb.  
Nov. 1841, 36.*Leucosticte griseinucha* BONAP. Consp. Av. I. 1850, 537.[B 323, C 144*a*, R 174, C 205.]

GEOG. DIST.—Aleutian Islands, including Kadiak, Unalaska, Pribilof, and Commander Islands.

524. ***Leucosticte tephrocotis*** SWAINS.

Gray-crowned Leucosticte.

*Linaria (Leucosticte) tephrocotis* SWAINS. in SW. & RICH. Fauna  
Bor. Am. II. 1831, 265, pl. 50.*Leucosticte tephrocotis* SWAINS. in SW. & RICH. Fauna Bor. Am.  
II. 1831, 494.

[B 322, C 144, R 175, C 203.]

GEOG. DIST.—Interior of British America, south in winter throughout the entire Rocky Mountain region of the United States, but most abundant on the eastern slope. Known to breed only in the Sierra Nevada in California.

524 *a. Leucosticte tephrocotis littoralis* (BAIRD).

Hepburn's Leucosticte.

*Leucosticte littoralis* BAIRD, Trans. Chicago Ac. Sci. I. i. 1869, 318, pl. 28, fig. 1.

*Leucosticte tephrocotis* var. *littoralis* COUES, Key, 1872, 130.

[B —, C —, R 175*a*, C 204.]

GEOG. DIST.—In summer, probably the interior mountainous regions of British Columbia; in winter, northwest coast, from Kadiak southward, and eastward in the Rocky Mountain region to Colorado.

525. *Leucosticte atrata* RIDGW.

Black Leucosticte.

*Leucosticte atrata* RIDGW. American Sportsman, July 18, 1874, 241; Bull. U. S. Geol. & Geog. Surv. Terr. 2d ser. No. 2, May 11, 1875, 69.

[B —, C —, R 176, C 201.]

GEOG. DIST.—Breeds in the Salmon River Mountains, Idaho, and probably other northern ranges; in winter, mountains of Colorado and Utah.

526. *Leucosticte australis* RIDGW.

Brown-capped Leucosticte.

*Leucosticte tephrocotis* var. *australis* "ALLEN MS." RIDGW. Bull. Essex Inst. V. Dec. 1873, 197.

*Leucosticte australis* RIDGW. Bull. U. S. Geol. & Geog. Surv. Terr. 2d ser. No. 2, May 11, 1875, 79.

[B —, C —, R 177, C 203.]

GEOG. DIST.—Mountains of Colorado, breeding above timber-line, descending into the valleys in winter; New Mexico.

GENUS **ACANTHIS** BECHSTEIN.

*Acanthis* BECHST. Orn. Tasch. Deutschl. 1803, 125. Type,  
*Fringilla linaria* LINN.

527. **Acanthis hornemannii** (HOLB.).

Greenland Redpoll.

*Linota hornemannii* HOLBÖLL, Naturh. Tidskr. IV. 1843, 398.

*Acanthis hornemannii* STEJN. Auk, I. April, 1884, 152.

[B 321, C —, R 178, C 209.]

GEOG. DIST.—Greenland and eastern Arctic America.

527 a. **Acanthis hornemannii exilipes** (COUES).

Hoary Redpoll.

*Ægiothus exilipes* COUES, Pr. Ac. Nat. Sci. Phila. 1861, 385.

*Acanthis hornemannii exilipes* STEJN. Auk, I. April, 1884, 152.

[B —, C 146b, R 178a, C 210.]

GEOG. DIST.—Arctic America and northeastern Asia, south in winter (rarely?) to the northern border of the United States.

528. **Acanthis linaria** (LINN.).

Redpoll.

*Fringilla linaria* LINN. S. N. ed. 10, I. 1758, 182.

*Acanthis linaria* BONAP. & SCHLEG. Mon. Lox. 1850, 48.

[B 320, C 146, 146a, R 179, C 207.]

GEOG. DIST.—Northern portions of northern hemisphere, south irregularly in winter, in North America, to the middle United States (Virginia, Kansas, southeastern Oregon).

528 a. **Acanthis linaria holbœllii** (BREHM).

Holböll's Redpoll.

*Linaria holbœllii* BREHM. Handb. Vög. Deutschl. 1831, 280.

*Acanthis linaria* β. *holbœllii* DUBOIS, Consp. Av. Europ. 1871, 18.

[B —, C —, R 179a, *part*, C 208, *part*.]

GEOG. DIST.—Northern portions of northern hemisphere, near the seacoast, south in winter to northern New York and Massachusetts.

528 *b.* ***Acanthis linaria rostrata*** (COUES).

Greater Redpoll.

*Ægiothus rostratus* COUES, Pr. Ac. Nat. Sci. Phila. 1861, 378.*Acanthis linaria rostrata* STEJN. Auk, I. April, 1884, 153.[B —, C —, R 179*a*, *part*, C 208, *part*.]

GEOG. DIST.—Greenland and northeastern North America, south irregularly in winter to New England, New York, and northern Illinois.

GENUS **SPINUS** KOCH.

*Spinus* KOCH, Bayr. Zool. 1816, 233. Type, *Fringilla spinus* LINN.

529. ***Spinus tristis*** (LINN.).

American Goldfinch.

*Fringilla tristis* LINN. S. N. ed. 10, I. 1758, 181.*Spinus tristis* STEJN. Auk, I. Oct. 1884, 362.

[B 313, C 149, R 181, C 213.]

GEOG. DIST.—Temperate North America generally, from southern Labrador, Manitoba and British Columbia south, in winter, to the northern boundary of Lower California; breeding southward to the middle districts of the United States (Virginia, Kentucky, Kansas, and California), and wintering mainly within the United States.

529 *a.* ***Spinus tristis pallidus*** MEARNs.

Western Goldfinch.

*Spinus tristis pallidus* MEARNs, Auk, VII. July, 1890, 244.[B 313, *part*, C 149, *part*, R 181, *part*, C 213, *part*.]

GEOG. DIST.—Arizona.

530. ***Spinus psaltria*** (SAY).

Arkansas Goldfinch.

*Fringilla psaltria* SAY, LONG's Exp. II. 1823, 40.*Spinus psaltria* STEJN. Auk, I. Oct. 1884, 362.

[B 314, C 151, R 182, C 215.]

GEOG. DIST.—Western United States, from the Plains to the Pacific, and from southern Oregon, Colorado, and Utah southward to Cape St. Lucas and Sonora.

530 a. **Spinus psaltria arizonæ** (COUES).

Arizona Goldfinch.

*Chrysomitris mexicana* var. *arizonæ* COUES, Pr. Ac. Nat. Sci. Phila. 1866, 82.

*Spinus psaltria arizonæ* STEJN. Auk, I. Oct. 1884, 362.

[B —, C 151a, R 182a, C 216.]

GEOG. DIST.—Southern California, southwestern Utah, southern New Mexico and Arizona, southward into northern Mexico.

530 b. **Spinus psaltria mexicanus** (SWAINS.).

Mexican Goldfinch.

*Carduelis mexicanus* SWAINS. Phil. Mag. I. 1827, 435.

*Spinus psaltria mexicanus* STEJN. Auk, I. Oct. 1884, 362.

[B 315, C 159b, R 182b, C 217.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, southward through Mexico.

531. **Spinus lawrencei** (CASS.).

Lawrence's Goldfinch.

*Carduelis lawrencei* CASS. Pr. Ac. Nat. Sci. Phila. 1851, 105, pl. v.

*Spinus lawrencei* STEJN. Auk, I. Oct. 1884, 362.

[B 316, C 150, R 183, C 214.]

GEOG. DIST.—California west of the Sierra Nevada, from Lat. 40° south to northern Lower California; Arizona (Fort Mohave and Fort Whipple) in winter.

[532.] **Spinus notatus** (DUBUS).

Black-headed Goldfinch.

*Carduelis notata* DUBUS, Bull. Ac. Brux. XIV. pt. 2, 1847, 106.

*Spinus notatus* STEJN. Auk, I. Oct. 1884, 362.

[B 310, C —, R 184, C 218.]

GEOG. DIST.—Mountains of Guatemala and southern Mexico, north at least to central Vera Cruz; accidental in Kentucky (AUDUBON).

533. **Spinus pinus** (WILS.).

Pine Siskin.

*Fringilla pinus* WILS. Am. Orn. II. 1810, 133, pl. 17, fig. 1.

*Spinus pinus* STEJN. Auk, I. Oct. 1884, 362.

[B 317, C 148, R 185, C 212.]

GEOG. DIST.—North America generally, breeding in the British Provinces, Rocky Mountains, Sierra Nevada, and high mountains of Arizona, south to Lower California and the mountains of Mexico to Orizaba. Also breeds sparingly in northeastern United States.

GENUS **PLECTROPHENAX** STEJNEGER.

*Plectrophenax* STEJN. Pr. U. S. Nat. Mus. V. June 5, 1882, 33.

Type, *Emberiza nivalis* LINN.

534. **Plectrophenax nivalis** (LINN.).

Snowflake.

*Emberiza nivalis* LINN. S. N. ed. 10, I. 1758, 176.

*Plectrophenax nivalis* STEJN. Pr. U. S. Nat. Mus. V. 1882, 33.

[B 325, C 152, R 186, C 219.]

GEOG. DIST.—Northern parts of the northern hemisphere, breeding in the arctic regions; in North America south in winter into the northern United States, irregularly to Georgia, southern Illinois, Kansas, and Oregon.

534 a. **Plectrophenax nivalis townsendi** RIDGW.

Pribilof Snowflake.

*Plectrophenax nivalis townsendi* RIDGW. Man. N. Am. B. 1887,

403.



[B —, C —, R 186, *part*, C 219, *part*.]

GEOG. DIST.—Pribilof and Aleutian Islands, Alaska, and Commander Islands, Kamchatka.

535. **Plectrophenax hyperboreus** RIDGW.

McKay's Snowflake.

*Plectrophenax hyperboreus* RIDGW. Pr. U. S. Nat. Mus.<sup>2</sup> VII.  
June 11, 1884, 68.

[B —, C —, R —, C —.]

GEOG. DIST.—Western Alaska, breeding on Hall Island (and probably St. Matthew Island), Bering Sea.

GENUS **CALCARIUS** BECHSTEIN.

*Calcarius* BECHST. Taschb. Vög. Deutschl. 1803, 130. Type,  
*Fringilla lapponica* LINN.

536. **Calcarius lapponicus** (LINN.).

Lapland Longspur.

*Fringilla lapponica* LINN. S. N. ed. 10, I. 1758, 180.

*Calcarius lapponicus* STEJN. Pr. U. S. Nat. Mus. V. June 5, 1882,  
33.

[B 326, C 153, R 187, C 220.]

GEOG. DIST.—Northern portions of the northern hemisphere, breeding far north; in North America south in winter to the northern United States, irregularly to the Middle States, accidentally to South Carolina, and abundantly in the interior to Kansas and Colorado.

537. **Calcarius pictus** (SWAINS.).

Smith's Longspur.

*Emberiza (Plectrophanes) picta* SWAINS. in Sw. and RICH. Fauna  
Bor. Am. II. 1831, 250, pl. 49.

*Calcarius pictus* STEJN. Pr. U. S. Nat. Mus. V. June 5, 1882, 33.

[B 327, C 154, R 188, C 221.]

GEOG. DIST.—Interior of North America, from the Arctic coast to Illinois and Texas, breeding far north.

538. **Calcarius ornatus** (TOWNS.).

Chestnut-collared Longspur.

*Plectrophanes ornatus* TOWNS. Journ. Ac. Nat. Sci. Phila. VII.  
1837, 189.*Calcarius ornatus* STEJN. Pr. U. S. Nat. Mus. V. June 5, 1882,  
33.

[B 328, 329, C 155, R 189, C 222.]

GEOG. DIST.—Interior of North America, from the Saskatchewan Plains south through Texas to Orizaba. Breeds from eastern Manitoba, western Minnesota and eastern Nebraska westward and north-westward into Montana and Assiniboia. Rare west of the Rocky Mountains. Accidental in Massachusetts.

GENUS **RHYNCHOPHANES** BAIRD.*Rhynchophanes* BAIRD, B. N. Am. 1858, 432 (in text). Type,  
*Plectrophanes mccownii* LAWR.539. **Rhynchophanes mccownii** (LAWR.).

McCown's Longspur.

*Plectrophanes mccownii* LAWR. Ann. Lyc. N. Y. V. 1851, 122.*Rhynchophanes maccownii* RIDGW. Field & Forest, II. May, 1877,  
197.

[B 330, C 156, R 190, C 223.]

GEOG. DIST.—Interior of North America, from the Saskatchewan Plains south to Texas and northern Mexico; breeds from about the northern border of western Kansas northward throughout Nebraska, western North and South Dakota, Wyoming, and Montana to the Plains of the Saskatchewan.

GENUS **POOCÆTES** BAIRD.*Poocætes* BAIRD, B. N. Am. 1858, 447. Type, *Fringilla graminea*  
GMEL.540. **Poocætes gramineus** (GMEL.).

Vesper Sparrow.

*Fringilla graminea* GMEL. S. N. I. ii. 1788, 992.*Poocætes gramineus* BAIRD, B. N. Am. 1858, 447.

[B 337, *part*, C 161, R 197, C 232.]

GEOG. DIST.—Eastern North America to the Plains, from Nova Scotia and Ontario southward; breeds from Virginia, Kentucky, and Missouri northward.

540 *a*. **Poocætes gramineus confinis** BAIRD.

Western Vesper Sparrow.

*Poocætes gramineus* var. *confinis* BAIRD, B. N. Am. 1858, 448 (in text).

[B 337, *part*, C 161*a*, R 197*a*, C 232.]

GEOG. DIST.—Western United States, from the Plains to the Pacific, north into Manitoba and Assiniboia, south into Lower California and through Mexico as far as Jalapa, Vera Cruz.

540 *b*. **Poocætes gramineus affinis** MILLER.

Oregon Vesper Sparrow.

*Poocætes gramineus affinis* MILLER, Auk, V. Oct. 1888, 404.

[B 337, *part*, C 161*a*, *part*, R 197*a*, *part*, C 232, *part*.]

GEOG. DIST.—Northern California and western Oregon.

#### GENUS **AMMODRAMUS** SWAINSON.

*Ammodramus* SWAINS. Zool. Journ. III. 1827, 348. Type,  
*Fringilla caudacuta* WILSON.

#### SUBGENUS **PASSERCULUS** BONAPARTE.

*Passerculus* BONAP. Geog. & Comp. List, 1838, 33. Type,  
*Fringilla savanna* WILS.

541. **Ammodramus princeps** (MAYN.).

Ipswich Sparrow.

*Passerculus princeps* MAYN. Am. Nat. VI. 1872, 637.

*Ammodramus princeps* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885,

[B —, C 158, R 192, C 225.]

GEOG. DIST.—Atlantic coast, from Nova Scotia south, in winter, to Georgia. Breeds on Sable Island, Nova Scotia.

542. ***Ammodramus sandwichensis*** (GMEL.).

**Sandwich Sparrow.**

*Emberiza sandwichensis* GMEL. S. N. I. ii. 1788, 875.

*Ammodramus sandwichensis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 333, C 159*b*, R 193, C 226.]

GEOG. DIST.—Northwest coast, from the Columbia River to Unalaska.

542 *a*. ***Ammodramus sandwichensis savanna*** (WILS.).

**Savanna Sparrow.**

*Fringilla savanna* WILS. Am. Orn. III. 1811, 55, pl. 22, fig. 2.

*Ammodramus sandwichensis savanna* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 332, C 159, R 193*a*, C 227.]

GEOG. DIST.—Eastern North America, breeding from the northern United States to Labrador and Hudson Bay Territory.

542 *b*. ***Ammodramus sandwichensis alaudinus*** (BONAP.).

**Western Savanna Sparrow.**

*Passerculus alaudinus* BONAP. Compt. Rend. XXXVII. 1853, 918.

*Ammodramus sandwichensis alaudinus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 335, C —, R 193*b*, C 229.]

GEOG. DIST.—Western North America, from the Plains to the Pacific coast region, north to the Arctic coast.

542 *c*. ***Ammodramus sandwichensis bryanti*** RIDGW.

**Bryant's Marsh Sparrow.**

*Passerculus sandwichensis bryanti* RIDGW. Pr. U. S. Nat. Mus. VII. Jan. 19, 1885, 517.

*Ammodramus sandwichensis bryanti* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 334, *part*, C 159*a*, *part*, R 194, *part*, C 228, *part*.]

GEOG. DIST.— Salt marshes about San Francisco Bay, and south along the coast in winter to southern California.

543. ***Ammodramus beldingi*** RIDGW.

Belding's Marsh Sparrow.

*Passerculus beldingi* RIDGW. Pr. U. S. Nat. Mus. VII. Jan. 19, 1885, 516.

*Ammodramus beldingi* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 334, *part*, C 159*a*, *part*, R 194, *part*, C 228, *part*.]

GEOG. DIST.— Salt marshes of the Pacific coast, from Santa Barbara south to Todos Santos Island, Lower California.

544. ***Ammodramus rostratus*** CASS.

Large-billed Sparrow.

*Emberiza rostrata* CASS. Pr. Ac. Nat. Sci. Phila. 1852, 348.

*Ammodramus rostratus* CASS. Illustr. B. Cal. Tex. etc. 1855, 226, pl. 38.

[B 336, C 160, R 196, C 230.]

GEOG. DIST.— Coast of southern California, south in winter to Cape St. Lucas and northwestern Mexico.

544 *a*. ***Ammodramus rostratus guttatus*** (LAWR.).

St. Lucas Sparrow.

*Passerculus guttatus* LAWR. Ann. Lyc. N. Y. VIII. 1867, 473.

*Ammodramus rostratus guttatus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B —, C 160*a*, R 195, C 231.]

GEOG. DIST.— Southern Lower California.

SUBGENUS **CENTRONYX** BAIRD.

*Centronyx* BAIRD, B. N. Am. 1858, 440. Type, *Emberiza bairdii* AUD.

545. **Ammodramus bairdii** (AUD.).

Baird's Sparrow.

*Emberiza bairdii* AUD. B. Am. VII. 1843, 359, pl. 500.*Ammodramus bairdi* GIEBEL, Thes. Orn. I. 1872, 328.[B 331, C 157, 157*bis*, R 191, C 224.]

GEOG. DIST.—Interior of North America, from the plains of the Red River and Saskatchewan south to Texas, New Mexico, Arizona, and Chihuahua.

SUBGENUS **COTURNICULUS** BONAPARTE.

*Coturniculus* BONAP. Geog. & Comp. List, 1838, 32. Type, *Fringilla passerina* WILS.

546. **Ammodramus savannarum passerinus** (WILS.).

Grasshopper Sparrow.

*Fringilla passerina* WILS. Am. Orn. III. 1811, 76, pl. 26, fig. 5.*Ammodramus savannarum passerinus* RIDGW. Pr. U. S. Nat. Mus.

VIII. 1885, 355.

[B 338, *part*, C 162, R 198, C 234.]

GEOG. DIST.—Eastern United States and southern Canada, west to the Plains, south, in winter, to Florida, Cuba, Porto Rico, and coast of Central America.

546 a. **Ammodramus savannarum perpallidus** (COUES).

Western Grasshopper Sparrow.

*Coturniculus passerinus* var. *perpallidus* "RIDGW. MS." COUES, Key, 1872, 137.

*Ammodramus savannarum perpallidus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 338, *part*, C 162*a*, R 198*a*, C 235.]

GEOG. DIST.—Western United States, from the Plains to the Pacific coast, south to Cape St. Lucas and the tablelands of Mexico.

547. **Ammodramus henslowii** (AUD.).

Henslow's Sparrow.

*Emberiza henslowii* AUD. Orn. Biog. I. 1831, 360, pl. 77.*Ammodromus henslowi* GRAY, Gen. B. II. June, 1849, 374.



[B 339, *part*, C 163, *part*, R 199, *part*, C 236, *part*.]

GEOG. DIST.—Eastern United States, west to the Plains, north to southern New England and Ontario.

- 547 *a.* ***Ammodramus henslowii occidentalis*** BREWST.  
Western Henslow's Sparrow.

*Ammodramus henslowii occidentalis* BREWST. Auk, VIII. April, 1891, 145.

[B 339, *part*, C 163, *part*, R 199, *part*, C 236, *part*.]

GEOG. DIST.—South Dakota.

548. ***Ammodramus leconteii*** (AUD.).  
Leconte's Sparrow.

*Emberiza leconteii* AUD. B. Am. VII. 1843, 338, pl. 488.

*Ammodromus leconteii* GRAY, Gen. B. II. June, 1849, 374.

[B 340, C 164, R 200, C 237.]

GEOG. DIST.—From the Plains eastward to Illinois and Indiana, and from Manitoba south in winter to South Carolina, Florida, and Texas.

#### SUBGENUS **AMMODRAMUS.**

*Ammodramus* SWAINS. Zool. Jour. III. 1827, 348. Type,  
*Oriolus caudacutus* GMEL.

549. ***Ammodramus caudacutus*** (GMEL.).  
Sharp-tailed Sparrow.

*Oriolus caudacutus* GMEL. S. N. I. i. 1788, 394.

*Ammodramus caudacuta* SWAINS. Classif. B. II. 1837, 289.

[B 341, C 166, R 201, C 240.]

GEOG. DIST.—Salt marshes of the Atlantic coast, from southern New England to Maryland, and south in winter to the Gulf coast.

- 549 *a.* ***Ammodramus caudacutus nelsoni*** ALLEN.  
Nelson's Sparrow.

*Ammodromus caudacutus* var. *nelsoni* ALLEN, Pr. Bost. Soc. Nat. Hist. XVII. March, 1875, 293.

[B —, C —, R 201a, C 241.]

GEOG. DIST.— Fresh marshes of the interior, from northern Illinois northward to North Dakota and Manitoba; south in winter to Texas; in migrations visits the Atlantic coast (New England and Lower Hudson Valley to Charleston, S. C.). Accidental in California.

549 b. ***Ammodramus caudacutus subvirgatus*** DWIGHT.

Acadian Sharp-tailed Sparrow.

*Ammodramus caudacutus subvirgatus* DWIGHT, Auk, IV. July, 1887, 233.

[B —, C —, R 201, part, C 240, part.]

GEOG. DIST.— Coast of southern New Brunswick, Prince Edward Island (and probably Nova Scotia), and southward in migration to South Carolina.

550. ***Ammodramus maritimus*** (WILS.).

Seaside Sparrow.

*Fringilla maritima* WILS. Am. Orn. VII. 1811, 68, pl. 24, fig. 2.

*Ammodramus maritimus* SWAINS. Classif. B. II. 1837, 289.

[B 342, C 165, R 202, C 238.]

GEOG. DIST.— Salt marshes of the Atlantic coast, from Connecticut southward to Georgia. Accidental in Massachusetts.

550 a. ***Ammodramus maritimus peninsulæ*** ALLEN.

Scott's Seaside Sparrow.

*Ammodramus maritimus peninsulæ* ALLEN, Auk, V. July, 1888, 284.

[B 342, part, C 165, part, R 202, part, C 238, part.]

GEOG. DIST.— South Carolina to northern Florida; Gulf coast from Florida to Texas.

550 b. ***Ammodramus maritimus sennetti*** ALLEN.

Texas Seaside Sparrow.

*Ammodramus maritimus sennetti* ALLEN, Auk, V. July, 1888, 286.

[B —, C —, R 202, part, C 238, part.]

GEOG. DIST.— Coast of Texas (Corpus Christi).

551. ***Ammodramus nigrescens* RIDGW.**

Dusky Seaside Sparrow.

*Ammodromus maritimus* var. *nigrescens* RIDGW. Bull. Essex Inst. V. Dec. 1873, 198.*Ammodramus nigrescens* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880, 178.[B —, C 165*a*, R 203, C 239.]

GEOG. DIST.— Salt Lake and Merritt Island, eastern Florida.

GENUS **CHONDESTES** SWAINSON.*Chondestes* SWAINS. Phil. Mag. I. 1827, 435. Type, *C. strigatus* SWAINS.552. ***Chondestes grammacus* (SAY).**

Lark Sparrow.

*Fringilla grammaca* SAY, LONG's Exp. II. 1823, 139.*Chondestes grammaca* BONAP. Geog. & Comp. List, 1838, 32.[B 344, *part*, C 186, *part*, R 204, C 281, *part*.]

GEOG. DIST.— Southern Ontario and Mississippi Valley region, from Ohio, Illinois, and Michigan to the Plains, south to eastern Texas and northwestern Alabama. Accidental near the Atlantic coast (Massachusetts, Long Island, New Jersey, Washington, D. C., Florida).

552 *a*. ***Chondestes grammacus strigatus* (SWAINS.).**

Western Lark Sparrow.

*Chondestes strigatus* SWAINS. Phil. Mag. I. 1827, 435.*Chondestes grammaca strigata* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880, 179.[B 344, *part*, C 186, *part*, R 204*a*, C 281, *part*.]

GEOG. DIST.— Western United States, from the Plains to the Pacific coast, north to British Columbia and Manitoba, south through Lower California and Mexico to Guatemala.

GENUS **ZONOTRICHIA** SWAINSON.

*Zonotrichia* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831, 493. Type, by elimination, *Emberiza leucophrys* FORST.

553. **Zonotrichia querula** (NUTT.).

Harris's Sparrow.

*Fringilla querula* NUTT. Man. I. 2d ed. 1840, 555.

*Zonotrichia querula* GAMB. Journ. Ac. Nat. Sci. Phila. 2d ser. I. 1847, 51.

[B 348, C 185, R 205, C 280.]

GEOG. DIST.—Middle United States, from Illinois, Missouri, and Iowa west to middle Kansas and the Dakotas, and from Texas north to Manitoba. Accidental on Vancouver Island and in British Columbia and Oregon.

554. **Zonotrichia leucophrys** (FORST.).

White-crowned Sparrow.

*Emberiza leucophrys* FORST. Philos. Trans. LXII. 1772, 426.

*Z[onotrichia] leucophrys* SWAINS. in SW. & RICH. Fauna. Bor. Am. II. 1831, 493.

[B 345, C 183, R 206, C 276.]

GEOG. DIST.—North America at large, breeding chiefly in the Rocky Mountains, the Sierra Nevada and northeast to Labrador. South in winter to the Valley of Mexico.

554a. **Zonotrichia leucophrys intermedia** RIDGW.

Intermediate Sparrow.

*Zonotrichia leucophrys* var. *intermedia* RIDGW. Bull. Essex Inst. V. Dec. 1873, 198.

[B 346, *part*, C 183*b*, R 207*a*, C 277.]

GEOG. DIST.—Western North America, from the Rocky Mountains to the Pacific, and from Lower California and Mazatlan, Mexico, to Alaska. Breeds, so far as known, mainly north of the United States.

554 *b. Zonotrichia leucophrys gambelii* (NUTT.).

Gambel's Sparrow.

*Fringilla gambelii* NUTT. Man. I. 2d ed. 1840, 556.*Zonotrichia leucophrys* var. *gambeli* COUES, Key, 1872, 145.[B 346, *part*, C 183*a*, R 207, C 278.]

GEOG. DIST.—Pacific coast region, from British Columbia southward into Lower California.

557. *Zonotrichia coronata* (PALL.).

Golden-crowned Sparrow.

*Emberiza coronata* PALL. Zoog. Rosso-As. II. 1826, 44.*Zonotrichia coronata* BAIRD, B. N. Am. 1858, 461.

[B 347, C 184, R 208, C 279.]

GEOG. DIST.—Pacific coast region, from Alaska to southern California. Casually to Guadalupe Island, and in Wisconsin.

558. *Zonotrichia albicollis* (GMEL.).

White-throated Sparrow.

*Fringilla albicollis* GMEL. S. N. I. ii. 1788, 926.*Zonotrichia albicollis* SWAINS. Classif. B. II. 1837, 288.

[B 349, C 182, R 209, C 275.]

GEOG. DIST.—Chiefly eastern North America, west to the Plains, north to Labrador and the Fur Countries. Breeds from Montana, northern Wyoming, northern Michigan, northern New York, and northern New England northward, and winters from Massachusetts southward. Accidental in Utah, California and Oregon.

GENUS **SPIZELLA** BONAPARTE.*Spizella* BONAP. Saggio Distr. Met. 1832, 140. Type, *Fringilla pusilla* WILS.559. *Spizella monticola* (GMEL.).

Tree Sparrow.

*Fringilla monticola* GMEL. S. N. I. ii. 1788, 912.*Spizella monticola* BAIRD, B. N. Am. 1858, 472.



[B 357, *part*, C 177, *part*, R 210, *part*, C 268, *part*.]

GEOG. DIST.— Eastern North America, west to the Plains, and from the Arctic Ocean south, in winter, to the Carolinas, Kentucky, and eastern Kansas. Breeds north of the United States, east of the Rocky Mountains.

559 a. ***Spizella monticola ochracea*** BREWST.

Western Tree Sparrow.

*Spizella monticola ochracea* BREWST. Bull. Nutt. Orn. Club, VII.  
Oct. 1882, 228.

[B 357, *part*, C 177, *part*, R 210, *part*, C 268, *part*.]

GEOG. DIST.— Western North America, east to the Dakotas and western Kansas, south in winter to New Mexico and Arizona, north to the arctic regions ; breeds in Alaska.

560. ***Spizella socialis*** (WILS.).

Chipping Sparrow.

*Fringilla socialis* WILS. Am. Orn. II. 1810, 127, pl. 16, fig. 5.

*Spizella socialis* BONAP. Geog. & Comp. List, 1838, 33.

[B 359, *part*, C 178, R 211, C 269.]

GEOG. DIST.— Eastern North America, west to the Rocky Mountains, north to Great Slave Lake, and south to eastern Mexico, breeding from the Gulf States northward.

560 a. ***Spizella socialis arizonæ*** COUES.

Western Chipping Sparrow.

*Spizella socialis* var. *arizonæ* COUES, Key, 1872, 143.

[B 359, *part*, C 178a, R 211a, C 270.]

GEOG. DIST.— Western United States, from the Rocky Mountains to the Pacific, south in winter to central and western Mexico and Cape St. Lucas.

561. ***Spizella pallida*** (SWAINS.).

Clay-colored Sparrow.

*Emberiza pallida* SWAINS. in SW. & RICH. Fauna Bor. Am. II.  
1831, 251.

*Spizella pallida* BONAP. Geog. & Comp. List, 1838, 33.



[B 360, C 180, R 212, C 272.]

GEOG. DIST.—Interior of North America, from Illinois and Iowa west to the Rocky Mountains, Arizona, and Cape St. Lucas, and from Guanajuato and Oaxaca north to the Saskatchewan Plains. Breeds from Iowa and Nebraska northward.

562. **Spizella breweri** CASS.

Brewer's Sparrow.

*Spizella breweri* CASS. Pr. Ac. Nat. Sci. Phila. Feb. 1856, 40.

[B 361, C 180a, R 213, C 273.]

GEOG. DIST.—Western United States, north to Montana and British Columbia, south to Cape St. Lucas and Durango. Breeds throughout its United States range. Accidental in Massachusetts.

563. **Spizella pusilla** (WILS.).

Field Sparrow.

*Fringilla pusilla* WILS. Am. Orn. II. 1810, 121, pl. 16, fig. 2.

*Spizella pusilla* BONAP. Geog. & Comp. List, 1838, 33.

[B 358, C 179, R 214, C 271.]

GEOG. DIST.—Eastern United States and southern Canada, west to the Plains, south to the Gulf States and Texas. Breeds from South Carolina, southern Illinois and Kansas northward.

563 a. **Spizella pusilla arenacea** CHADB.

Western Field Sparrow.

*Spizella pusilla arenacea* CHADB. Auk, III. April, 1886, 248.

[B—, C—, R—, C—.]

GEOG. DIST.—Great Plains, from Texas to Montana and Dakota. Casual at New Orleans, La.

564. **Spizella wortheni** RIDGW.

Worthen's Sparrow.

*Spizella wortheni* RIDGW. Pr. U. S. Nat. Mus. VII. Aug. 22, 1884, 259.

[B—, C—, R—, C—.]

GEOG. DIST.—New Mexico (Silver City), and Chalchicomula, Puebla.

565. **Spizella atrigularis** (CAB.).

Black-chinned Sparrow.

*Spinites atrigularis* CAB. Mus. Hein. I. 1851, 133.*Spizella atrigularis* BAIRD, B. N. Am. 1858, 476.

[B 362, C 181, R 215, C 274.]

GEOG. DIST.—Southern California (north in the deserts to lat. 37°), Arizona, and southern New Mexico, south to Cape St. Lucas and on the tablelands of Mexico to Puebla.

GENUS **JUNCO** WAGLER.

*Junco* WAGLER, Isis, 1831, 526. Type, *J. phæonotus* WAGL. =  
*Fringilla cinerea* SWAINS.

566. **Junco aikenii** RIDGW.

White-winged Junco.

*Junco hyemalis* var. *aikenii* RIDGW. Am. Nat. VII. Oct. 1873, 612,  
614.

*Junco aikenii* RIDGW. Field & Forest, May, 1877, 198.

[B—, C 174a, R 216, C 262.]

GEOG. DIST.—Colorado, north to the Black Hills, where it breeds.

567. **Junco hyemalis** (LINN.).

Slate-colored Junco.

*Fringilla hyemalis* LINN. S. N. ed. 10, I. 1758, 183.*Junco hyemalis* SCL. P. Z. S. 1857, 7.

[B 354, C 174, R 217, C 261.]

GEOG. DIST.—North America, chiefly east of the Rocky Mountains, breeding from the higher parts of the Alleghanies, the Catskills, and the mountainous parts of southern New England northward. South in winter to the Gulf States. Casual in California and Arizona.

567 a. **Junco hyemalis oregonus** (TOWNS.).

Oregon Junco.

*Fringilla oregana* [err. typ.] TOWNS. Journ. Ac. Nat. Sci. Phila.  
VII. 1837, 188.

*Junco hyemalis* var. *oregonus* RIDGW. Am. Nat. VII. Oct. 1873, 612.

[B 352, C 175, R 218, C 263.]

GEOG. DIST.—Pacific coast, from Oregon north to Alaska, south in winter into California and Arizona.

567 *b. Junco hyemalis shufeldti* COALE.

Shufeldt's Junco.

*Junco hyemalis shufeldti* COALE, Auk, IV. Oct. 1887, 330.

[B 352, *part*, C 175, *part*, R 218, *part*, C 263, *part*.]

GEOG. DIST.—Rocky Mountain region, west in the mountains of the Great Basin to California; in winter south to Arizona, New Mexico, Texas, and northern Mexico. Accidental in Michigan, Illinois, Massachusetts, Maryland, etc.

567 *c. Junco hyemalis thurberi* ANTHONY.

Thurber's Junco.

*Junco hyemalis thurberi* ANTHONY, Zoe, I. Oct. 1890, 238.

[B 352, *part*, C 175, *part*, R 218, *part*, C 263, *part*.]

GEOG. DIST.—Sierra Nevada, and deserts and southern coast ranges of California.

567 *d. Junco hyemalis pinosus* LOOMIS.

Point Pinos Junco.

*Junco pinosus* LOOMIS, Auk, X. April, 1893, 47.

[B —, C —, R —, C —.]

GEOG. DIST.—Vicinity of Monterey, California.

567 *e. Junco hyemalis carolinensis* BREWST.

Carolina Junco.

*Junco hyemalis carolinensis* BREWST. Auk, III. Jan. 1886, 108.

[B 354, *part*, C 174, *part*, R 217, *part*, C 261, *part*.]

GEOG. DIST.—Southern Alleghanies (Western North Carolina, etc.).

568. *Junco annectens* BAIRD.

Pink-sided Junco.

*Junco annectens* BAIRD, Orn. Cal. I. 1870, 564.

[B —, C —, R 219, C 264.]

GEOG. DIST.— Rocky Mountain region, from Idaho and Montana south, in winter, to Arizona, New Mexico and northern Mexico.

568.1. **Junco ridgwayi** MEARNs.

**Ridgway's Junco.**

*Junco ridgwayi* MEARNs, Auk, VII. July, 1890, 243.

[B —, C —, R —, C —.]

GEOG. DIST.— Wyoming, Colorado, Arizona, and New Mexico.

569. **Junco caniceps** (WOODH.).

**Gray-headed Junco.**

*Struthus caniceps* WOODH. Pr. Ac. Nat. Sci. Phila. Dec. 1852, 202.

*Junco caniceps* BAIRD, B. N. Am. 1858, 468.

[B 353, C 176, R 220, C 265.]

GEOG. DIST.— Rocky Mountain region, from the Black Hills to the Wahsatch and Uintah Mountains, south to New Mexico, Arizona, and northern Mexico.

570. **Junco phæonotus palliatus** RIDGW.

**Arizona Junco.**

*Junco cinereus palliatus* RIDGW. Auk, II. Oct. 1885, 364.

*Junco phæonotus palliatus* RIDGW. Auk, XII. Oct. 1895, 391.

[B 350, *part*, C —, R 222, *part*, C 267, *part*.]

GEOG. DIST.— Mountains of southern Arizona, and southward into northern Mexico.

570 a. **Junco phæonotus dorsalis** (HENRY).

**Red-backed Junco.**

*Junco dorsalis* HENRY, Pr. Ac. Nat. Sci. Phila. 1858, 117.

*Junco phæonotus dorsalis* RIDGW. Auk, XII. Oct. 1895, 391.

[B 351, C —, R 221, C 266.]

GEOG. DIST.— Mountains of New Mexico and eastern Arizona, south into northern Mexico.

571. **Junco bairdi** BELDING.

Baird's Junco.

*Junco bairdi* BELDING, Pr. U. S. Nat. Mus. VI. Oct. 5, 1883, 155.

[B—, C—, R—, C—.]

GEOG. DIST.—Southern Lower California.

571.1. **Junco townsendi** ANTHONY.

Townsend's Junco.

*Junco townsendi* ANTHONY, Pr. Cal. Ac. Sci. 2d. ser. II. Oct. 11, 1889, 76.

[B—, C—, R—, C—.]

GEOG. DIST.—San Pedro Martir Mountains, Lower California.

572. **Junco insularis** RIDGW.

Guadalupe Junco.

*Junco insularis* RIDGW. Bull. U. S. Geol. & Geog. Surv. Terr. II. No. 2, April 1, 1876, 188.

[B—, C—, R 223, C—.]

GEOG. DIST.—Guadalupe Island, Lower California.

GENUS **AMPHISPIZA** COUES.*Amphispiza* COUES, B. Northwest, 1875, 234. Type, *Emberiza bilineata* CASS.573. **Amphispiza bilineata** (CASS.).

Black-throated Sparrow.

*Emberiza bilineata* CASSIN, Pr. Ac. Nat. Sci. Phila. Oct. 1850, 104, pl. 3.*Amphispiza bilineata* COUES, B. Northwest, 1875, 234.

[B 355, C 172, R 224, C 258.]

GEOG. DIST.—Western United States, from western Texas and Oklahoma west to the west slope of the Sierra Nevada, north throughout the Great Basin, and south in Mexico to Cape St. Lucas and San Luis Potosi. Breeds throughout its range.

574. ***Amphispiza belli*** (CASS.).

Bell's Sparrow.

*Emberiza belli* CASSIN, Pr. Ac. Nat. Sci. Phila. Oct. 1850, 104,  
pl. 4.*Amphispiza bellii* COUES, B. Northwest, 1875, 234.

[B 356, C 173, R 225, C 259.]

GEOG. DIST.—California west of the Sierra Nevada, north to about Lat. 38°, south to Lower California. Breeds throughout most of its range.

574 a. ***Amphispiza belli nevadensis*** (RIDGW.).

Sage Sparrow.

*Poospiza belli* var. *nevadensis* RIDGW. Bull. Essex Inst. V. Nov.  
1873, 191.*Amphispiza bellii* var. *nevadensis* COUES, B. Northwest, 1875, 234.

[B —, C 173a, R 225a, C 260.]

GEOG. DIST.—Great Basin, from Oregon and Idaho south to southern Arizona and New Mexico, breeding throughout its range.

574 b. ***Amphispiza belli cinerea*** TOWNSEND.

Gray Sage Sparrow.

*Amphispiza belli cinerea* TOWNSEND, Pr. U. S. Nat. Mus. XIII.  
1890, 136.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California.

GENUS **PEUCÆA** AUDUBON.*Peucæa* AUD. Synop. 1839, 112. Type, *Fringilla bachmanii* AUD.575. ***Peucæa æstivalis*** (LICHT.).

Pine-woods Sparrow.

*Fringilla æstivalis* LICHT. Verz. Doubl. 1823, 25.*Peucæa æstivalis* CAB. Mus. Hein. I. 1850, 132.[B 370, *part*, C 170, *part*, R 226, C 251.]

GEOG. DIST.—Florida and southern Georgia, migrating to southern Florida in winter.



575 a. **Peucæa æstivalis bachmanii** (AUD.).

Bachman's Sparrow.

*Fringilla bachmanii* AUD. Orn. Biog. II. 1834, 366, pl. 165.*Peucæa æstivalis bachmani* BREWST. Auk, II. Jan. 1885, 106.[B 370, *part*, C 170, *part*, R 226 a, C 252.]

GEOG. DIST.—North and South Carolina, Georgia, and Alabama, west to Texas, and north to southern Illinois and southern Indiana. Florida in winter.

576. **Peucæa arizonæ** RIDGW.

Arizona Sparrow.

*Peucæa æstivalis* var. *arizonæ* RIDGW. Am. Nat. VII. Oct. 1873, 615.*Peucæa arizonæ* RIDGW. Pr. U. S. Nat. Mus. I. Aug. 15, 1878, 127.

[B —, C 170 a, R 227, C 253.]

GEOG. DIST.—Southern Arizona and Sonora.

577. **Peucæa mexicana** (LAWR.).

Mexican Sparrow.

*Coturniculus mexicanus* LAWR. Ann. Lyc. N. Y. VIII. May, 1867, 474. (Mts. of Colima.)*Peucæa mexicana* RIDGW. Pr. U. S. Nat. Mus. VIII. No. 7, May 23, 1885, 99.

[B —, C —, R —, C —.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas south into central and western Mexico.

578. **Peucæa cassini** (WOODH.).

Cassin's Sparrow.

*Zonotrichia cassini* WOODH. Pr. Ac. Nat. Sci. Phila. April, 1852, 60.*Peucæa cassini* BAIRD, B. N. Am. 1858, 485.[B 371, C 170 *bis*, R 228, C 254.]

GEOG. DIST.—Central and western Kansas, southward and westward through Texas, New Mexico, Arizona and southern Nevada.

579. ***Peucaea carpalis*** COUES.

Rufous-winged Sparrow.

*Peucaea carpalis* COUES, Am. Nat. VII. June, 1873, 322.

[B —, C 171 bis, R 229, C 257.]

GEOG. DIST.— Arizona and Sonora.

580. ***Peucaea ruficeps*** (CASS.).

Rufous-crowned Sparrow.

*Ammodromus ruficeps* CASS. Pr. Ac. Nat. Sci. Phila. Oct. 1852, 184.

*Peucaea ruficeps* BAIRD, B. N. Am. 1858, 486.

[B 372, C 171, R 230, C 255.]

GEOG. DIST.— Coast of California, from Lat. 40° south to Cape St. Lucas.

580 a. ***Peucaea ruficeps boucardi*** (SCL.).

Boucard's Sparrow.

*Zonotrichia boucardi* SCLATER, P. Z. S. 1867, 1, pl. i.

*Peucaea ruficeps boucardi* RIDGW. Hist. N. Am. B. II. 1874, 38.

[B —, C —, R 230a, C 256.]

GEOG. DIST.— Southern New Mexico and southern Arizona, south into Mexico to Puebla.

580 b. ***Peucaea ruficeps eremæca*** BROWN.

Rock Sparrow.

*Peucaea ruficeps eremæca* BROWN, Bull. Nutt. Orn. Cl. VII. Jan. 1882, 26.

[B —, C —, R —, C —.]

GEOG. DIST.— Southwestern Texas, south in eastern Mexico to Orizaba.

GENUS **MELOSPIZA** BAIRD.

*Melospiza* BAIRD, B. N. Am. 1858, 478. Type, *Fringilla melodia* WILS. = *F. fasciata* GMEL.

**581. *Melospiza fasciata* (GMEL.).**

Song Sparrow.

*Fringilla fasciata* GMEL. S. N. I. ii. 1788, 922.*Melospiza fasciata* SCOTT, Am. Nat. X. 1876, 18.

[B 363, C 169, R 231, C 244.]

GEOG. DIST.—Eastern United States to the Plains, breeding from Virginia and the southern portion of the Lake States northward to the Fur Countries.

**581 a. *Melospiza fasciata fallax* (BAIRD).**

Desert Song Sparrow.

*Zonotrichia fallax* BAIRD, Pr. Ac. Nat. Sci. Phila. June, 1854, 119  
(nec *Melospiza fallax* auctorum plurimorum!).

*Melospiza fasciata fallax* HENSH. Auk, I. July, 1884, 224.

[B 367, C 169a, part, R 231a, part, C 245, part.]

GEOG. DIST.—Parts of New Mexico, Arizona, southern Nevada, and southwestern Utah.

**581 b. *Melospiza fasciata montana* HENSH.**

Mountain Song Sparrow.

*Melospiza fasciata montana* HENSHAW, Auk, I. July, 1884, 224.

[B —, C 169a, part, R 231a, part, C 245, part.]

GEOG. DIST.—Northern Mexico, mountains of New Mexico, Colorado, Utah, Nevada, and northward.

**581 c. *Melospiza fasciata heermanni* (BAIRD).**

Heerman's Song Sparrow.

*Melospiza heermanni* BAIRD, B. N. Am. 1858, 478.

*Melospiza fasciata* δ. *heermanni* RIDGW. Bull. Nutt. Orn. Cl. III.  
April, 1878, 66.

[B 364, C 169d, R 231b, C 248.]

GEOG. DIST.—California, east into western Nevada.

**581 d. *Melospiza fasciata samuelis* (BAIRD).**

Samuels's Song Sparrow.

*Ammodromus samuelis* BAIRD, B. N. Am. 1858, 455.

*Melospiza fasciata samuelis* RIDGW. Pr. U. S. Nat. Mus. III. Mar.  
1880, 3.

[B 343, 365, C 169e, R 231c, C 249.]

GEOG. DIST.—Coast region of California.

581 e. **Melospiza fasciata guttata** (NUTT.).

Rusty Song Sparrow.

*Fringilla guttata* NUTTALL, Man. Orn. I. ed. 2, 1840, 581.

*Melospiza fasciata* β. *guttata* RIDGW. Bull. Nutt. Orn. Cl. III.

April, 1878, 66.

[B—, C 169b, R 231d, C 246.]

GEOG. DIST.—Coast region of Washington and Oregon, south in winter to San Francisco, California.

581 f. **Melospiza fasciata rufina** (BONAP.).

Sooty Song Sparrow.

*Passerella rufina* BONAP. Consp. Av. I. July 15, 1850, 477.

*Melospiza fasciata rufina* RIDGW. Pr. U. S. Nat. Mus. III. Mar.

1880, 3.

[B 366, C 169c, R 231e, C 247.]

GEOG. DIST.—Coast region of British Columbia, north to Sitka.

581 g. **Melospiza fasciata rivularis** BRYANT.

Brown's Song Sparrow.

*Melospiza fasciata rivularis* BRYANT, Proc. Cal. Ac. Sci. 2d ser.

I. Sept. 29, 1888, 197.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California, from Comodu northward.

581 h. **Melospiza fasciata graminea** TOWNSEND.

Santa Barbara Song Sparrow.

*Melospiza fasciata graminea* TOWNSEND, Pr. U. S. Nat. Mus.

XIII. 1890, 139.

[B —, C —, R —, C —.]

GEOG. DIST.—Santa Barbara Island, and, in winter, adjacent mainland of California.

581 *i.* **Melospiza fasciata clementæ** TOWNSEND.

San Clemente Song Sparrow.

*Melospiza fasciata clementæ* TOWNSEND, Pr. U. S. Nat. Mus.  
XIII. 1890, 139.

[B —, C —, R —, C —.]

GEOG. DIST.— San Clemente and Santa Rosa Islands, California.

581.1. **Melospiza insignis** BAIRD.

Bischoff's Song Sparrow.

*Melospiza insignis* BAIRD, Trans. Chicago Acad. Sci. I. 1869,  
319, pl. xxix, fig. 2.[B —, C 169 *f*, R 232, *part*, C 250, *part*.]

GEOG. DIST.— Kadiak Island, Alaska.

582. **Melospiza cinerea** (GMEL.).

Aleutian Song Sparrow.

*Fringilla cinerea* GMEL. S. N. I. ii. 1788, 922.*Melospiza cinerea* FINSCH, Abhandl. Nat. Ver. Bremen, III.  
1872, 20.[B —, C 169 *f*, R 232, C 250.]

GEOG. DIST.— Aleutian and Pribilof Islands (except Kadiak Island), and east to Fort Kenai, Alaska.

583. **Melospiza lincolnii** (AUD.).

Lincoln's Sparrow.

*Fringilla lincolnii* AUD. Orn. Biog. II. 1834, 539, pl. 193.*Melospiza lincolni* BAIRD, B. N. Am. 1858, 482.

[B 368, C 167, R 234, C 242.]

GEOG. DIST.— North America at large, breeding chiefly north of the United States (as far north as Fort Yukon) and in the higher parts of the Rocky Mountains and Sierra Nevada; south, in winter, to Panama.

583 *a.* **Melospiza lincolnii striata** BREWST.

Forbush's Sparrow.

*Melospiza lincolni striata* BREWST. Auk, VI. April, 1889, 89.

[B 368, *part*, C 167, *part*, R 234, *part*, C 242, *part*.]

GEOG. DIST.—British Columbia.

584. **Melospiza georgiana** (LATH.).

Swamp Sparrow.

*Fringilla georgiana* LATH. Ind. Orn. I. 1790, 460.

*Melospiza georgiana* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 355.

[B 369, C 168, R 233, C 243.]

GEOG. DIST.—Eastern North America to the Plains, accidentally to Utah, north to the British Provinces, including Newfoundland and Labrador. Breeds from the Northern States northward, and winters from Massachusetts southward to the Gulf States.

GENUS **PASSERELLA** SWAINSON.

*Passerella* SWAINS. Classif. B. II. 1837, 288. Type, *Fringilla iliaca* MERR.

585. **Passerella iliaca** (MERR.).

Fox Sparrow.

*Fringilla iliaca* MERREM, "Beitr. zur besond. Gesch. der Vögel, II. 1786–87, 40, pl. x."

*Passerella iliaca* SWAINS. Classif. B. II. 1837, 288.

[B 374, C 188, R 235, C 282.]

GEOG. DIST.—Eastern North America, west to the Plains and Alaska (valley of the Yukon to the Pacific), and from the Arctic coast south to the Gulf States. Breeds north of the United States; winters chiefly south of the Potomac and Ohio Rivers. Accidental in winter in California.

585a. **Passerella iliaca unalaschcensis** (GMEL.).

Townsend's Sparrow.

*Emberiza unalaschcensis* GMEL. S. N. I. ii. 1788, 875.

*Passerella iliaca unalaschcensis* RIDGW. Pr. U. S. Nat. Mus. III.

Mar. 1880, 3.



[B 375, C 189, R 235*a*, C 283.]

GEOG. DIST.—Pacific coast region, from Kadiak south, in winter, to southern California, and casually to Guadalupe Island, Lower California. Breeds north of the United States.

585 *b*. ***Passerella iliaca megarhyncha*** (BAIRD).

Thick-billed Sparrow.

*Passerella megarhynchus* BAIRD, B. N. Am. 1858, 925.

*Passerella iliaca* ♂. *megarhyncha* BELDING, Pr. U. S. Nat. Mus. I. Mar. 1879, 418.

[B 376*a*, C —, R 235*b*, C 285.]

GEOG. DIST.—Sierra Nevada and Coast Range, California.

585 *c*. ***Passerella iliaca schistacea*** (BAIRD).

Slate-colored Sparrow.

*Passerella schistacea* BAIRD, B. N. Am. 1858, 490.

*Passerella iliaca* var. *schistacea* ALLEN, Bull. M. C. Z. III. 1872, 168.

[B 376, C 189*a*, R 235*c*, C 284.]

GEOG. DIST.—Rocky Mountain region of the United States, east in winter, to the Plains (Kansas), west to Nevada and California.

GENUS **EMBERNAGRA** LESSON.

*Embernagra* LESS. Traité, 1831, 465. Type, *E. dumetorum* LESS.  
= *Emberiza platensis* GMEL.

586. ***Embernagra rufivirgata*** LAWR.

Texas Sparrow.

*Embernagra rufivirgata* LAWR. Ann. Lyc. N. Y. V. May, 1851, 112, pl. 5, fig. 2.

[B 373, C 209, R 236, C 311.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas and eastern Mexico; casually north to the coast of Louisiana.

GENUS **PIPILO** VIEILLOT.

*Pipilo* VIEILL. Analyse, 1816, 32. Type, *Fringilla erythrophthalma* LINN.

587. **Pipilo erythrophthalmus** (LINN.).

Towhee.

*Fringilla erythrophthalma* LINN. S. N. ed. 10, I. 1758, 180.*Pipilo erythrophthalmus* VIEILL. Gal. Ois. I. 1824, 109, pl. 80.

[B 391, C 204, R 237, C 301.]

GEOG. DIST.—Eastern United States and southern Canada, west to the Plains, breeding from the lower Mississippi Valley and Georgia northward; in winter from the middle districts southward.

587 a. **Pipilo erythrophthalmus alleni** COUES.

White-eyed Towhee.

*Pipilo erythrophthalmus* var. *alleni* COUES, Am. Nat. V. Aug. 1871, 366.

[B—, C 204a, R 237a, C 302.]

GEOG. DIST.—Florida, and northward along the Atlantic coast to Charleston, South Carolina.

588. **Pipilo maculatus arcticus** (SWAINS.).

Arctic Towhee.

*Pyrgita* (*Pipilo*) *arctica* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831, 260, pll. 51, 52.

*Pipilo maculatus* var. *arcticus* COUES, Key, 1872, 152.

[B 393, C 205a, R 238, C 304.]

GEOG. DIST.—Plains of the Platte, Upper Missouri, Yellowstone and Saskatchewan Rivers, west to the eastern slope of the Rocky Mountains, south in winter to Kansas, Colorado, and Texas.

588 a. **Pipilo maculatus megalonyx** (BAIRD).

Spurred Towhee.

*Pipilo megalonyx* BAIRD, B. N. Am. 1858, 515.*Pipilo maculatus* var. *megalonyx* COUES, Key, 1872, 152.

[B 394, C 205b, R 238a, C 305.]

GEOG. DIST.—Rocky Mountain region of the United States, west to the Sierra Nevada and southern California, south into northern Mexico and Lower California.

588 *b. Pipilo maculatus oregonus* (BELL).

Oregon Towhee.

*Pipilo oregonus* BELL, Ann. Lyc. N. Y. V. 1852, 6.*Pipilo maculatus* var. *oregonus* COUES, Key, 1872, 152.[B 392, C 205, R 238*b*, C 303.]

GEOG. DIST.—Pacific coast region, from British Columbia south to San Francisco, and, in winter, to southern California.

589. *Pipilo consobrinus* RIDGW.

Guadalupe Towhee.

*Pipilo maculatus consobrinus* RIDGW. Bull. U. S. Geol. & Geog. Surv. Terr. II. No. 2, April 1, 1876, 189.

*Pipilo consobrinus* RIDGW. Bull. Nutt. Orn. Club, II. July, 1877, 60.[B —, C —, R 238*c*, C —.]

GEOG. DIST.—Guadalupe Island, Lower California.

590. *Pipilo chlorurus* (TOWNS.).

Green-tailed Towhee.

*Fringilla chlorura* TOWNS. in AUD. Orn. Biog. V. 1839, 336.*Pipilo chlorurus* BAIRD, B. N. Am. 1858, 519.

[B 398, C 208, R 239, C 310.]

GEOG. DIST.—Interior Plateau region of the United States, from the western border of the Plains to the Sierra Nevada, from central Idaho and Yellowstone National Park south to Cape St. Lucas, and on the tablelands of Mexico to Guanajuato.

591. *Pipilo fuscus mesoleucus* (BAIRD).

Cañon Towhee.

*Pipilo mesoleucus* BAIRD, Pr. Ac. Nat. Sci. Phila. June, 1854, 119.*Pipilo fuscus* var. *mesoleucus* RIDGW. Bull. Essex Inst. V. Nov. 1873, 183.

[B 397, C 206, R 240, C 306.]

GEOG. DIST.—Southern border of the United States, from the valley of the Upper Rio Grande west to the valley of the Gila, south into Sonora and Chihuahua.

591 *a.* **Pipilo fuscus albigula** (BAIRD).

Saint Lucas Towhee.

*Pipilo albigula* BAIRD, Pr. Ac. Nat. Sci. Phila. Nov. 1859, 305.*Pipilo fuscus* var. *albigula* COUES, Key, 1872, 152.[B —, C 206*a*, R 240*a*, C 307.]

GEOG. DIST.—Lower California, north to Lat. 30°.

591 *b.* **Pipilo fuscus crissalis** (VIG.).

California Towhee.

*Fringilla crissalis* VIG. Zool. Bloss. 1839, 19.*Pipilo fuscus* var. *crissalis* COUES, Key, 1872, 153.[B 396, C 206*b*, R 240*b*, C 308.]

GEOG. DIST.—Pacific coast region west of the Cascades and Sierra Nevada, from the Umpqua Valley, Oregon, south to northern Lower California.

591 *c.* **Pipilo fuscus senicula** ANTHONY.

Anthony's Towhee.

*Pipilo fuscus senicula* ANTHONY, Auk, XII. April, 1895, 111.[B 396, *part*, C 206*b*, *part*, R 240*b*, *part*, C 308, *part*.]

GEOG. DIST.—Southern California, and Lower California, south to Lat. 29°.

592. **Pipilo aberti** BAIRD.

Abert's Towhee.

*Pipilo aberti* BAIRD, STANSBURY'S Rep. Exped. Utah, 1852, 325.

[B 395, C 207, R 241, C 309.]

GEOG. DIST.—New Mexico and Arizona, north into southern Colorado, southwestern Utah, and southern Nevada.

GENUS **CARDINALIS** BONAPARTE.*Cardinalis* BONAP. P. Z. S. 1837 (June, 1838), 111. Type, *C.**virginianus* BONAP. = *Loxia cardinalis* LINN.

593. **Cardinalis cardinalis** (LINN.).

Cardinal.

*Loxia cardinalis* LINN. S. N. ed. 10, 1758, 172.*Cardinalis cardinalis* LICHT. Nomencl. Mus. Berol. 1854, 44.

[B 390, C 203, R 242, C 299.]

GEOG. DIST.—Eastern United States, north to the lower Hudson Valley and the Great Lakes, casually further north, and west to the Plains. Resident in Bermuda.

593 a. **Cardinalis cardinalis superbus** RIDGW.

Arizona Cardinal.

*Cardinalis cardinalis superbus* RIDGW. Auk, II. Oct. 1885, 344.

[B —, C 203a, part, R 242a, part, C 300, part.]

GEOG. DIST.—Southern Arizona and western Mexico, south to Mazatlan.

593 b. **Cardinalis cardinalis igneus** (BAIRD).

Saint Lucas Cardinal.

*Cardinalis igneus* BAIRD, Pr. Ac. Nat. Sci. Phila. 1859, 305.*Cardinalis cardinalis igneus* STEJN. Auk, I. 1884, 171.

[B —, C 203a, part, R 242a, part, C 300, part.]

GEOG. DIST.—Lower California, north to Lat. 29°.

593 c. **Cardinalis cardinalis canicaudus** CHAPM.

Gray-tailed Cardinal.

*Cardinalis cardinalis canicaudus* CHAPM. Bull. Am. Mus. Nat. Hist. III. Aug. 1891, 324.

[B 390, part, C 203, part, R 242, part, C 299, part.]

GEOG. DIST.—Southwestern Texas, south into northeastern Mexico.

GENUS **PYRRHULOXIA** BONAPARTE.

*Pyrrhuloxia* BONAP. Consp. Av. I. 1850, 500. Type, *Cardinalis sinuatus* BONAP.



594. **Pyrrhuloxia sinuata** BONAP.

Texas Cardinal.

*Cardinalis sinuatus* BONAP. P. Z. S. 1837 (June, 1838), 111.*Pyrrhuloxia sinuata* BONAP. Consp. Av. I. 1850, 500.

[B 389, C 202, R 243, C 298.]

GEOG. DIST.—Southern border of the United States, from the valley of the Lower Rio Grande south to San Luis Potosi, Puebla, etc. North casually to the coast of Louisiana.

594 a. **Pyrrhuloxia sinuata beckhami** RIDGW.

Arizona Pyrrhuloxia.

*Pyrrhuloxia sinuata beckhami* RIDGW. Auk, IV. Oct. 1887, 347.[B 389, *part*, C 202, *part*, R 243, *part*, C 298, *part*.]

GEOG. DIST.—Southern Arizona to western Texas, and south into northwestern Mexico.

594 b. **Pyrrhuloxia sinuata peninsulæ** RIDGW.

Saint Lucas Pyrrhuloxia.

*Pyrrhuloxia sinuata peninsulæ* RIDGW. Auk, IV. Oct. 1887, 347.[B 389, *part*, C 202, *part*, R 243, *part*, C 298, *part*.]

GEOG. DIST.—Lower California.

GENUS **HABIA** REICHENBACH.

*Habia* REICH. Syst. Av. June 1, 1850, pl. lxxviii. Type, *Guiraca melanocephala* SWAINS.

595. **Habia ludoviciana** (LINN.).

Rose-breasted Grosbeak.

*Loxia ludoviciana* LINN. S. N. ed. 12, I. 1766, 306.*Habia ludoviciana* STEJN. Auk, I. Oct. 1884, 367.

[B 380, C 193, R 244, C 289.]

GEOG. DIST.—Eastern United States and southern Canada, west to Manitoba and the eastern border of the Plains, breeding from Kansas and the mountains of the Carolinas northward; south, in winter, to Cuba, Central America, and northern South America.



596. **Habia melanocephala** (SWAINS.).

Black-headed Grosbeak.

*Guiraca melanocephala* SWAINS. Philos. Mag. I. 1827, 438.*Habia melanocephala* STEJN. Auk, I. Oct. 1884, 367.

[B 381, C 194, R 245, C 290.]

GEOG. DIST.—Western United States, from middle Kansas to the Pacific coast, north to British Columbia and Montana, and south in Mexico to Oaxaca and the Cape region of Lower California.

GENUS **GUIRACA** SWAINSON.

*Guiraca* SWAINS. Zool. Jour. III. Nov. 1827, 350. Type, *Loxia cærulea* LINN.

597. **Guiraca cærulea** (LINN.).

Blue Grosbeak.

*Loxia cærulea* LINN. S. N. ed. 10, I. 1758, 175.*Guiraca cærulea* SWAINS. Phil. Mag. I. 1827, 438.

[B 382, C 195, R 246, C 291.]

GEOG. DIST.—Eastern United States, from southern New Jersey, southern Illinois and central Nebraska, south to Cuba and Mexico. Casual northward to New England.

597 a. **Guiraca cærulea eurhyncha** COUES.

Western Blue Grosbeak.

*Guiraca cærulea*, var. *eurhyncha* COUES, Am. Nat. VIII. Sept. 1874, 563.

[B 382, *part*, C 195, *part*, R 246, *part*, C 291, *part*.]

GEOG. DIST.—Southwestern United States, from South Dakota, Colorado, southern Utah, southern Nevada and California to Lower California and southern Mexico.

GENUS **PASSERINA** VIEILLOT.

*Passerina* VIEILL. Analyse, 1816, 30. Type, by elimination, *Tanagra cyanea* LINN.

598. **Passerina cyanea** (LINN.).

Indigo Bunting.

*Tanagra cyanea* LINN. S. N. ed. 12, I. 1766, 315.

*Passerina cyanea* VIEILL. Nouv. Dict. d'Hist. Nat. XXV. 1817, 7.

[B 387, C 199, R 248, C 295.]

GEOG. DIST.—Eastern United States, west to Kansas, north to New Brunswick, southern Ontario and Minnesota; south in winter to Central America.

599. **Passerina amoena** (SAY).

Lazuli Bunting.

*Emberiza amoena* SAY, LONG'S Exp. II. 1823, 47.

*Passerina amoena* GRAY, Handl. II. 1870, 97.

[B 386, C 200, R 249, C 296.]

GEOG. DIST.—Western United States, from the Plains to the Pacific, north to the dry interior of British Columbia and Idaho, south to the Valley of Mexico.

600. **Passerina versicolor** (BONAP.).

Varied Bunting.

*Spiza versicolor* BONAP. P. Z. S. 1837 (June, 1838), 120.

*Passerina versicolor* GRAY, Handl. II. 1870, 97.

[B 385, C 197, R 250, C 293.]

GEOG. DIST.—From the valley of the Lower Rio Grande in Texas southward to Guatemala. Accidental in southern Michigan.

600 a. **Passerina versicolor pulchra** RIDGW.

Beautiful Bunting.

*Passerina versicolor pulchra* RIDGW. Man. N. Am. B. 1887, 448.

[B 385, *part*, C 197, *part*, R 250, *part*, C 293, *part*.]

GEOG. DIST.—Lower California and northwestern Mexico.

601. **Passerina ciris** (LINN.).

Painted Bunting.

*Emberiza ciris* LINN. S. N. ed. 10, I. 1758, 179.

*Passerina ciris* VIEILL. Nouv. Dict. d'Hist. Nat. XXV. 1817, 17.

[B 384, C 196, R 251, C 292.]

GEOG. DIST.—South Atlantic and Gulf States to western Texas, north to North Carolina and southern Illinois, and south to Panama.

GENUS **SPOROPHILA** CABANIS.

*Sporophila* CABANIS, Arch. f. Naturg. X. i. 1844, 291.

602. **Sporophila morelleti sharpei** LAWR.

Sharpe's Seed-eater.

*Sporophila morelleti sharpei* LAWR. Auk, VI. Jan. 1889, 53.

[B 388, C 200, R 252, C 296.]

GEOG. DIST.—Southeastern Texas and adjacent parts of Mexico.

GENUS **EUETHEIA** REICHENBACH.

*Euetheia* REICH. Av. Syst. Nat. June 1, 1850, pl. lxxix. Type,  
*Emberiza lepida* LINN.

[603.] **Euetheia bicolor** (LINN.).

Grassquit.

*Fringilla bicolor* LINN. S. N. ed. 12, I. 1766, 324.

*Euetheia bicolor* GUNDLACH, J. f. O. XXII. 1874, 312.

[B —, C 201, R 253, C 297.]

GEOG. DIST.—West Indies. Accidental or casual in southern Florida.

[603.1.] **Euetheia canora** (GMEL.).

Melodious Grassquit.

*Loxia canora* GMEL. S. N. I. ii. 1788, 858.

*Euetheia canora* BREWER, Proc. Bost. Soc. Nat. Hist. VII. 1860,  
307.

[B —, C —, R —, C —.]

GEOG. DIST.—Cuba. Accidental in southern Florida (Sombrero Key).

GENUS **SPIZA** BONAPARTE.

*Spiza* BONAP. Journ. Ac. Nat. Sci. Phila. IV. i. Aug. 1824, 45.

Type, *Emberiza americana* GMEL.

604. **Spiza americana** (GMEL.).

Dickeissel.

*Emberiza americana* GMEL. S. N. I. ii. 1788, 872.

*Spiza americana* RIDGW. Pr. U. S. Nat. Mus. III. March 27, 1880, 3.

[B 378, C 191, R 254, C 287.]

GEOG. DIST.—Eastern United States to the Rocky Mountains, breeding from Texas north to Massachusetts, New York, southern Ontario, Wisconsin, Minnesota, and North Dakota; south in winter through Central America to northern South America; southwest in migrations to Arizona. Now rare or entirely absent east of the Alleghanies.

GENUS **CALAMOSPIZA** BONAPARTE.

*Calamospiza* BONAP. Geog. & Comp. List, 1838, 30. Type,

*Fringilla bicolor* TOWNS. = *Calamospiza melanocorys* STEJN.

605. **Calamospiza melanocorys** STEJN.

Lark Bunting.

*Calamospiza melanocorys* STEJN. Auk, II. Jan. 1885, 49.

[B 377, C 190, R 256, C 286.]

GEOG. DIST.—From the Plains of middle Kansas north to Manitoba and Assiniboia, west to the Rocky Mountains, less commonly to the Pacific in southern California, and south to Guanajuato and Lower California. Accidental in Massachusetts, New York, and South Carolina.

FAMILY **TANAGRIDÆ**. TANAGERS.GENUS **EUPHONIA** DESMAREST.

*Euphonia* DESM. Hist. Nat. Tang. 1805, pl. xix. Type, *Pipra musica* GMEL.?

[606.] **Euphonia elegantissima** (BONAP.).

Blue-headed Euphonia.

*Pipra elegantissima* BONAP. P. Z. S. 1837 (June, 1838), 112.*Euphonia elegantissima* GRAY, Gen. B. App. 1849, 17.

[B 224, C —, R 160, C —.]

GEOG. DIST.—Eastern Mexico, and south to Veragua. Texas (GIRAUD).

GENUS **PIRANGA** VIEILLOT.*Piranga* VIEILL. Ois. Am. Sept. I. 1807, p. iv. Type, *Muscicapa rubra* LINN. 1766 = *Fringilla rubra* LINN. 1758.607. **Piranga ludoviciana** (WILS.).

Louisiana Tanager.

*Tanagra ludoviciana* WILS. Am. Orn. III. 1811, 27, pl. 20, fig. 1.*Pyranga ludoviciana* RICHARDSON, Rep. Brit. Ass. Adv. Sci. V. 1837, 175.

[B 223, C 110, R 162, C 158.]

GEOG. DIST.—Western United States, from the Great Plains to the Pacific, north to South Dakota and British Columbia. In winter south to Guatemala. Accidental in New York and New England.

[607.1.] **Piranga rubriceps** GRAY.

Gray's Tanager.

*Pyranga rubriceps* GRAY, Gen. B. II. 1844, pl. 89.

[B —, C —, R —, C —.]

GEOG. DIST.—Colombia. Accidental in California (Dos Pueblos, Santa Barbara Co.).

608. **Piranga erythromelas** VIEILL.

Scarlet Tanager.

*Pyranga erythromelas* VIEILL. Nouv. Dict. d'Hist. Nat. XXVIII.1819, 293 (= *Pyranga rubra* AUCT, nec *Fringilla rubra* LINN.).

[B 220, C 107, R 161, C 154.]

GEOG. DIST.—Eastern United States, west to the Plains, and north to southern Ontario and Manitoba. In winter the West Indies, eastern Mexico, Central America and northern South America.

609. **Piranga hepatica** SWAINS.

Hepatic Tanager.

*Piranga hepatica* SWAINS. Phil. Mag. I. 1827, 438.

[B 222, C 109, R 163, C 157.]

GEOG. DIST.—Southern New Mexico and southern Arizona southward to Guatemala.

610. **Piranga rubra** (LINN.).

Summer Tanager.

*Fringilla rubra* LINN. S. N. ed. 10, I. 1758, 181.

*Piranga rubra* VIEILL. Ois. Am. Sept. I. 1807, p. iv.

[B 221, C 108, R 164, C 155.]

GEOG. DIST.—Eastern United States, to the Plains, north to southern New Jersey and southern Illinois, casually north to Massachusetts and Ontario, and accidentally to Nova Scotia. In winter, Cuba, eastern Mexico, Central America, and northern South America to Peru.

610 a. **Piranga rubra cooperi** RIDGW.

Cooper's Tanager.

*Piranga cooperi* RIDGW. Pr. Ac. Nat. Sci. Phila. 1869, 130.

*Piranga rubra cooperi* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B —, C 108a, R 164a, C 156.]

GEOG. DIST.—New Mexico and Arizona, south in western Mexico to Colima.

FAMILY **HIRUNDINIDÆ**. SWALLOWS.

GENUS **PROGNE** BOIE.

*Progne* BOIE, Isis, 1826, 971. Type, *Hirundo subis* LINN.



**611. *Progne subis* (LINN.).****Purple Martin.***Hirundo subis* LINN. S. N. ed. 10, l. 1758, 192.*Progne subis* BAIRD, Rev. Am. B. I. May, 1865, 274.

[B 231, C 117, R 152, C 165.]

GEOG. DIST.—Temperate North America, north to Ontario and the Saskatchewan, south to the higher parts of Mexico, wintering in South America.

**611 a. *Progne subis hesperia* BREWST.****Western Martin.***Progne subis hesperia* BREWST. Auk, VI. April, 1889, 92.[B 231, *part*, C 117, *part*, R 152, *part*, C 165, *part*.]

GEOG. DIST.—California, south of Latitude 40°, and southern Arizona; in winter, south to Nicaragua.

**611.1. *Progne cryptoleuca* BAIRD.****Cuban Martin.***Progne cryptoleuca* BAIRD, Rev. Am. B. I. May, 1865, 277.

[B 231a, C —, R 152a, C —.]

GEOG. DIST.—Cuba, and southern Florida.

**GENUS PETROCHELIDON CABANIS.**

*Petrochelidon* CAB. Mus. Hein. I. 1850, 47. Type, *Hirundo melanogastra* SWAINS.

**612. *Petrochelidon lunifrons* (SAY).****Cliff Swallow.***Hirundo lunifrons* SAY, LONG'S Exp. II. 1823, 47.

*Petrochelidon lunifrons* CASSIN, Cat. Hirun. Mus. Phila. Acad. Nat. Sci. 1853, 4.

[B 226, C 114, R 153, C 162.]

GEOG. DIST.—North America, north to the limit of trees, breeding south to the valleys of the Potomac and the Ohio, southern Texas, southern Arizona, and California; Central and South America in winter. Not recorded from Florida or the West Indies.

[612.1.] **Petrochelidon fulva** (VIEILL.).

Cuban Cliff Swallow.

*Hirundo fulva* VIEILL. Ois. Am. Sept. I. 1807, 62, pl. 30.*Petrochelidon fulva* CAB. Mus. Hein. I. 1850, 47.

[B—, C—, R—, C—.]

GEOG. DIST.—Greater Antilles and coast of Central America.  
 Accidental on the Dry Tortugas, Florida.

GENUS **CHELIDON** FORSTER.

*Chelidon* FORST. Synop. Cat. Brit. B. 1817, 55. Type, *Hirundo rustica* LINN.

613. **Chelidon erythrogastra** (BODD.).

Barn Swallow.

*Hirundo erythrogaster* BODD. Tabl. P. E. 1783, 45.

*Chelidon erythrogastra* STEJN. Pr. U. S. Nat. Mus. V. June 5,  
 1882, 31.

[B 225, C 111, R 154, C 159.]

GEOG. DIST.—North America in general, breeding from the Fur Countries south into Mexico; visits the West Indies in migrations, and winters in Central America and South America.

GENUS **TACHYCINETA** CABANIS.

*Tachycineta* CAB. Mus. Hein. I. 1850, 48. Type, *Hirundo thalassina* SWAINS.

614. **Tachycineta bicolor** (VIEILL.).

Tree Swallow.

*Hirundo bicolor* VIEILL. Ois. Am. Sept. I. 1807, 61, pl. 31.*Tachycineta bicolor* CAB. Mus. Hein. I. 1850, 48.

[B 227, C 112, R 155, C 160.]

GEOG. DIST.—North America at large, breeding from the Fur Countries south to New Jersey, the Ohio Valley, Kansas, and Colorado, etc., wintering from South Carolina and the Gulf States southward to the West Indies and Guatemala.

615. **Tachycineta thalassina** (SWAINS.).

Violet-green Swallow.

*Hirundo thalassinus* SWAINS. Phil. Mag. I. 1827, 366.*Tachycineta thalassina* CAB. Mus. Hein. I. 1850, 48.

[B 228, C 113, R 156, C 161.]

GEOG. DIST.—Western United States, from the eastern base of the Rocky Mountains to the Pacific, north to British Columbia, south, in winter, to Guatemala and Costa Rica.

GENUS **CALlicHELIDON** BAIRD.*Callichelidon* BRYANT, MS. BAIRD, Rev. Am. Bds. I. 1865, 303.Type, *Hirundo cyaneoviridis* BRYANT.[615.1.] **Callichelidon cyaneoviridis** (BRYANT).

Bahaman Swallow.

*Hirundo cyaneoviridis* BRYANT, Pr. Boston Soc. Nat. Hist., VII. 1859, 111.*Callichelidon cyaneoviridis* BRYANT, MS. BAIRD, Rev. Am. Birds, I. 1865, 303.

[B —, C —, R —, C —.]

GEOG. DIST.—Bahamas. Accidental on the Dry Tortugas, Florida.

GENUS **CLIVICOLA** FORSTER.*Clivicola* FORST. Synop. Cat. Brit. B. 1817, 55. Type, *Hirundo riparia* LINN.616. **Clivicola riparia** (LINN.).

Bank Swallow.

*Hirundo riparia* LINN. S. N. ed. 10, I. 1758, 192.*Clivicola riparia* STEJN. Pr. U. S. Nat. Mus. V. 1882, 32.

[B 229, C 115, R 157, C 163.]

GEOG. DIST.—Northern hemisphere; in America, south to the West Indies, Central America, and northern South America, breeding from the middle districts of the United States northward to about the limit of trees.

GENUS **STELGIDOPTERYX** BAIRD.

*Stelgidopteryx* BAIRD, B. N. Am. 1858, 312. Type, *Hirundo serripennis* AUD.

617. **Stelgidopteryx serripennis** (AUD.).

Rough-winged Swallow.

*Hirundo serripennis* AUD. Orn. Biog. IV. 1838, 593.

*Stelgidopteryx serripennis* BAIRD, B. N. Am. 1858, 312.

[B 230, C 116, R 158, C 164.]

GEOG. DIST.—United States at large, north to Connecticut, southern Ontario, southern Minnesota, southern Montana, and British Columbia, south through Mexico to Costa Rica. Breeds throughout its United States range and south into Mexico.

FAMILY **AMPELIDÆ**. WAXWINGS, ETC.SUBFAMILY **AMPELINÆ**. WAXWINGS.GENUS **AMPELIS** LINNÆUS.

*Ampelis* LINN. S. N. ed. 12, I. 1766, 297. Type, by elimination, *Lanius garrulus* LINN.

618. **Ampelis garrulus** LINN.

Bohemian Waxwing.

*Lanius garrulus* LINN. S. N. ed. 10, I. 1758, 95.

*Ampelis garrulus* LINN. S. N. ed. 12, I. 1766, 297.

[B 232, C 118, R 150, C 166.]

GEOG. DIST.—Northern parts of the northern hemisphere. In North America, south in winter, irregularly, to Pennsylvania, Illinois, Kansas, southern Colorado, and northern California. Accidental at Fort Mohave, Arizona. Breeds north of the United States.

619. **Ampelis cedrorum** (VIEILL.).

Cedar Waxwing.

*Bombycilla cedrorum* VIEILL. Ois. Am. Sept. I. 1807, 88, pl. 57.

*Ampelis cedrorum* GRAY, Gen. B. I. 1846, 278.

[B 233, C 119, R 151, C 167.]

GEOG. DIST.—North America at large, from the Fur Countries southward. In winter, from the northern border of the United States south to the West Indies and Costa Rica. Breeds from Virginia, the southern Alleghanies, Kentucky, Kansas, Arizona, etc., northward.

## SUBFAMILY PTILIOGONATINÆ.

## GENUS PHAINOPEPLA SCLATER.

*Phainopepla* SCL. P. Z. S. 1858, 543. Type, *Ptiliogonys nitens* SWAINS.

620. *Phainopepla nitens* (SWAINS.).

*Phainopepla*.

*Ptiliogonys nitens* SWAINS. Anim. in Menag. 1838, 285.

*Phainopepla nitens* SCL. P. Z. S. 1858, 543.

[B 234, C 120, R 26, C 168.]

GEOG. DIST.—Southwestern United States, from southwestern Texas westward; north to southern Utah, Nevada, Fort Crook, California, and south to Cape St. Lucas and the Valley of Mexico.

## FAMILY LANIIDÆ. SHRIKES.

## GENUS LANIUS LINNÆUS.

*Lanius* LINN. S. N. ed. 10, I. 1758, 93. Type, by elimination, *L. excubitor* LINN.

621. *Lanius borealis* VIEILL.

Northern Shrike.

*Lanius borealis* VIEILL. Ois. Am. Sept. 1. 1807, 80, pl. 50.

[B 236, C 134, R 148, C 186.]

GEOG. DIST.—Northern North America, south in winter to the middle portions of the United States (Virginia, Kentucky, Kansas, Colorado, Arizona, northern California). Breeds north of the United States.

622. **Lanius ludovicianus** LINN.

Loggerhead Shrike.

*Lanius ludovicianus* LINN. S. N. ed. 12, I. 1766, 134.

[B 237, C 135, R 149, C 187.]

GEOG. DIST.— Eastern United States, west to the Plains; north to northern New England. Breeds from the Gulf States to Virginia and casually north, on the Atlantic coast, to southern New Jersey; in the interior, northward to the Great Lakes, and through western Pennsylvania and New York to New Hampshire, Vermont, and Maine.

622 *a.* **Lanius ludovicianus excubitorides** (SWAINS.).

White-rumped Shrike.

*Lanius excubitorides* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831, 115, pl. 34.

*Lanius ludovicianus* var. *excubitoroides* COUES, Key, 1872, 125.

[B 238, C 135*a*, R 149*a*, C 188.]

GEOG. DIST.— Western North America, from the eastern border of the Plains to the Pacific, except coast of California, and from Manitoba and the Plains of the Saskatchewan south over the tablelands of Mexico.

622 *b.* **Lanius ludovicianus gambeli** RIDGW.

California Shrike.

*Lanius ludovicianus gambeli* RIDGW. Man. N. Am. B. 1887, 467.

[B 238, *part*, C 153*a*, *part*, R 149*a*, *part*, C 188, *part*.]

GEOG. DIST.— Coast of California.

FAMILY **VIREONIDÆ**. VIREOS.GENUS **VIREO** VIEILLOT.SUBGENUS **VIREOSYLVA** BONAPARTE.

*Vireosylva* BONAP. Geog. & Comp. List, 1838, 26. Type, *Muscicapa olivacea* LINN.



**623. Vireo calidris barbatulus (CAB.).****Black-whiskered Vireo.***Phyllomanes barbatulus* CAB. J. f. O. 1855, 467.*Vireo calidris barbatulus* COUES, B. Col. Vall. 1878, 491.

[B 243, C 123, R 137, C 172.]

GEOG. DIST.—Bahamas, Cuba, and southern Florida.

**624. Vireo olivaceus (LINN.).****Red-eyed Vireo.***Muscicapa olivacea* LINN. S. N. ed. 12, I. 1766, 327.*Vireo olivaceus* BONAP. Ann. Lyc. N. Y. II. 1826, 71.

[B 240, C 122, R 135, C 170.]

GEOG. DIST.—Eastern North America, west to Colorado, Utah, and British Columbia; north to the arctic regions; south, in winter, from Florida to northern South America. Breeds nearly throughout its North American range.

**625. Vireo flavoviridis (CASS.).****Yellow-green Vireo.***Vireosylvia flavoviridis* CASS. Pr. Ac. Nat. Sci. Phila. V. Feb. 1851, 152.*Vireo flavoviridis* BAIRD, B. N. Am. 1858, 332.

[B 241, C —, R 136, C 171.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, southward to Panama, Ecuador, Peru and Upper Amazon. Accidental at Godbout, Province of Quebec, and at Riverside, California.

**626. Vireo philadelphicus (CASS.).****Philadelphia Vireo.***Vireosylvia philadelphica* CASS. Pr. Ac. Nat. Sci. Phila. V. Feb. 1851, 153, pl. 10, fig. 2.*Vireo philadelphicus* BAIRD, B. N. Am. 1858, 335.

[B 244, C 124, R 138, C 173.]

GEOG. DIST.—Eastern North America, north to Hudson Bay; south, in winter, to Costa Rica and Panama. Not recorded from Mexico or the West Indies. Breeds from Maine, New Hampshire, and Manitoba northward.

627. **Vireo gilvus** (VIEILL.).

Warbling Vireo.

*Muscicapa gilva* VIEILL. Ois. Am. Sept. I. 1807, 65, pl. 34.*Vireo gilvus* BONAP. Journ. Ac. Nat. Sci. Phila. IV. 1824, 176.

[B 245, C 125, 125a, R 139, 139a, C 174, 175.]

GEOG. DIST.—North America in general, from the Fur Countries to Oaxaca, Mexico. Breeds throughout the greater part of its range.

SUBGENUS **LANIVIREO** BAIRD.

*Lanivireo* BAIRD, Rev. Am. B. I. May, 1866, 345. Type, *Vireo flavifrons* VIEILL.

628. **Vireo flavifrons** VIEILL.

Yellow-throated Vireo.

*Vireo flavifrons* VIEILL. Ois. Am. Sept. I. 1807, 85, pl. 54.

[B 252, C 126, R 140, C 176.]

GEOG. DIST.—Eastern United States, north to Ontario and Manitoba; south, in winter, to Colombia. Breeds from Florida and the Gulf States northward.

629. **Vireo solitarius** (WILS.).

Blue-headed Vireo.

*Muscicapa solitaria* WILS. Am. Orn. II. 1810, 43, pl. 17, fig. 6.*Vireo solitarius* VIEILL. Nouv. Dict. d'Hist. Nat. XXXVI. 1819, 103.

[B 250, C 127, R 141, C 177.]

GEOG. DIST.—Eastern North America to the Plains, north to Hudson Bay and Fort Simpson. South, in winter, to Guatemala. Breeds from southern New England and the northern part of the Lake States northward.

629<sup>a</sup>. **Vireo solitarius cassinii** (XANTUS).

Cassin's Vireo.

*Vireo cassinii* XANT. Pr. Ac. Nat. Sci. Phila. 1858, 117.*Vireo solitarius* var. *cassini* HENSH. Rep. Orn. Spec. (Wheeler's Exp.), 1874, 105.

[B 251, C —, R 141a, C 178.]

GEOG. DIST.—Western United States; confined to the Pacific slope, from British Columbia southward, during the breeding season.

629 *b*. **Vireo solitarius plumbeus** (COUES).

**Plumbeous Vireo.**

*Vireo plumbeus* COUES, Pr. Ac. Nat. Sci. Phila. 1866, 74.

*Vireo solitarius* var. *plumbeus* ALLEN, Bull. M. C. Z. III. 1872, 176.

[B —, C 127a, R 141b, C 179.]

GEOG. DIST.—Southern Rocky Mountain region from the eastern base of the mountains westward to the desert ranges of the Great Basin, and from southern Wyoming south, in winter, to Oaxaca, Mexico. Accidental in New York.

629 *c*. **Vireo solitarius alticola** BREWST.

**Mountain Solitary Vireo.**

*Vireo solitarius alticola* BREWST. Auk, III. Jan. 1886, 111.

[B 250, *part*, C 127, *part*, R 141, *part*, C 177, *part*.]

GEOG. DIST.—Southern Alleghanies (western North Carolina, etc.), south in winter to Florida.

629 *d*. **Vireo solitarius lucasanus** BREWST.

**St. Lucas Solitary Vireo.**

*Vireo solitarius lucasanus* BREWST. Auk, VIII. April, 1891, 147.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California.

SUBGENUS **VIREO** VIEILLOT.

*Vireo* VIEILL. Ois. Am. Sept. I. 1807, 83. Type, *V. musicus*

VIEILL. = *Muscicapa noveboracensis* GMEL.

630. **Vireo atricapillus** WOODH.

**Black-capped Vireo.**

*Vireo atricapillus* WOODH. Pr. Ac. Nat. Sci. Phila. 1852, 60.

[B 427, C 133, R 142, C 185.]

GEOG. DIST.—Central and western Texas, from the Rio Grande north to southwestern Kansas.

631. **Vireo noveboracensis** (GMEL.).

White-eyed Vireo.

*Muscicapa noveboracensis* GMEL. S. N. I. ii. 1788, 947.*Vireo noveboracensis* BONAP. Journ. Ac. Nat. Sci. Phila. IV. 1824, 176.

[B 248, C 129, R 143, C 181.]

GEOG. DIST.—Eastern United States, west to the Rocky Mountains; north to southern New England and Minnesota; south in winter, from Florida to Guatemala and Honduras. Breeds from Florida and the Gulf States northward. Resident in Bermuda.

631 a. **Vireo noveboracensis maynardi** BREWST.

Key West Vireo.

*Vireo noveboracensis maynardi* BREWST. Auk, IV. April, 1887, 148.

[B —, C —, R —, C —.]

GEOG. DIST.—Southern Florida.

632. **Vireo huttoni** CASS.

Hutton's Vireo.

*Vireo huttoni* CASS. Pr. Ac. Nat. Sci. Phila. 1851, 150, pl. 10, fig. 1.

[B 249, C 130, R 144, C 182.]

GEOG. DIST.—Southern and central California, west of the Sierra Nevada.

632 a. **Vireo huttoni stephensi** BREWST.

Stephens's Vireo.

*Vireo huttoni stephensi* BREWST. Bull. Nutt. Orn. Club, VII. July, 1882, 142.

[B —, C —, R —, C —.]

GEOG. DIST.—Arizona, western Mexico, and Lower California.

632 c. **Vireo huttoni obscurus** ANTHONY.

Anthony's Vireo.

*Vireo huttoni obscurus* ANTHONY, Zoe, I. Dec. 1890, 306.

[B—, C —, R —, C —.]

GEOG. DIST.—Pacific coast, from Oregon to southern British Columbia, south in winter to California.

633. **Vireo bellii** AUD.

Bell's Vireo.

*Vireo bellii* AUD. B. Am. VII. 1844, 333, pl. 485.

[B 246, C 131, R 145, C 183.]

GEOG. DIST.—Upper Mississippi Valley and Great Plains, from Dakota, Minnesota, Illinois, and western Indiana, southwestward to Mexico.

633 a. **Vireo bellii pusillus** (COUES).

Least Vireo.

*Vireo pusillus* COUES, Pr. Ac. Nat. Sci. Phila. 1866, 76.*Vireo bellii pusillus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B—, C 132, R 146, C 184.]

GEOG. DIST.—Arizona and California, south to Cape St. Lucas, and throughout western Mexico.

634. **Vireo vicinior** COUES.

Gray Vireo.

*Vireo vicinior* COUES, Pr. Ac. Nat. Sci. Phila. 1866, 75.

[B—, C 128, R 147, C 180.]

GEOG. DIST.—Southern border of the United States, from western Texas and New Mexico to southern and Lower California and north-western Mexico, north to southern Nevada (Grapevine Mts., Lat. 37°) and the Grand Cañon of the Colorado in Arizona.

FAMILY **CÆREBIDÆ**. HONEY CREEPERS.GENUS **CÆREBA** VIEILLOT.

*Cæreba* VIEILLOT, Ois. Am. Sept. I. 1807, 70. Type, *Certhia flaveola* LINN.

635. *Cœreba bahamensis* (REICH.).

Bahama Honey Creeper.

*Certhiola bahamensis* REICH. Handb. I. 1853, 253.*Cœreba bahamensis* A. O. U. Code and Check-List, Suppl. 1889, 23.

[B 301, C 106, R 159, C 153.]

GEOG. DIST.—Bahamas, and the Keys of the southern coast of Florida.

FAMILY **MNIOTILTIDÆ**. WOOD WARBLERS.GENUS **MNIOTILTA** VIEILLOT.*Mniotilta* VIEILL. Analyse, 1816, 45. Type, *Motacilla varia* LINN.636. *Mniotilta varia* (LINN.).

Black and White Warbler.

*Motacilla varia* LINN. S. N. ed. 12, I. 1766, 333.*Mniotilta varia* VIEILL. Nouv. Dict. d'Hist. Nat. XXI. 1818, 230.

[B 167, C 57, R 74, 74a, C 91, 92.]

GEOG. DIST.—Eastern United States to the Plains, north to Fort Simpson, south, in winter, through Central America and the West Indies to Venezuela and Colombia. Breeds from Virginia and southern Kansas northward, and winters from Florida and the Gulf States southward.

GENUS **PROTONOTARIA** BAIRD.*Protonotaria* BAIRD, B. N. Am. 1858, 239. Type, *Motacilla citrea* BODD.637. *Protonotaria citrea* (BODD.).

Prothonotary Warbler.

*Motacilla citrea* BODD. Tabl. P. E. 1783, 44.*Protonotaria citrea* BAIRD, B. N. Am. 1858, 239.



[B 169, C 59, R 75, C 95.]

GEOG. DIST.—Eastern United States, west to Nebraska and Kansas, north to Virginia, southern Michigan, and Iowa, casually to New England, Ontario, and Minnesota; in winter Cuba and northern South America. Breeds throughout its United States range.

GENUS **HELINAIA** AUDUBON.

*Helinaia* AUD. Synop. 1839, 66. Type, *Sylvia swainsonii* AUD.

638. **Helinaia swainsonii** AUD.

Swainson's Warbler.

*Sylvia swainsonii* AUD. Orn. Biog. II. 1834, 563, pl. 198.

*Helinaia swainsonii* AUD. Synop. 1839, 66.

[B 179, C 61, R 76, C 97.]

GEOG. DIST.—Southeastern United States, north to southern Virginia (Dismal Swamp), southwestern Indiana, southeastern Missouri, and west to Texas; in winter, south to Vera Cruz, Mexico, and Jamaica.

GENUS **HELMITHERUS** RAFINESQUE.

*Helmitherus* RAFIN. Journ. de Phys. LXXXVIII. 1819, 417.

Type, *Motacilla vermivora* GMEL.

639. **Helmitherus vermivorus** (GMEL.).

Worm-eating Warbler.

*Motacilla vermivora* GMEL. S. N. I. ii. 1788, 951.

*Helmitheros vermivora* BONAP. Consp. Av. I. April 20, 1850, 314.

[B 178, C 60, R 77, C 96.]

GEOG. DIST.—Eastern United States, north to southern New York and southern New England, west to eastern Nebraska and Texas; south, in winter, to Cuba and northern South America. Breeds throughout its United States range.

GENUS **HELMINTHOPHILA** RIDGWAY.

*Helminthophila* RIDGW. Bull. Nutt. Orn. Club, VII. Jan. 1882, 53.

Type, *Sylvia ruficapilla* WILS.

640. ***Helminthophila bachmanii*** (AUD.).**Bachman's Warbler.***Sylvia bachmanii* AUD. Orn. Biog. II. 1834, 483, pl. 183.*Helminthophila bachmani* RIDGW. Bull. Nutt. Orn. Club, VII.

Jan. 1882, 53.

[B 182, C 64, R 78, C 103.]

GEOG. DIST.—South Atlantic States (southern Virginia to Florida), and westward to Louisiana; Cuba, in winter.

641. ***Helminthophila pinus*** (LINN.).**Blue-winged Warbler.***Certhia pinus* LINN. S. N. ed. 12, I. 1766, 187.*Helminthophila pinus* RIDGW. Bull. Nutt. Orn. Club, VII. Jan. 1882, 53.

[B 180, C 62, R 79, C 98.]

GEOG. DIST.—Eastern United States, from southern New York, southern New England, and southern Minnesota southward, and west to Nebraska and Texas. In winter, south to Mexico, Guatemala and Nicaragua.

642. ***Helminthophila chrysoptera*** (LINN.).**Golden-winged Warbler.***Motacilla chrysoptera* LINN. S. N. ed. 12, I. 1766, 333.*Helminthophila chrysoptera* RIDGW. Bull. Nutt. Orn. Club, VII.

Jan. 1882, 53.

[B 181, C 63, R 81, C 102.]

GEOG. DIST.—Eastern United States, north to southern New England, southwestern Ontario, and southern Minnesota; breeding from northern New Jersey and northern Indiana northward, and southward along the Alleghanies to South Carolina. Central America and northern South America in winter.

643. ***Helminthophila luciae*** (COOPER).**Lucy's Warbler.***Helminthophaga luciae* COOPER, Pr. Cal. Ac. Sci. July, 1862, 120.*Helminthophila luciae* RIDGW. Bull. Nutt. Orn. Club, VII. Jan.

1882, 54.

[B —, C 65, R 83, C 104.]

GEOG. DIST.— Arizona and extreme southwestern Utah, from the Santa Clara Valley southward to Sonora, Mexico.

644. *Helminthophila virginiae* (BAIRD).

Virginia's Warbler.

*Helminthophaga virginiae* BAIRD, B. N. Am. ed. 1860, Atlas, p. xi. foot-note, pl. 79, fig. 1.

*Helminthophila virginiae* RIDGW. Bull. Nutt. Orn. Club, VII. Jan. 1882, 54.

[B —, C 66, R 84, C 105.]

GEOG. DIST.— Rocky Mountain region of the United States, from Wyoming, Colorado, Utah, and Nevada southward on the tableland of Mexico to Guanajuato.

645. *Helminthophila ruficapilla* (WILS.).

Nashville Warbler.

*Sylvia ruficapilla* WILS. Am. Orn. III. 1811, 120, pl. 27, fig. 3.

*Helminthophila ruficapilla* RIDGW. Bull. Nutt. Orn. Club, VII. Jan. 1882, 54.

[B 183, *part*, C 67, *part*, R 85, *part*, C 106, *part*.]

GEOG. DIST.— Eastern North America to the Plains, north to the Fur Countries, breeding from the northern United States northward. Mexico and Guatemala in winter.

645a. *Helminthophila ruficapilla gutturalis* RIDGW.

Calaveras Warbler.

*Helminthophaga ruficapilla* var. *gutturalis* RIDGW. in Hist. N. Am. B. I. Jan. 1874, 191.

*Helminthophila ruficapilla gutturalis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 183, *part*, C 67, *part*, R 85, *part*, C 106, *part*.]

GEOG. DIST.— Pacific coast of North America, eastward, during migrations, to the Rocky Mountains, northward to Kadiak, Alaska, and southward to Lower California and western Mexico.

646. **Helminthophila celata** (SAY).

Orange-crowned Warbler.

*Sylvia celata* SAY, LONG'S EXP. I. 1823, 169.*Helminthophila celata* RIDGW. Bull. Nutt. Orn. Club, VII. Jan. 1882, 54.[B 184, *part*, C 68, R 86, C 107.]

GEOG. DIST.— Eastern North America, breeding as far northward as the Yukon and Mackenzie River districts, and southward through the Rocky Mountains, and wintering in the south Atlantic and Gulf States and Mexico. Rare east of the Alleghanies, north of Virginia.

646 a. **Helminthophila celata lutescens** (RIDGW.).

Lutescent Warbler.

*Helminthophaga celata* var. *lutescens* RIDGW. Am. Jour. Sci. & Arts, 1872, 457.*Helminthophila celata lutescens* BREWST. Bull. Nutt. Orn. Club, VII. April, 1882, 85.[B 184, *part*, C 68a, R 86a, C 108.]

GEOG. DIST.— Western United States, from the Rocky Mountains to the Pacific.

646 b. **Helminthophila celata sordida** TOWNSEND.

Dusky Warbler.

*Helminthophila celata sordida* TOWNSEND, Pr. U. S. Nat. Mus. XIII. 1890, 139.

[B —, C —, R —, C —.]

GEOG. DIST.— San Clemente, Santa Cruz, and Santa Rosa Islands, California.

647. **Helminthophila peregrina** (WILS.).

Tennessee Warbler.

*Sylvia peregrina* WILS. Am. Orn. III. 1811, 83, pl. 25, fig. 2.*Helminthophila peregrina* RIDGW. Bull. Nutt. Orn. Club, VII. Jan. 1882, 54.

[B 185, C 69, R 87, C 109.]

GEOG. DIST.— Eastern North America, breeding from northern New York and northern New England northward to Hudson Bay Territory; in winter south through eastern Mexico to Costa Rica and Colombia.

GENUS **COMPSOTHLYPIS** CABANIS.

*Compsothlypis* CAB. Mus. Hein. I. 1850, 20. Type, *Parus americanus* LINN.

648. **Compsothlypis americana** (LINN.).

Parula Warbler.

*Parus americanus* LINN. S. N. ed. 10, I. 1758, 190.

*Compsothlypis americana* CAB. Mus. Hein. I. 1850, 20.

[B 168, C 58, R 88, C 93.]

GEOG. DIST.—Eastern United States, west to the Plains, north to Canada, and south in winter to the West Indies, Eastern Mexico and Nicaragua. Breeds locally throughout its United States range.

649. **Compsothlypis nigrilora** (COUES).

Sennett's Warbler.

*Parula nigrilora* COUES, Bull. U. S. Geol. & Geog. Surv. Terr. IV. 1878, 11.

*Compsothlypis nigrilora* STEJN. Auk, I. April, 1884, 170.

[B —, C —, R 89a, C 94.]

GEOG. DIST.—Valley of the Lower Rio Grande in Texas, and southward in Nuevo Leon, Tamaulipas, and southeastern San Luis Potosi.

GENUS **DENDROICA** GRAY.SUBGENUS **PERISSOGLOSSA** BAIRD.

*Perissoglossa* BAIRD, Rev. Am. B. I. April, 1865, 180. Type, *Motacilla tigrina* GMEL.

650. **Dendroica tigrina** (GMEL.).

Cape May Warbler.

*Motacilla tigrina* GMEL. S. N. I. ii. 1788, 985.

*Dendroica tigrina* BAIRD, B. N. Am. 1858, 286.



[B 206, C 85, R 90, C 126.]

GEOG. DIST.— Eastern North America, north to Lake Winnipeg and Hudson Bay Territory, west to the Plains. Breeds from northern New England northward; winters in the West Indies.

SUBGENUS **PEUCEDRAMUS** COUES.

*Peucedramus* COUES, in Zool. Wheeler's Exp. 1876, 201. Type, *Sylvia olivacea* GIRAUD.

651. **Dendroica olivacea** (GIRAUD).

Olive Warbler.

*Sylvia olivacea* GIRAUD, Sixteen Sp. Tex. B. 1841, 29, pl. 7, fig. 2.

*Dendroica olivacea* BAIRD, B. N. Am. 1858, 305.

[B—, C—, R 92, C 110.]

GEOG. DIST.— Highlands of Guatemala and Mexico, north to southern New Mexico and Mt. Graham, Arizona.

SUBGENUS **DENDROICA** GRAY.

*Dendroica* GRAY, List Gen. B. App. 1842, 8. Type, *Motacilla coronata* LINN.

652. **Dendroica æstiva** (GMEL.).

Yellow Warbler.

*Motacilla æstiva* GMEL. S. N. I. ii. 1788, 996.

*Dendroica æstiva* BAIRD, B. N. Am. 1858, 282.

[B 203, C 70, R 93, C 111.]

GEOG. DIST.— North America at large, except southwestern part, south in winter to Central America and northern South America. Breeds nearly throughout its North American range.

652 a. **Dendroica æstiva sonorana** BREWST.

Sonora Yellow Warbler.

*Dendroica æstiva sonorana* BREWST. Auk, V. April, 1888, 137.



[B 203, *part*, C 70, *part*, R 93, *part*, C 111, *part*.]

GEOG. DIST.—Southern Arizona to western Texas and northwestern Mexico.

653. ***Dendroica bryanti castaneiceps*** RIDGW.

Mangrove Warbler.

*Dendroica bryanti castaneiceps* RIDGW. Pr. U. S. Nat. Mus. VIII.  
Sept. 2, 1885, 350.

[B—, C—, R—, C—.]

GEOG. DIST.—Western Mexico, and southern part of Lower California.

654. ***Dendroica cærulescens*** (GMEL.).

Black-throated Blue Warbler.

*Motacilla cærulescens* GMEL. S. N. I. 1788, 960.

*Dendroica cærulescens* BAIRD, Rev. Am. B. I. April, 1865, 186.

[B 193, C 76, R 94, C 117.]

GEOG. DIST.—Eastern North America to the Plains, breeding from northern New England and northern New York northward to Labrador, and in the Alleghanies south to northern Georgia; West Indies and Guatemala in winter. Accidental on the Farallon Islands, California.

655. ***Dendroica coronata*** (LINN.).

Myrtle Warbler.

*Motacilla coronata* LINN. S. N. ed. 12, I. 1766, 333.

*Dendroica coronata* GRAY, List Gen. B. App. 1842, 8.

[B 194, C 78, R 95, C 119.]

GEOG. DIST.—Eastern North America, chiefly, straggling more or less commonly westward to the Pacific; breeds from the northern United States northward, and winters from southern New England and the Ohio valley southward to the West Indies, and through Mexico to Panama.

656. ***Dendroica auduboni*** (TOWNS.).

Audubon's Warbler.

*Sylvia auduboni* TOWNS. Journ. Ac. Nat. Sci. Phila. VII. 1837, 191.

*Dendroica audubonii* BAIRD, B. N. Am. 1858, 273.

[B 195, C 79, R 96, C 120.]

GEOG. DIST.—Western United States, east to the western border of the Plains and north to British Columbia; south in winter to Guatemala. Accidental in Pennsylvania and Massachusetts.

657. ***Dendroica maculosa*** (GMEL.).

Magnolia Warbler.

*Motacilla maculosa* GMEL. S. N. I. ii. 1788, 984.*Dendroica maculosa* BAIRD, B. N. Am. 1858, 284.

[B 204, C 84, R 97, C 125.]

GEOG. DIST.—Eastern North America, west to the base of the Rocky Mountains, and casually to British Columbia; breeding from northern New England, northern New York, and northern Michigan, to Hudson Bay Territory and southward in the Alleghanies to Pennsylvania. In winter, Bahamas, Cuba, and south through eastern Mexico to Panama.

658. ***Dendroica cærulea*** (WILS.).

Cerulean Warbler.

*Sylvia cærulea* WILS. Am. Orn. II. 1810, 141, pl. 17, fig. 5.*Dendroica cærulea* BAIRD, B. N. Am. 1858, 280.

[B 201, C 77, R 98, C 118.]

GEOG. DIST.—Eastern United States and southern Ontario, west to the Plains. Rare or casual east of central New York and the Alleghanies. In winter south to Cuba (rare), southeastern Mexico, Central America, Colombia, Peru, and Bolivia. Breeds from West Virginia, Tennessee, Missouri, and Kansas northward to Minnesota.

659. ***Dendroica pensylvanica*** (LINN.).

Chestnut-sided Warbler.

*Motacilla pensylvanica* LINN. S. N. ed. 12, I. 1766, 333.*Dendroica pensylvanica* BAIRD, B. N. Am. 1858, 279.

[B 200, C 83, R 99, C 124.]

GEOG. DIST.—Eastern United States and southern Ontario, west to Manitoba and the Plains, breeding southward to central Illinois and northern New Jersey, and in the Appalachian highlands probably to northern Georgia. Visits the Bahamas, eastern Mexico, Central America and Panama in winter.

**660. *Dendroica castanea* (WILS.).****Bay-breasted Warbler.***Sylvia castanea* WILS. Am. Orn. II. 1810, 97, pl. 14, fig. 4.*Dendroica castanea* BAIRD, B. N. Am. 1858, 276.

[B 197, C 82, R 100, C 123.]

GEOG. DIST.—Eastern North America, north to Hudson Bay. Breeds from northern New England and northern Michigan northward; in winter south through eastern Mexico (rare) and Guatemala to Colombia.

**661. *Dendroica striata* (FORST.).****Black-poll Warbler.***Muscicapa striata* FORST. Philos. Trans. LXII. 1772, 406, 428.*Dendroica striata* BAIRD, B. N. Am. 1858, 280.

[B 202, C 81, R 101, C 122.]

GEOG. DIST.—Eastern North America, west to the Rocky Mountains, north to Greenland, the Barren Grounds, and Alaska, breeding from northern New England and the Catskills northward. South in winter to northern South America, but not recorded from Mexico or Central America.

**662. *Dendroica blackburniæ* (GMEL.).****Blackburnian Warbler.***Motacilla blackburniæ* GMEL. S. N. I. ii. 1788, 977.*Dendroica blackburniæ* BAIRD, B. N. Am. 1858, 274.

[B 196, C 80, R 102, C 121.]

GEOG. DIST.—Eastern North America, west to eastern Kansas and Manitoba, breeding from the southern Alleghanies, Massachusetts, and Michigan northward to Labrador. In winter, south to the Bahamas, eastern Mexico, Central America, Colombia, Ecuador, and Peru.

**663. *Dendroica dominica* (LINN.).****Yellow-throated Warbler.***Motacilla dominica* LINN. S. N. ed. 12, I. 1766, 334.*Dendroica dominica* BAIRD, Rev. Am. B. I. April, 1865, 209.

[B 209, *part*, C 88, R 103, C 129.]

GEOG. DIST.—Southeastern United States, north to southern Maryland and Virginia, and casually to southern New England; south to the West Indies.

663 *a.* ***Dendroica dominica albilora*** RIDGW.

Sycamore Warbler.

*Dendroica dominica* var. *albilora* BAIRD MSS. RIDGW. Am. Nat. VII. Oct. 1873, 606.

[B 209, *part*, C 88*a*, R 103*a*, C 130.]

GEOG. DIST.—Mississippi Valley, west to the Plains, north to Lake Erie and southern Michigan, and east to western North Carolina; in winter south to southern Mexico, Honduras, Guatemala and Nicaragua.

664. ***Dendroica graciae*** BAIRD.

Grace's Warbler.

*Dendroica graciae* "COUES MSS." BAIRD, Rev. Am. B. I. April, 1865, 210.

[B —, C 87, R 104, C 128.]

GEOG. DIST.—Southern New Mexico and Arizona, and southward into Sonora.

665. ***Dendroica nigrescens*** (TOWNS.).

Black-throated Gray Warbler.

*Sylvia nigrescens* TOWNS. Journ. Ac. Nat. Sci. Phila. VII. 1837, 191.

*Dendroica nigrescens* BAIRD, B. N. Am. 1858, 270.

[B 192, C 75, R 105, C 116.]

GEOG. DIST.—Western United States, north to Colorado, Oregon, and British Columbia west of the Cascades, migrating to southern Mexico in winter.

666. ***Dendroica chrysoparia*** SCL. & SALV.

Golden-cheeked Warbler.

*Dendræca chrysoparia* SCL. & SALV. P. Z. S. 1860, 298.

[B—, C 74, R 106, C 115.]

GEOG. DIST.—Southern Texas, and southward to Guatemala.

667. **Dendroica virens** (Gmel.).**Black-throated Green Warbler.***Motacilla virens* GMEL. S. N. I. ii. 1788, 985.*Dendroica virens* BAIRD, B. N. Am. 1858, 267.

[B 189, C 71, R 107, C 112.]

GEOG. DIST.—Eastern North America to the Plains, north to Hudson Bay Territory, breeding from Connecticut and northern Illinois northward, and south along the Alleghanies to South Carolina. In winter, south to Cuba and Panama. Accidental in Greenland and Europe.

668. **Dendroica townsendi** (TOWNS.).**Townsend's Warbler.***Sylvia townsendi* "NUTT." TOWNS. Journ. Ac. Nat. Sci. Phila. VII. 1837, 191.*Dendroica townsendi* BAIRD, B. N. Am. 1858, 269.

[B 191, C 73, R 108, C 114.]

GEOG. DIST.—Western North America, east to central Colorado, north to Sitka, south in winter to Mexico and Guatemala. Accidental near Philadelphia. Breeds from the southern border of the United States northward.

669. **Dendroica occidentalis** (TOWNS.).**Hermit Warbler.***Sylvia occidentalis* TOWNS. Journ. Ac. Nat. Sci. Phila. VII. 1837, 190.*Dendroica occidentalis* BAIRD, B. N. Am. 1858, 268.

[B 190, C 72, R 109, C 113.]

GEOG. DIST.—Western United States, from the Rocky Mountains to the Pacific coast, and from Washington southward; in winter, Lower California and Mexico to the highlands of Guatemala.



670. **Dendroica kirtlandi** BAIRD.

Kirtland's Warbler.

*Sylvicola kirtlandi* BAIRD, Ann. Lyc. N. Y. V. 1852, 216, pl. 6.*Dendroica kirtlandii* BAIRD, B. N. Am. 1858, 286.

[B 205, C 89, R 110, C 131.]

GEOG. DIST.—Eastern United States (South Carolina, Virginia, Ohio, Indiana, Illinois, Missouri, Michigan, Wisconsin, Minnesota), and the Bahamas in winter.

671. **Dendroica vigorsii** (AUD.).

Pine Warbler.

*Sylvia vigorsii* AUD. Orn. Biog. I. 1832, 153, pl. 30.*Dendroica vigorsii* STEJN. Auk, II. Oct. 1885, 343.

[B 198, C 91, R 111, C 134.]

GEOG. DIST.—Eastern United States, west to the Plains, north to Manitoba, Ontario, and New Brunswick, wintering in the South Atlantic and Gulf States, and the Bahamas.

672. **Dendroica palmarum** (GMEL.).

Palm Warbler.

*Motacilla palmarum* GMEL. S. N. I. ii. 1788, 951.*Dendroica palmarum* BAIRD, B. N. Am. 1858, 288.[B 208, *part*, C 90, *part*, R 113, C 132.]

GEOG. DIST.—Northern interior to Great Slave Lake; in winter South Atlantic and Gulf States, the West Indies and Mexico. Of rare but regular occurrence in the Atlantic States in migrations.

672 a. **Dendroica palmarum hypochrysea** RIDGW.

Yellow Palm Warbler.

*Dendroica palmarum hypochrysea* RIDGW. Bull. Nutt. Orn. Club,  
I. Nov. 1876, 85.

[B 208, *part*, C 90, *part*, R 113a, C 133.]

GEOG. DIST.—Atlantic States, north to Hudson Bay. Breeds from eastern Maine, New Brunswick, and Nova Scotia northward; winters in the South Atlantic and Gulf States.



**673. *Dendroica discolor* (VIEILL.).****Prairie Warbler.***Sylvia discolor* VIEILL. Ois. Am. Sept. I. 1807, 37, pl. 98.*Dendroica discolor* BAIRD, B. N. Am. 1858, 290.

[B 210, C 86, R 114, C 127.]

GEOG. DIST.—Eastern United States to the Plains, breeding from Florida north to Michigan and southern New England. Winters in southern Florida and the West Indies.

**GENUS *SEIURUS* SWAINSON.**

*Seiurus* SWAINS. Phil. Mag. I. May, 1827, 369. Type, *Motacilla aurocapilla* LINN.

**674. *Seiurus aurocapillus* (LINN.).****Oven-bird.***Motacilla aurocapilla* LINN. S. N. ed. 12, I. 1766, 334.*Seiurus aurocapillus* SWAINS. Zool. Journ. III. 1827, 171.

[B 186, C 92, R 115, C 135.]

GEOG. DIST.—Eastern North America, north to Hudson Bay Territory and Alaska, breeding from Kansas, the Ohio Valley, and Virginia northward. In winter, Florida, the West Indies, southern Mexico, and Central America to Panama.

**675. *Seiurus noveboracensis* (GMEL.).****Water-Thrush.***Motacilla noveboracensis* GMEL. S. N. I. ii. 1788, 958.*Seiurus noveboracensis* BONAP. Geog. & Comp. List, 1838, 21.[B 187, *part*, C 93, *part*, R 116, C 136.]

GEOG. DIST.—Eastern United States to Illinois, and northward to Arctic America, breeding from the northern United States northward. South in winter to the West Indies, Central America, and northern South America.

**675 a. *Seiurus noveboracensis notabilis* (RIDGW.).****Grinnell's Water-Thrush.**

*Seiurus naevius notabilis* "GRINNELL," RIDGW. Pr. U. S. Nat. Mus. III. 1880, 12.

*Seiurus noveboracensis notabilis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 187, *part*, C 93, *part*, R 116a, C 137.]

GEOG. DIST.—Western United States, from Indiana and Illinois westward to California, and north into British America. Casual in migrations eastward to the Atlantic coast. Winters from the southern border of the United States southward to Lower California, Mexico, and northern South America.

676. **Seiurus motacilla** (VIEILL.).

Louisiana Water-Thrush.

*Turdus motacilla* VIEILL. Ois. Am. Sept. II. 1807, 9, pl. 65.

*Seiurus motacilla* BONAP. Consp. Av. I. 1850, 306.

[B 188, C 94, R 117, C 138.]

GEOG. DIST.—Eastern United States, north to southern New England and southern Michigan, casually north to Lake George, north-eastern New York, west to the Plains. In winter, West Indies, southern Mexico, and Central America to Panama.

GENUS **GEOTHYLPIS** CABANIS.

SUBGENUS **OPORORNIS** BAIRD.

*Oporornis* BAIRD, B. N. Am. 1858, 246. Type, *Sylvia agilis* WILS.

677. **Geothlypis formosa** (WILS.).

Kentucky Warbler.

*Sylvia formosa* WILS. Am. Orn. III. 1811, 85, pl. 25, fig. 3.

*Geothlypis formosa* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 175, C 96, R 119, C 140.]

GEOG. DIST.—Eastern United States, west to the Plains, breeding from the Gulf States north to southern New England and southern Michigan. In winter, West Indies, eastern Mexico, and Central America to Panama.

678. **Geothlypis agilis** (WILS.).

Connecticut Warbler.

*Sylvia agilis* WILS. Am. Orn. V. 1812, 64, pl. 39, fig. 4.

*Geothlypis agilis* GREGG, Pr. Elmira Acad. 1870,—(p. 7 of reprint).

[B 174, C 95, R 118, C 139.]

GEOG. DIST.— Eastern North America, breeding north of the United States (Manitoba, Ontario). Northern South America in winter.

SUBGENUS **GEOTHELYPIS** CABANIS.

*Geothlypis* CAB. Wieg. Archiv, 1847, i. 316, 349. Type, *Turdus trichas* LINN.

679. **Geothlypis philadelphia** (WILS.).

Mourning Warbler.

*Sylvia philadelphia* WILS. Am. Orn. II. 1810, 101, pl. 14, fig. 6.

*Geothlypis philadelphia* BAIRD, B. N. Am. 1858, 243.

[B 172, C 98, R 120, C 142.]

GEOG. DIST.— Eastern North America to the Plains, breeding from the mountainous portions of Pennsylvania, New England, New York, and northern Michigan northward. Central America and northern South America in winter. Accidental in Greenland.

680. **Geothlypis macgillivrayi** (AUD.).

Macgillivray's Warbler.

*Sylvia macgillivrayi* AUD. Orn. Biog. V. 1839, 75, pl. 399, figs. 4, 5.

*Geothlypis macgillivrayi* BAIRD, B. N. Am. 1858, 244.

[B 173, C 99, R 121, C 143.]

GEOG. DIST.— Western United States, from the eastern foothills of the Rocky Mountains to the Pacific coast, north into British Columbia. Lower California, Mexico, and Central America to Colombia in winter.

681. **Geothlypis trichas** (LINN.).

Maryland Yellow-throat.

*Turdus trichas* LINN. S. N. ed. 12, I. 1766, 293.

*Geothlypis trichas* CAB. Mus. Hein. I. 1850, 16.

[B 170, *part*, C 97, *part*, R 122, *part*, C 141, *part*.]

GEOG. DIST.— Eastern United States, north to Ontario, Nova Scotia, and southern Labrador, breeding from Georgia northward. In winter, South Atlantic and Gulf States, the West Indies, eastern Mexico, and Central America; casually northward to Massachusetts.

681 *a.* ***Geothlypis trichas occidentalis*** BREWST.

Western Yellow-throat.

*Geothlypis trichas occidentalis* BREWST. Bull. Nutt. Orn. Club, VIII. July, 1883, 159.[B 170, *part*, C 97, *part*, R 122, *part*, C 141, *part*.]

GEOG. DIST.—Western United States, from the Mississippi Valley west to the Pacific coast, and north to British Columbia; south, in winter, to Central America.

681 *b.* ***Geothlypis trichas ignota*** CHAPM.

Florida Yellow-throat.

*Geothlypis trichas ignota* CHAPM. Auk, VII. Jan. 1890, 11.[B 170, *part*, C 97, *part*, R 122, *part*, C 141, *part*.]

GEOG. DIST.—Florida, north to southern Georgia.

682. ***Geothlypis beldingi*** RIDGW.

Belding's Yellow-throat.

*Geothlypis beldingi* RIDGW. Pr. U. S. Nat. Mus. V. 1882, 344.

[B —, C —, R —, C —.]

GEOG. DIST.—Southern portion of Lower California, north to San Ignacio (Lat. 27°).

682.1. ***Geothlypis poliocephala ralphi*** RIDGW.

Rio Grande Yellow-throat.

*Geothlypis poliocephala ralphi* RIDGW. Proc. U. S. Nat. Mus. 1893, 692.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower Rio Grande Valley.

GENUS **ICTERIA** VIEILLOT.*Icteria* VIEILL. Ois. Am. Sept. I. 1807, pp. iii. 85. Type, *Muscicapa viridis* GMEL. = *Turdus virens* LINN.

683. **Icteria virens** (LINN.).

Yellow-breasted Chat.

*Turdus virens* LINN. S. N. ed. 10, I. 1758, 171.*Icteria virens* BAIRD, Rev. Am. B. I. April, 1865, 228.

[B 176, C 100, R 123, C 144.]

GEOG. DIST.— Eastern United States to the Plains, breeding north to Ontario and southern New England; south, in winter, to eastern Mexico, Guatemala, Nicaragua, and Costa Rica.

683 a. **Icteria virens longicauda** (LAWR.).

Long-tailed Chat.

*Icteria longicauda* LAWR. Ann. Lyc. N. Y. VI. 1853, 4.*Icteria virens* var. *longicauda* COUES, Key, 1872, 108.

[B 177, C 100a, R 123a, C 145.]

GEOG. DIST.— Western United States, from the Plains to the Pacific, north to southern Montana, Washington, and the dry interior of British Columbia, south into Lower California and Mexico.

GENUS **SYLVANIA** NUTTALL.

*Sylvania* NUTT. Man. Land Birds, I. 1832, 290. Type, by elimination, *Muscicapa selbii* AUD. = *Motacilla mitrata* GMEL.

684. **Sylvania mitrata** (GMEL.).

Hooded Warbler.

*Motacilla mitrata* GMEL. S. N. I. ii. 1788, 977.*Sylvania mitrata* NUTT. Man. Land B. ed. 1840, 333.

[B 211, C 101, R 124, C 146.]

GEOG. DIST.— Eastern United States, west to the Plains, north and east to southern Michigan, southern Ontario, western and south-eastern New York, and southern New England. Breeds from the Gulf of Mexico northward. In winter, West Indies, eastern Mexico, and Central America to Panama.

685. **Sylvania pusilla** (WILS.).

Wilson's Warbler.

*Muscicapa pusilla* WILS. Am. Orn. III. 1811, 103, pl. 26, fig. 4.*Sylvania pusilla* NUTT. Man. Land B. ed. 1840, 335.



[B 213, *part*, C 102, R 125, C 147.]

GEOG. DIST.—Eastern North America, west to and including the Rocky Mountains, north to Labrador, Hudson Bay Territory, and Alaska. Breeds chiefly north of the United States, migrating south to eastern Mexico and Central America.

685 *a*. ***Sylvania pusilla pileolata*** (PALL.).

Pileolated Warbler.

*Motacilla pileolata* PALL. Zoog. Rosso-As. I. 1826, 497.

*Sylvania pusilla pileolata* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 213, *part*, C 102*a*, R 125*a*, C 148.]

GEOG. DIST.—Western North America, from the Great Basin to the Pacific, north to Alaska (Kadiak), and south, in winter, to Costa Rica.

686. ***Sylvania canadensis*** (LINN.).

Canadian Warbler.

*Muscicapa canadensis* LINN. S. N. ed. 12, I. 1766, 327.

*Sylvania canadensis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B 214, 215, C 103, R 127, C 149.]

GEOG. DIST.—Eastern North America, west to the Plains, and north to Newfoundland, southern Labrador, and Lake Winnipeg, south, in winter, to Central America and northern South America. Breeds from the higher parts of the Alleghanies, and the more elevated parts of southern New York and southern New England, northward.

GENUS **SETOPHAGA** SWAINSON.

*Setophaga* SWAINS. Phil. Mag. I. May, 1827, 368. Type, *Motacilla ruticilla* LINN.

687. ***Setophaga ruticilla*** (LINN.).

American Redstart.

*Motacilla ruticilla* LINN. S. N. ed. 10, I. 1758, 186.

*Setophaga ruticilla* SWAINS. Phil. Mag. I. May, 1827, 368.



[B 217, C 104, R 128, C 152.]

GEOG. DIST.—North America, north to Fort Simpson, west regularly to the Great Basin, casually to California and Lower California, breeding from the middle portion of the United States northward. In winter, the West Indies, southern Mexico, Central America and northern South America.

688. **Setophaga picta** SWAINS.

Painted Redstart.

*Setophaga picta* SWAINS. Zool. Illustr. 2d ser. I. 1829, pl. 3.

[B 218, C 105, R 129, C 151.]

GEOG. DIST.—Mountains of Mexico, north to southern Arizona.

[689.] **Setophaga miniata** SWAINS.

Red-bellied Redstart.

*Setophaga miniata* SWAINS. Phil. Mag. I. 1827, 368.

[B 219, C —, R 130, C —.]

GEOG. DIST.—Highlands of Mexico. Texas (GIRAUD).

GENUS **CARDELLINA** DU BUS.*Cardellina* DU BUS, Esq. Orn. 1850, pl. 25. Type, *C. amicta*DU BUS = *Muscicapa rubrifrons* GIRAUD.690. **Cardellina rubrifrons** (GIRAUD).

Red-faced Warbler.

*Muscicapa rubrifrons* GIRAUD, Sixteen Sp. Texas B. 1841, pl. 7, fig. 1.*Cardellina rubrifrons* SCL. P. Z. S. 1855, 66.

[B —, C —, R 131, C 150.]

GEOG. DIST.—Southern Arizona and southwestern New Mexico, through Mexico, to Guatemala. Texas (GIRAUD).

GENUS **ERGATICUS** BAIRD.*Ergaticus* BAIRD, Rev. Am. B. I. April, 1865, 237. Type, *Setophaga rubra* SWAINS.

[691.] **Ergaticus ruber** (SWAINS.).

Red Warbler.

*Setophaga rubra* SWAINS. Phil. Mag. I. 1827, 368.*Ergaticus ruber* SCL. & SALV. Nom. Neotr. 1873, 11.

[B 216, C —, R 132, C —.]

GEOG. DIST.— Highlands of Mexico. Texas (GIRAUD).

GENUS **BASILEUTERUS** CABANIS.*Basileuterus* CAB. in SCHOMB. Guiana, III. 1848, 666. Type,*Sylvia vermivora* VIEILL. = *Setophaga auricapilla* SWAINS.[692.] **Basileuterus culicivorus** (LICHT.).

Brasher's Warbler.

*Sylvia culicivora* (LICHT.) Preis-Verzeich. 1830, no. 78.*Basileuterus culicivorus* BONAP. Consp. Av. I. 1850, 313.

[B —, C —, R 133, C —.]

GEOG. DIST.— Central America, from Panama north to eastern Mexico. Texas (GIRAUD.)

[693.] **Basileuterus belli** (GIRAUD).

Bell's Warbler.

*Muscicapa belli* GIRAUD, Sixteen Sp. Texas B. 1841, pl. 4, fig. 1.*Basileuterus belli* SCL. P. Z. S. 1855, 65.

[B —, C —, R 134, C —.]

GEOG. DIST.— Guatemala and Mexico, north to the temperate regions of Vera Cruz. Texas (GIRAUD.)

FAMILY **MOTACILLIDÆ**. WAGTAILS.GENUS **MOTACILLA** LINNÆUS.*Motacilla* LINN. S. N. ed. 10, I. 1758, 184. Type, by elimination, *M. alba* LINN.

[694.] **Motacilla alba** LINN.

White Wagtail.

*Motacilla alba* LINN. S. N. ed. 10, I. 1758, 185.

[B —, C —, R 69, C 86.]

GEOG. DIST.—Northern Europe and northern Asia, south, in winter, to North Africa and India. Accidental in Greenland.

[695.] **Motacilla ocularis** SWINH.

Swinhoe's Wagtail.

*Motacilla ocularis* SWINH. Ibis, Jan. 1860, 55.

[B —, C —, R —, C —.]

GEOG. DIST.—Eastern Asia. Accidental in Lower California. Aleutian Islands?

GENUS **BUDYTES** CUVIER.*Budytes* CUV. Règne An. I. 1817, 371. Type, *Motacilla flava* LINN.696. **Budytes flavus leucostriatus** (HOM.).

Siberian Yellow Wagtail.

*Budytes leucostriatus* HOMEYER, J. f. O. 1878, 128.*Budytes flavus leucostriatus* STEJN. Orn. Expl. Kamtsch. 1885, 280.

[B —, C 54, R 70, C 87.]

GEOG. DIST.—Alaska and northern Siberia to China, wintering in the Moluccas.

GENUS **ANTHUS** BECHSTEIN.SUBGENUS **ANTHUS**.*Anthus* BECHST. Gem. Naturg. Deutschl. III. 1807, 704. Type, by elimination, *A. aquaticus* = *Alauda spinoletta* LINN.697. **Anthus pensilvanicus** (LATH.).

American Pipit.

*Alauda pensilvanica* LATH. Synop. Suppl. I. 1787, 287.*Anthus pensilvanicus* THIENEM. Rhea, II. 1849, 171.

[B 165, C 55, R 71, C 89.]

GEOG. DIST.—North America at large, breeding in the higher parts of the Rocky Mountains and subarctic districts, and wintering in the Gulf States, Mexico, and Central America. Accidental in Europe.

[698.] **Anthus pratensis** (LINN.).

Meadow Pipit.

*Alauda pratensis* LINN. S. N. ed. 10, I. 1758, 166.

*Anthus pratensis* BECHST. Gem. Naturg. Deutschl. III. 1807, 732.

[B —, C 55 *bis*, R 72, C 88.]

GEOG. DIST.—Europe, straggling to Greenland (and Alaska?).

[699.] **Anthus cervinus** (PALLAS).

Red-throated Pipit.

*Motacilla cervina* PALLAS, Zoog. Rosso-As. I. 1826, 511.

*Anthus cervinus* KEYS. & BLAS. Wirb. Eur. I. 1840, p. xlviii.

[B —, C —, R —, C —.]

GEOG. DIST.—Northern parts of the Old World. Accidental in Lower California. St. Michael and Aleutian Islands, Alaska?

SUBGENUS **NEOCORYS** SCLATER.

*Neocorys* SCL. P. Z. S. 1857, 5. Type, *Alauda spragueii* AUD.

700. **Anthus spragueii** (AUD.).

Sprague's Pipit.

*Alauda spragueii* AUD. B. Am. VII. 1843, 335, pl. 486.

*Anthus spraguei* BAIRD, Rev. Am. B. I. Oct. 1864, 155.

[B 166, C 56, R 73, C 90.]

GEOG. DIST.—Interior plains of North America, breeding from the plains of the Yellowstone northward to the Saskatchewan district, and from the Red River westward (probably to the Rocky Mountains). South in winter on the tablelands of Mexico to Puebla. Accidental in South Carolina.

FAMILY **CINCLIDÆ**. DIPPERS.GENUS **CINCLUS** BECHSTEIN.

*Cinclus* BECHST. Orn. Taschenb. Deutschl. 1802, 205. Type,  
*Sturnus cinclus* LINN.

701. **Cinclus mexicanus** SWAINS.

American Dipper.

*Cinclus mexicanus* SWAINS. Phil. Mag. I. 1827, 368.

[B 164, C 19, R 19, C 30.]

GEOG. DIST.—The mountainous parts of central and western North America, from the Yukon Valley and Unalaska to Guatemala; east, in the United States, to the eastern base of the Rocky Mountains. Apparently resident throughout its range.

FAMILY **TROGLODYTIDÆ**. WRENS, THRASHERS, ETC.SUBFAMILY **MIMINÆ**. THRASHERS.GENUS **OROSOPTES** BAIRD.

*Oroscoptes* BAIRD, B. N. Am. 1858, 346. Type, *Orpheus montanus* TOWNS.

702. **Oroscoptes montanus** (TOWNS.).

Sage Thrasher.

*Orpheus montanus* TOWNS. Journ. Ac. Nat. Sci. Phila. VII. 1837,  
193.

*Oroscoptes montanus* BAIRD, B. N. Am. 1858, 347.

[B 255, C 7, R 10, C 14.]

GEOG. DIST.—Western United States, from the western part of the Plains (western South Dakota, western Nebraska, and eastern Colorado), north to Montana, west to the Cascades and Sierra Nevada, south into northern Mexico, Lower California, and casually to Guadalupe Island.

GENUS **MIMUS** BOIE.

*Mimus* BOIE, Isis, Oct. 1826, 972. Type, *Turdus polyglottos* LINN.

703. **Mimus polyglottos** (LINN.).

Mockingbird.

*Turdus polyglottos* LINN. S. N. ed. 10, I. 1758, 169.

*Mimus polyglottus* BONAP. Geog. & Comp. List, 1838, 17.

[B 253, 253a, C 8, R 11, C 15.]

GEOG. DIST.—United States, south into Mexico. Rare and of irregular distribution from Maryland northward to Massachusetts, and north of southern Ohio, Colorado, and southern California. Bahamas. Casual on Guadalupe Island, Lower California.

GENUS **GALEOSOPTES** CABANIS.

*Galeoscoptes* CAB. Mus. Hein. I. 1850, 82. Type, *Muscicapa carolinensis* LINN.

704. **Galeoscoptes carolinensis** (LINN.).

Catbird.

*Muscicapa carolinensis* LINN. S. N. ed. 12, I. 1766, 328.

*Galeoscoptes carolinensis* CAB. Mus. Hein. I. 1850, 82.

[B 254, C 9, R 12, C 16.]

GEOG. DIST.—Eastern United States and British Provinces, west to and including the Rocky Mountains; occasional on the Pacific coast, from British Columbia south to central California. Breeds from the Gulf States northward to the Saskatchewan. Winters in the Southern States, Cuba, and Middle America to Panama. Bermuda, resident. Accidental in Europe.

GENUS **HARPORHYNCHUS** CABANIS.SUBGENUS **METHRIOPTERUS** REICHENBACH.

*Methriopterus* REICH. Syst. Nat. 1850, pl. iv. Type, *Turdus rufus* LINN.



705. **Harporhynchus rufus** (LINN.).

Brown Thrasher.

*Turdus rufus* LINN. S. N. ed. 10, I. 1758, 169.*Harporhynchus rufus* CAB. Mus. Hein. I. 1850, 82.

[B 261, 261a, C 10, R 13, C 17.]

GEOG. DIST.—Eastern United States, west to the Rocky Mountains, north to southern Maine, Ontario, and Manitoba. Breeds from the Gulf States, including eastern Texas, northward. Accidental in Europe.

706. **Harporhynchus longirostris sennetti** (RIDGW.).

Sennett's Thrasher.

*Harporhynchus longirostris sennetti* RIDGW. Proc. U. S. Nat. Mus. X. Aug. 6, 1888, 506.

[B 260, C 10a, R 13a, C 18.]

GEOG. DIST.—Southeastern Texas, from Corpus Christi and Laredo southward to central Nuevo Leon and Tamaulipas.

707. **Harporhynchus curvirostris** (SWAINS.).

Curve-billed Thrasher.

*Orpheus curvirostris* SWAINS. Phil. Mag. I. 1827, 369.*Harporhynchus curvirostris* CAB. Mus. Hein. I. 1850, 81.

[B 259, 259a, C—, R 15, C 19.]

GEOG. DIST.—Tablelands of Mexico, from Oaxaca and Puebla northward to southern New Mexico and southeastward in Texas to the mouth of the Rio Grande.

707 a. **Harporhynchus curvirostris palmeri** COUES.

Palmer's Thrasher.

*Harporhynchus curvirostris* var. *palmeri* "RIDGW." COUES, Key, 1872, 351.

[B —, C 11, R 15a, C 20.]

GEOG. DIST.—Southern Arizona, from about fifty miles northwest of Phoenix south to Guaymas, Sonora.

708. **Harporhynchus bendirei** COUES.

Bendire's Thrasher.

*Harporhynchus bendirei* COUES, Am. Nat. VII. 1873, 330.

[B —, C 11 *bis*, R 14*a*, C 21.]

GEOG. DIST.— Arizona, from Phoenix and the Painted Desert south to Guaymas, Sonora. Westward casually to Agua Caliente, California. Accidental (?) at Colorado Springs, Colorado.

709. **Harporhynchus cinereus** XANTUS.

St. Lucas Thrasher.

*Harporhynchus cinereus* XANTUS, Pr. Ac. Nat. Sci. Phila. 1859, 298.

[B —, C 12, R 14, C 22.]

GEOG. DIST.— Lower California, from Cape St. Lucas north to San Quintin (Lat. 30° 30').

709 *a*. **Harporhynchus cinereus mearnsi** ANTHONY.

Mearns's Thrasher.

*Harporhynchus cinereus mearnsi* ANTHONY, Auk, XII. Jan. 1895, 53.

[B —, C —, R —, C —.]

GEOG. DIST.— Northern Lower California.

SUBGENUS **HARPORHYNCHUS** CABANIS.

*Harporhynchus* CAB. Wieg. Archiv, 1848, i. 98. Type, *Harpes rediviva* GAMB.

710. **Harporhynchus redivivus** (GAMB.).

Californian Thrasher.

*Harpes rediviva* GAMB. Pr. Ac. Nat. Sci. Phila. 1845, 264.

*Harporhynchus redivivus* CAB. Wieg. Archiv, 1848, i. 98.

[B 256, C 13, R 16, C 23.]

GEOG. DIST.— Coast region of California, from the heads of the Sacramento and Russian River valleys south to El Rosario, Lower California (Lat. 30°).

711. **Harporhynchus lecontei** (LAWR.).

Leconte's Thrasher.

*Toxostoma lecontei* LAWR. Ann. Lyc. N. Y. V. 1852, 121.

*Harporhynchus lecontei* BONAP. Notes Coll. Delattre, 1854, 39.

[B 257, C 13a, R 16a, C 24.]

GEOG. DIST.—Desert region of southern California, Nevada, and extreme southwestern Utah, from Benton, Cal. (Lat. 38°), southeastward through Arizona to Sonora (Lat. 30°). Local in the southern San Joaquin Valley.

712. **Harporhynchus crissalis** (HENRY).

Crissal Thrasher.

*Toxostoma crissalis* HENRY, Pr. Ac. Nat. Sci. Phila. 1858, 117.

*Harporhynchus crissalis* BAIRD, B. N. Am. 1858, 350.

[B 258, C 14, R 17, C 25.]

GEOG. DIST.—Southwestern United States, from western Texas to the Colorado Desert, California, and northern Lower California; north to the Charleston Mountains, Nevada, and St. George, Utah.

SUBFAMILY TROGLODYTINÆ. WRENS.

GENUS **HELEODYTES** CABANIS.

*Heleodytes* CABANIS, Mus. Hein. I. 1850, 80. Type, *Furnarius griseus* SWAIN.

713. **Heleodytes brunneicapillus** (LAFR.).

Cactus Wren.

*Picolaptes brunneicapillus* LAFR. Mag. de Zool. 1835, 61, pl. 47.

*Heleodytes brunneicapillus* FISHER, N. Am. Fauna, No. 7, May 31, 1893, 130.

[B 262, C 43, R 56, C 63.]

GEOG. DIST.—Southern border of the United States, from the Lower Rio Grande in Texas to southern California, north to southwestern Utah, and south to central Mexico.

713 a. **Heleodytes brunneicapillus bryanti** ANTHONY.

Bryant's Cactus Wren.

*Heleodytes brunneicapillus bryanti* ANTHONY, Auk, XI. July, 1894, 212.

[B 262, *part*, C 43, *part*, R 56, *part*, C 63, *part*.]

GEOG. DIST.—Northern Lower California, north into southern California.

713 *b. Heleodytes brunneicapillus affinis* (XANTUS).

St. Lucas Cactus Wren.

*Campylorhynchus affinis* XANTUS, Pr. Ac. Nat. Sci. Phila. 1859, 298.*Heleodytes brunneicapillus affinis* ANTHONY, Auk, XII. July, 1895, 280.

[B —, C 44, R 57, C 64.]

GEOG. DIST.— Southern Lower California.

GENUS **SALPINCTES** CABANIS.*Salpinctes* CAB. Wieg. Archiv, 1847, i. 323. Type, *Troglodytes obsoletus* SAY.715. *Salpinctes obsoletus* (SAY).

Rock Wren.

*Troglodytes obsoletus* SAY, LONG'S Exp. II. 1823, 4.*Salpinctes obsoletus* CAB. Wieg. Archiv, 1847, i. 323.

[B 264, C 45, R 58, C 65.]

GEOG. DIST.— Western United States, from the western border of the Plains to the Pacific, north to Dakota, Montana, and British Columbia; south on the tablelands of Mexico and Guatemala to Salvador. Breeds throughout its range, and is resident from about the southern border of the United States southward.

716. *Salpinctes guadeloupensis* RIDGW.

Guadalupe Rock Wren.

*Salpinctes obsoletus guadeloupensis* RIDGW. Bull. U. S. Geol. & Geog. Serv. Terr. II. No. 2, April, 1876, 185.*Salpinctes guadeloupensis* RIDGW. Bull. Nutt. Orn. Club, II. July, 1877, 60.

[B —, C —, R 58a, C —.]

GEOG. DIST.— Guadalupe Island, Lower California.

GENUS **CATHERPES** BAIRD.*Catherpes* BAIRD, B. N. Am. 1858, 356. Type, *Thryothorus mexicanus* SWAINS.

[717.] **Catherpes mexicanus** (SWAINS.).

White-throated Wren.

*Thryothorus mexicanus* SWAINS. Zool. Ill. 2d ser. I. 1829, pl. 11.*Catherpes mexicanus* BAIRD, B. N. Am. 1858, 356.

[B 263, C —, R 59, C 66.]

GEOG. DIST.— Mexico, from Oaxaca and Orizaba northward on the tablelands. Texas (GIRAUD).

717 a. **Catherpes mexicanus conspersus** RIDGW.

Cañon Wren.

*Catherpes mexicanus* var. *conspersus* RIDGW. Am. Nat. VII. Oct. 1873, 602.[B 263, *part*, C 46, R 59a, C 67.]

GEOG. DIST.— Great Basin and Rocky Mountain region, from the Sierra Nevada and Cascades eastward to southern Idaho, Wyoming, Colorado, and western Texas; south on the tablelands of Mexico to Aguas Calientes. Breeds nearly throughout its range; resident in the southern parts of its United States distribution.

717 b. **Catherpes mexicanus punctulatus** RIDGW.

Dotted Cañon Wren.

*Catherpes mexicanus punctulatus* RIDGW. Proc. U. S. Nat. Mus. V. Sept. 5, 1882, 343.[B 263, *part*, C 46, *part*, R 59, *part*, C 67, *part*.]

GEOG. DIST.— Oregon and California, west of the Cascades and Sierra Nevada, and Lower California. Resident from central California southward.

GENUS **THRYOTHORUS** VIEILLOT.SUBGENUS **THRYOTHORUS**.*Thryothorus* VIEILL. Analyse, 1816, 45. Type, *Troglodytes arundinaceus* VIEILL. = *Sylvia ludoviciana* LATH.

718. **Thryothorus ludovicianus** (LATH.).

Carolina Wren.

*Sylvia ludoviciana* LATH. Ind. Orn. II. 1790, 548.*Thryothorus ludovicianus* BONAP. Geog. & Comp. List, 1838, 11.

[B 265, C 47, R 60, C 68.]

GEOG. DIST.—Eastern United States, north to southern New York, southern Michigan, and southern Nebraska; west to the Plains. Rare or casual in southern New England and southern Ontario. Resident nearly throughout its range.

718 a. **Thryothorus ludovicianus miamensis** RIDGW.

Florida Wren.

*Thryothorus ludovicianus* var. *miamensis* RIDGW. Am. Nat. IX.

Aug. 1875, 469.

[B 265, *part*, C 47, *part*, R 60b, C 69.]

GEOG. DIST.—Southern Florida.

718 b. **Thryothorus ludovicianus lomitensis** SENN.

Lomita Wren.

*Thryothorus ludovicianus lomitensis* SENN. Auk, VII. Jan. 1890, 58.[B 265, *part*, C 47, *part*, R 60, *part*, C 68, *part*.]

GEOG. DIST.—Southeastern Texas.

SUBGENUS **THRYOMANES** SCLATER.*Thryomanes* SCL. Cat. Am. B. 1861, 22. Type, *Troglodytes bewickii* AUD.719. **Thryothorus bewickii** (AUD.).

Bewick's Wren.

*Troglodytes bewickii* AUD. Orn. Biog. I. 1831, 96, pl. 18.*Thryothorus bewickii* BONAP. Geog. & Comp. List, 1838, 11.

[B 267, C 48, R 61, C 71.]

GEOG. DIST.—Eastern United States, west to the eastern border of the Plains and eastern Texas; rare east of the Alleghanies north of Maryland and Delaware; north irregularly in the Mississippi Valley to southern Minnesota. Migratory only along the northern border of its range.



**719 a. *Thryothorus bewickii spilurus* (VIG.).****Vigors's Wren.***Troglodytes spilurus* VIG. Zool. Voy. Bloss. 1839, 18, pl. 4, fig. 1.*Thryothorus bewickii* var. *spilurus* BAIRD, Rev. Am. B. I. 1864, 126.[B —, C 48*b*, R 61*a*, C 73.]

GEOG. DIST.—Pacific coast region of North America, from British Columbia southward to Lower California and western Mexico. Breeds nearly throughout its range; resident from central California southward.

**719 b. *Thryothorus bewickii bairdi* (SALV. & GODM.).****Baird's Wren.***Thryothorus bairdi* SALV. & GODM. Biol. Centr.-Am. Aves, I. April, 1880, 95.*Thryothorus bewickii bairdi* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.[B —, C 48*a*, R 61*b*, C 72.]

GEOG. DIST.—Southern Texas, Arizona, and California east of the Sierra Nevada, north to middle Kansas, Colorado, and southern Utah, south into Mexico. Breeds throughout its range; resident from Arizona and Kansas southward.

**719.1. *Thryothorus leucophrys* ANTHONY.****San Clemente Wren.***Thryothorus leucophrys* ANTHONY, Auk, XII. Jan. 1895, 52.

[B —, C —, R —, C —.]

GEOG. DIST.—San Clemente Island, California.

**720. *Thryothorus brevicauda* (RIDGW.).****Guadalupe Wren.***Thryomanes brevicauda* RIDGW. Bull. U. S. Geol. & Geog. Surv. Terr. II. No. 2, April 1, 1876, 186.*Thryothorus brevicauda* SHARPE, Cat. B. Brit. Mus. VI. 1881, 227.

[B —, C —, R 62, C —.]

GEOG. DIST.—Guadalupe Island, Lower California.

GENUS **TROGLODYTES** VIEILLLOT.SUBGENUS **TROGLODYTES**.

*Troglodytes* VIEILL. Ois. Am. Sept. II. 1807, 52. Type, *T. ædon* VIEILL.

721. **Troglodytes ædon** VIEILL.

House Wren.

*Troglodytes ædon* VIEILL. Ois. Am. Sept. II. 1807, 52, pl. 107.

[B 270, 272, C 49, R 63, C 74.]

GEOG. DIST.— Eastern United States and southern Ontario, west to Indiana and Louisiana. Resident from the middle districts southward.

721 *a*. **Troglodytes ædon parkmanii** (AUD.).

Parkman's Wren.

*Troglodytes parkmanii* AUD. Orn. Biog. V. 1839, 310.

*Troglodytes ædon* var. *parkmanni* COUES, Key, 1872, 87.

[B 271, C 49*a*, R 63*a*, C 75.]

GEOG. DIST.— Pacific coast region of British Columbia, Washington, Oregon, and northern California.

721 *b*. **Troglodytes ædon aztecus** BAIRD.

Western House Wren.

*Troglodytes ædon* var. *aztecus* BAIRD, Rev. Am. B. I. Sept. 1864, 139.

[B 271, *part*, C 49*a*, *part*, R 63*a*, *part*, C 75, *part*.]

GEOG. DIST.— Western United States, except Pacific coast, east to Manitoba and Illinois, south into Mexico.

SUBGENUS **ANORTHURA** RENNIE.

*Anorthura* RENNIE, MONT. Orn. Dict. ed. 2, 1831, 570. Type, *Motacilla troglodytes* LINN.

**722. Troglodytes hiemalis** VIEILL.

Winter Wren.

*Troglodytes hiemalis* VIEILL. Nouv. Dict. d'Hist. Nat. XXXIV.  
1819, 514.

[B 273, C 50, R 65, C 76.]

GEOG. DIST.—Eastern North America generally, breeding from the northern parts of the United States northward, and in the Alleghanies south to North Carolina, and wintering from about its southern breeding limit southward.

**722 a. Troglodytes hiemalis pacificus** BAIRD.

Western Winter Wren.

*Troglodytes hyemalis* var. *pacificus* BAIRD, Rev. Am. B. I. Sept.  
1864, 145.[B 273, *part*, C 50, *part*, R 65a, C 77.]

GEOG. DIST.—Pacific coast, from Sitka to southern California, and eastward to the mountains of Idaho; south, in winter, to Mexico.

**723. Troglodytes alascensis** BAIRD.

Alaskan Wren.

*Troglodytes alascensis* BAIRD, Trans. Chic. Ac. Sci. I. 1869, 315,  
pl. 30, fig. 3.

[B —, C 50a, R 66, C 78.]

GEOG. DIST.—Aleutian and Pribilof Islands, Alaska.

GENUS **CISTOTHORUS** CABANIS.SUBGENUS **CISTOTHORUS**.*Cistothorus* CAB. Mus. Hein. I. 1850, 77. Type, *Troglodytes stellaris* LICHT.**724. Cistothorus stellaris** (LICHT.).

Short-billed Marsh Wren.

*Troglodytes stellaris* LICHT. in NAUM. Vög. Deutschl. III. 1823,  
tab. ad p. 724.*Cistothorus stellaris* CAB. Mus. Hein. I. 1850, 77.

[B 269, C 52, R 68, C 81.]

GEOG. DIST.— Eastern United States, north to southern New Hampshire, southern Ontario, southern Michigan, and southern Manitoba, and west to the Plains. Winters in the South Atlantic and Gulf States.

SUBGENUS **TELMATODYTES** CABANIS.

*Telmatodytes* CAB. Mus. Hein. I. 1850, 78. Type, *Certhia palustris* WILS.

725. **Cistothorus palustris** (WILS.).

Long-billed Marsh Wren.

*Certhia palustris* WILS. Am. Orn. II. 1810, 58, pl. 12, fig. 4.

*Cistothorus* (*Telmatodytes*) *palustris* BAIRD, B. N. Am. 1858, 364.

[B 268, C 51, R 67, C 79.]

GEOG. DIST.— Eastern United States, north to Massachusetts, Ontario, and southern Manitoba, wintering from the Gulf States south to eastern Mexico, and locally as far north as southern New England. Breeds throughout its United States and British American range.

725 a. **Cistothorus palustris paludicola** BAIRD.

Tulé Wren.

*Cistothorus palustris*, var. *paludicola* BAIRD, Rev. Am. B. I. Sept. 1864, 148.

[B 268, *part*, C 51, *part*, R 67a, C 80.]

GEOG. DIST.— Western United States, east to the Rocky Mountains and north to British Columbia; south to southern Mexico. Breeds nearly throughout its range, and winters from Oregon southward.

725 b. **Cistothorus palustris griseus** BREWST.

Worthington's Marsh Wren.

*Cistothorus palustris griseus* BREWST. Auk, X. July, 1893, 216.

[B 268, *part*, C 51, *part*, R 67, *part*, C 79, *part*.]

GEOG. DIST.— Coast region of South Carolina and Georgia.

725.1. **Cistothorus marianæ** SCOTT.

Marian's Marsh Wren.

*Cistothorus marianæ* SCOTT, Auk, V. April, 1888, 188.

[B —, C —, R —, C —.]

GEOG. DIST.— Western Florida.

FAMILY **CERTHIIDÆ**. CREEPERS.GENUS **CERTHIA** LINNÆUS.*Certhia* LINN. S. N. ed. 10, I. 1758, 118. Type, by elimination,  
*C. familiaris* LINN.726. **Certhia familiaris americana** (BONAP.).

Brown Creeper.

*Certhia americana* BONAP. Geog. & Comp. List, 1838, 11.*Certhia familiaris* var. *americana* RIDGW. Bull. Essex Inst. V.  
1873, 180.

[B 275, C 42, R 55, C 62.]

GEOG. DIST.— Eastern North America, breeding from the northern  
and more elevated parts of the United States northward, and casually  
further south, migrating southward in winter.726 a. **Certhia familiaris alticola** MILLER.

Mexican Creeper.

*Certhia familiaris alticola* MILLER, Auk, XII. April, 1895, 186.

[B 276, C —, R 55a, C —.]

GEOG. DIST.— Guatemala, Mexico, and southern Arizona.

726 b. **Certhia familiaris montana** RIDGW.

Rocky Mountain Creeper.

[*Certhia familiaris*] *montana* RIDGW. Proc. U. S. Nat. Mus. V.  
July 8, 1882, 114.[B 275, *part*, C 42, *part*, R 55, *part*, C 62, *part*.]

GEOG. DIST.— Rocky Mountains, from northern Mexico to Alaska.

726 *c.* ***Certhia familiaris occidentalis*** RIDGW.

Californian Creeper.

[*Certhia familiaris*] *occidentalis* RIDGW. Proc. U. S. Nat. Mus. V.  
July 8, 1882, 114.

[B 275, *part*, C 42, *part*, R 55, *part*, C 62, *part*.]

GEOG. DIST.—Pacific coast, from California to southern Alaska.

FAMILY **PARIDÆ**. NUTHATCHES AND TITS.

SUBFAMILY **SITTINÆ**. NUTHATCHES.

GENUS **SITTA** LINNÆUS.

*Sitta* LINN. S. N. ed. 10, I. 1758, 115. Type, *S. europæa* LINN.

727. ***Sitta carolinensis*** LATH.

White-breasted Nuthatch.

*Sitta carolinensis* LATH. Ind. Orn. I. 1790, 262.

[B 277, C 38, R 51, C 57.]

GEOG. DIST.—Eastern United States, from Georgia north to the southern British Provinces, and west to the Rocky Mountains.

727 *a.* ***Sitta carolinensis aculeata*** (CASS.).

Slender-billed Nuthatch.

*Sitta aculeata* CASS. Pr. Ac. Nat. Sci. Phila. Oct. 1856, 254.

*Sitta carolinensis* var. *aculeata* ALLEN, Bull. M. C. Z. III. No. 6,  
July, 1872, 161.

[B 278, C 38*a*, R 51*a*, C 58.]

GEOG. DIST.—Western North America, east to the Plains, and south into Mexico.

727 *b.* ***Sitta carolinensis atkinsi*** SCOTT.

Florida White-breasted Nuthatch.

*Sitta carolinensis atkinsi* SCOTT, Auk, VII. April, 1890, 118.



[B 277, *part*, C 38, *part*, R 51, *part*, C 57, *part*.]

GEOG. DIST.—Florida, and northward along the coast to South Carolina.

728. ***Sitta canadensis* LINN.**

Red-breasted Nuthatch.

*Sitta canadensis* LINN. S. N. ed. 12, I. 1766, 177.

[B 279, C 39, R 52, C 59.]

GEOG. DIST.—North America at large, breeding from northern New England, northern New York, and northern Michigan northward, and southward in the Alleghanies, Rocky Mountains, and Sierra Nevada; in winter south to about the southern border of the United States.

729. ***Sitta pusilla* LATH.**

Brown-headed Nuthatch.

*Sitta pusilla* LATH. Ind. Orn. I. 1790, 263.

[B 280, C 40, R 53, C 60.]

GEOG. DIST.—South Atlantic and Gulf States, north to southern Maryland and (casually?) Ohio, Missouri, etc.

730. ***Sitta pygmæa* VIG.**

Pygmy Nuthatch.

*Sitta pygmæa* VIG. Zool. Beechey's Voy. 1839, 25, pl. 4.

[B 281, C 41, R 54, C 61.]

GEOG. DIST.—Western United States, from New Mexico, Colorado and Montana to southern California, Washington, and eastern British Columbia; southward in Mexico to Mt. Orizaba.

730 a. ***Sitta pygmæa leuconucha* ANTHONY.**

White-naped Nuthatch.

*Sitta pygmæa leuconucha* ANTHONY, Proc. Cal. Ac. Sci. 2d ser. II. Oct. 11, 1889, 77.

[B 281, *part*, C 41, *part*, R 54, *part*, C 61, *part*.]

GEOG. DIST.—San Pedro Mountains, Lower California.

SUBFAMILY **PARINÆ**. TITMICE.GENUS **PARUS** LINNÆUS.SUBGENUS **LOPHOPHANES** KAUP.

*Lophophanes* KAUP, Entw. Gesch. Eur. Thierw. 1829, 92. Type,  
*Parus cristatus* LINN.

731. **Parus bicolor** LINN.

Tufted Titmouse.

*Parus bicolor* LINN. S. N. ed. 12, I. 1766, 340.

[B 285, C 27, R 36, C 40.]

GEOG. DIST.—Eastern United States to the Plains, north to northern New Jersey and southern Iowa; casual in southern New England. Resident throughout its breeding range.

731 a. **Parus bicolor texensis** SENN.

Texan Tufted Titmouse.

*Parus bicolor texensis* SENN. Auk, IV. Jan. 1887, 29.

[B —, C —, R —, C —.]

GEOG. DIST.—Southeastern Texas (Bee and Cameron Counties).

732. **Parus atricristatus** CASS.

Black-crested Titmouse.

*Parus atricristatus* CASS. Pr. Ac. Nat. Sci. Phila. 1850, 103, pl. 2.

[B 286, C 29, R 37, C 42.]

GEOG. DIST.—Southeastern Texas and eastern Mexico.

733. **Parus inornatus** GAMB.

Plain Titmouse.

*Parus inornatus* GAMB. Pr. Ac. Nat. Sci. Phila. Aug. 1845, 265.

[B 287, *part*, C 28, *part*, R 38, *part*, C 41, *part*.]

GEOG. DIST.—Pacific coast of California and western Oregon.

733 *a.* **Parus inornatus griseus** RIDGW.

Gray Titmouse.

*Lophophanes inornatus griseus* RIDGW. Pr. U. S. Nat. Mus. V.  
Sept. 5, 1882, 344.*Parus inornatus griseus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885,  
354.[B 287, *part*, C 28, *part*, R 38, *part*, C 41, *part*.]

GEOG. DIST.—New Mexico and Colorado to Arizona, Nevada and California east of the Sierra Nevada.

733 *b.* **Parus inornatus cineraceus** RIDGW.

Ashy Titmouse.

*Lophophanes inornatus cineraceus* RIDGW. Pr. U. S. Nat. Mus. VI.  
Oct. 5, 1883, 154.*Parus inornatus cineraceus* RIDGW. Pr. U. S. Nat. Mus. VIII.  
1885, 354.

[B —, C —, R —, C —.]

GEOG. DIST.—Lower California.

734. **Parus wollweberi** (BONAP.).

Bridled Titmouse.

*Lophophanes wollweberi* BONAP. Compt. Rend. XXXI. Sept. 1850,  
478.*Parus wollweberi* HENRY, Pr. Ac. Nat. Sci. Phila. 1855, 309.

[B 288, C 30, R 39, B 43.]

GEOG. DIST.—Western Texas, southern New Mexico, southern Arizona, and southward on the tableland of Mexico to Orizaba.

SUBGENUS **PARUS** LINNÆUS.*Parus* LINN. S. N. ed. 10, I. 1758, 189. Type, by elimination,  
*P. major* LINN.735. **Parus atricapillus** LINN.

Chickadee.

*Parus atricapillus* LINN. S. N. ed. 12, I. 1766, 341.

[B 290, C 31, R 41, C 44.]

GEOG. DIST.— Eastern North America, north of the Potomac and Ohio Valleys.

735 a. **Parus atricapillus septentrionalis** (HARRIS).

Long-tailed Chickadee.

*Parus septentrionalis* HARRIS, Pr. Ac. Nat. Sci. Phila. 1845, 300.

*Parus atricapillus* var. *septentrionalis* ALLEN, Bull. M. C. Z. III. 1872, 174.

[B 289, 289a, C 31a, R 41a, C 45.]

GEOG. DIST.— Rocky Mountain Plateau region, east to Manitoba and the Plains.

735 b. **Parus atricapillus occidentalis** (BAIRD).

Oregon Chickadee.

*Parus occidentalis* BAIRD, B. N. Am. 1858, 391.

*Parus atricapillus* var. *occidentalis* COUES, Key, 1872,<sup>4</sup>81.

[B 291, C 31c, R 41b, C 46.]

GEOG. DIST.— Northwest coast region, from northern California to Sitka.

736. **Parus carolinensis** AUD.

Carolina Chickadee.

*Parus carolinensis* AUD. Orn. Biog. II. 1834, 341, pl. 160.

[B 293, C 31b, R 42, C 47.]

GEOG. DIST.— Southeastern States, north to New Jersey and Illinois, west to Missouri.

736 a. **Parus carolinensis agilis** SENN.

Plumbeous Chickadee.

*Parus carolinensis agilis* SENN. Auk, V. Jan. 1888, 46.

[B —, C —, R —, C —.]

GEOG. DIST.— Eastern and central Texas (Bee, Victoria, Cook, and Concho Counties, etc.).

737. **Parus meridionalis** SCL.

Mexican Chickadee.

*Parus meridionalis* SCL. P. Z. S. 1856, 293.

[B 292, C —, R 43, C 879.]

GEOG. DIST.—Mountains of Mexico, from Orizaba north to southern Arizona.

738. **Parus gambeli** RIDGW.

Mountain Chickadee.

*Parus gambeli* RIDGW. in A. O. U. Check-List, 1886, 335.

[B 294, C 32, R 40, C 48.]

GEOG. DIST.—Mountainous parts of the western United States, from the eastern base of the Rocky Mountains to the Sierra Nevada, north to British Columbia, Idaho, etc., and south to northern Lower California.

739. **Parus cinctus obtectus** (CAB.).

Siberian Chickadee.

*Parus (Pœcila) obtectus* CAB. J. f. O. 1871, 237.*Parus cinctus obtectus* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B —, C —, R 44, C 52.]

GEOG. DIST.—Northern Alaska and eastern Siberia.

740. **Parus hudsonicus** FORST.

Hudsonian Chickadee.

*Parus hudsonicus* FORST. Phil. Trans. LXII. 1772, 383, 430.

[B 296, C 33, R 45, C 49.]

GEOG. DIST.—Northern North America, from the more elevated parts of the northern United States (northern New England, northern New York, northern Michigan, etc.) northward.

740 a. **Parus hudsonicus stoneyi** (RIDGW.).

Kowak Chickadee.

*Parus stoneyi* RIDGW. Man. N. Am. B. Aug. 1887, 591.*Parus hudsonicus stoneyi* A. O. U. Check-List, 1st Suppl. 1889, 17.

[B —, C —, R —, C —.]

GEOG. DIST.— Valley of the Kowak River, northwestern Alaska.

740 *b.* ***Parus hudsonicus columbianus*** RHOADS.

Columbian Chickadee.

*Parus hudsonicus columbianus* RHOADS, Auk, X. Jan. 1893, 23.[B 296, *part*, C 33, *part*, R 45, *part*, C 49, *part*.]

GEOG. DIST.— Rocky Mountains, from Liard River south into Montana.

741. ***Parus rufescens*** TOWNS.

Chestnut-backed Chickadee.

*Parus rufescens* TOWNS. Journ. Ac. Nat. Sci. Phila. VII. ii. 1837, 190.[B 295, *part*, C 34, *part*, R 46, C 50.]

GEOG. DIST.— Coast district of Oregon, Washington, British Columbia, and southern Alaska.

741 *a.* ***Parus rufescens neglectus*** RIDGW.

California Chickadee.

*Parus rufescens* *β. neglectus* RIDGW. Pr. U. S. Nat. Mus. I. April 25, 1879, 485.[B 295, *part*, C 34, *part*, R 46*a*, C 51.]

GEOG. DIST.— Coast of California, from Monterey County northward.

SUBFAMILY **CHAMÆINÆ**. WREN-TITS AND BUSH-TITS.GENUS **CHAMÆA** GAMBEL.*Chamæa* GAMB. Pr. Ac. Nat. Sci. Phila. 1847, 154. Type, *Parus fasciatus* GAMB.742. ***Chamæa fasciata*** GAMB.

Wren-Tit.

*Parus fasciatus* GAMB. Pr. Ac. Nat. Sci. Phila. Aug. 1845, 265.*Chamæa fasciata* GAMB. Pr. Ac. Nat. Sci. Phila. 1847, 154.



[B 274, *part*, C 26, *part*, R 35, *part*, C 39, *part*.]

GEOG. DIST.—Coast region of California north at least to Humboldt Bay.

742 *a*. **Chamæa fasciata henshawi** RIDGW.

Pallid Wren-Tit.

*Chamæa fasciata henshawi* RIDGW. Pr. U. S. Nat. Mus. V. June 5, 1882, 13.

[B 274, *part*, C 26, *part*, R 35, *part*, C 39, *part*.]

GEOG. DIST.—Interior of California, including the western slope of the Sierra Nevada, from the head of the Sacramento Valley south to northern Lower California.

GENUS **PSALTRIPARUS** BONAPARTE.

*Psaltriparus* BONAP. Compt. Rend. XXXI. 1850, 478. Type,  
*Parus melanotis* HARTL.

743. **Psaltriparus minimus** (TOWNS.).

Bush-Tit.

*Parus minimus* TOWNS. Journ. Ac. Nat. Sci. Phila. VII. ii. 1837, 190.

*Psaltriparus minimus* BONAP. Compt. Rend. XXXVIII. 1854, 62.

[B 298, *part*, C 35, *part*, R 47, *part*, C 53, *part*.]

GEOG. DIST.—Pacific coast region, from northern California to Washington.

743 *a*. **Psaltriparus minimus californicus** RIDGW.

California Bush-Tit.

*Psaltriparus minimus californicus* RIDGW. Pr. Biol. Soc. Wash. II. April 10, 1884, 89.

[B 298, *part*, C 35, *part*, R 47, *part*, C 53, *part*.]

GEOG. DIST.—California, except the northern coast district.

743 *b. Psaltriparus minimus grindæ* (BELDING).

Grinda's Bush-Tit.

*Psaltriparus grindæ* BELD. Pr. U. S. Nat. Mus. VI. Oct. 5, 1883,  
155.*Psaltriparus minimus grindæ* RIDGW. Pr. U. S. Nat. Mus. VIII.  
1885, 354.

[B —, C —, R —, C —.]

GEOG. DIST.— Lower California.

744. *Psaltriparus plumbeus* BAIRD.

Lead-colored Bush-Tit.

*Psaltria plumbea* BAIRD, Pr. Ac. Nat. Sci. Phila. June, 1854, 118.*Psaltriparus plumbeus* BAIRD, B. N. Am. 1858, 398.

[B 299, C 36, R 48, C 54.]

GEOG. DIST.— New Mexico and Arizona, north to eastern Oregon  
and western Wyoming.744.1. *Psaltriparus santaritæ* RIDGW.

Santa Rita Bush-Tit.

*Psaltriparus santaritæ* RIDGW. Proc. U. S. Nat. Mus. X. Sept. 19,  
1888, 697.

[B —, C —, R —, C —.]

GEOG. DIST.— Santa Rita Mountains, southern Arizona.

745. *Psaltriparus lloydi* SENNETT.

Lloyd's Bush-Tit.

*Psaltriparus lloydi* SENN. Auk, V. Jan. 1888, 43.

[B 297, C —, R 49, C 55.]

GEOG. DIST.— Western Texas (mountains between Pecos River  
and Rio Grande), and west to eastern Sonora.GENUS **AURIPARUS** BAIRD.*Auriparus* BAIRD, Rev. Am. B. I. July, 1864, 85. Type,  
*Ægithalus flaviceps* SUND.

746. **Auriparus flaviceps** (SUND.).

Verdin.

*Ægithalus flaviceps* SUND. Öfv. Vet. Ak. Förh. VII. 1850, 129.*Auriparus flaviceps* BAIRD, Rev. Am. B. I. July, 1864, 85.

[B 300, C 37, R 50, C 56.].

GEOG. DIST.—Southern border of the United States, from the Valley of the Rio Grande to Arizona and southern California, north to southern Nevada and southwestern Utah, and south to Lower California, Zacatecas, and southern Tamaulipas. Resident nearly throughout its range.

FAMILY **SYLVIIDÆ**. WARBLERS, KINGLETS, GNATCATCHERS.SUBFAMILY **SYLVIINÆ**. WARBLERS.GENUS **PHYLLOPSEUSTES** MEYER.

*Phyllopseustes* MEYER, Vög. Lifl. Estl. 1815, 122. Type, *Sylvia sibilatrix* BECHST.

747. **Phyllopseustes borealis** (BLAS.).

Kennicott's Willow Warbler.

*Phyllopseuste borealis* BLASIUS, Naumannia, 1858, 313.*Phyllopseustes borealis* MEVES, J. f. O. 1875, 429.

[B —, C 20, R 34, C 32.]

GEOG. DIST.—Northeastern Asia and Alaska.

SUBFAMILY **REGULINÆ**. KINGLETS.GENUS **REGULUS** CUVIER.

*Regulus* CUV. Leç. d'Anat. Comp. I. 1799–1800, tab. ii. Type, *Motacilla regulus* LINN.

748. **Regulus satrapa** LICHT.

Golden-crowned Kinglet.

*Regulus satrapa* LICHT. Verz. Doubl. 1823, 35.

[B 162, *part*, C 22, *part*, R 33, C 34.]

GEOG. DIST.—North America generally, breeding in the northern and elevated parts of the United States and northward, migrating south in winter to Guatemala.

748 *a*. **Regulus satrapa olivaceus** BAIRD.

Western Golden-crowned Kinglet.

*Regulus satrapa* var. *olivaceus* BAIRD, Rev. Am. B. I. July, 1864, 65 (in text under *R. satrapa*).

[B 162, *part*, C 22, *part*, R 33*a*, C 35.]

GEOG. DIST.—Pacific coast region of North America, from California northward.

749. **Regulus calendula** (LINN.).

Ruby-crowned Kinglet.

*Motacilla calendula* LINN. S. N. ed. 12, I. 1766, 337.

*Regulus calendula* LICHT. Verz. Doubl. 1823, 35.

[B 161, C 21, R 30, C 33.]

GEOG. DIST.—North America, south to Guatemala, north to the Arctic coast, breeding chiefly north of the United States, and in the Rocky Mountains, the Sierra Nevada, and the mountains of Arizona.

750. **Regulus obscurus** RIDGW.

Dusky Kinglet.

*Regulus calendula obscurus* RIDGW. Bull. U. S. Geol. & Geog. Surv. Terr. II. No. 2, April 1, 1876, 184.

*Regulus obscurus* RIDGW. Bull. Nutt. Orn. Club, II. July, 1877, 59.

[B —, C —, R 31, C —.]

GEOG. DIST.—Guadalupe Island, Lower California.

SUBFAMILY POLIOPTILINÆ. GNATCATCHERS.

GENUS **POLIOPTILA** SCLATER.

*Polioptila* SCL. P. Z. S. 1855, 11. Type, *Motacilla cærulea* LINN.

751. ***Polioptila cærulea*** (LINN.).

Blue-gray Gnatcatcher.

*Motacilla cærulea* LINN. S. N. ed. 12, I. 1766, 337.*Polioptila cærulea* SCL. P. Z. S. 1855, 11.

[B 282, C 23, R 27, C 36.]

GEOG. DIST.—Middle and southern portions of the eastern United States, south, in winter, to Guatemala, Cuba, and the Bahamas; rare north toward the Great Lakes, southern New York, and southern New England, accidental north to Massachusetts and Maine. Breeds throughout its United States range, and winters from the South Atlantic and Gulf States southward.

751 a. ***Polioptila cærulea obscura*** RIDGW.

Western Gnatcatcher.

*Polioptila cærulea obscura* RIDGW. Proc. U. S. Nat. Mus. V. March 21, 1883, 535, foot-note.

[B 282, *part*, C 23, *part*, R 27, *part*, C 36, *part*.]

GEOG. DIST.—Arizona, California, Lower California, and western Mexico.

752. ***Polioptila plumbea*** BAIRD.

Plumbeous Gnatcatcher.

*Culicivora plumbea* BAIRD, Pr. Ac. Nat. Sci. Phila. June, 1854, 118.*Polioptila plumbea* BAIRD, B. N. A. 1858, 382.

[B 283, C 25, R 28, C 38.]

GEOG. DIST.—Western Texas to eastern edge of Mohave Desert and eastern coast of Lower California.

753. ***Polioptila californica*** BREWST.

Black-tailed Gnatcatcher.

*Polioptila californica* BREWST. Bull. Nutt. Orn. Club, VI. April, 1881, 103.

[B 284, C 24, R 29, C 37.]

GEOG. DIST.—Southern California and Pacific coast of Lower California.

FAMILY **TURDIDÆ**. THRUSHES, SOLITAIRES, STONECHATS,  
BLUEBIRDS, ETC.

SUBFAMILY **MYADESTINÆ**. SOLITAIRES.

GENUS **MYADESTES** SWAINSON.

*Myadestes* SWAINS. Nat. Libr. XIII. Flycatchers, 1838, 132.  
Type, *M. genibarbis* SWAINS.

754. **Myadestes townsendii** (AUD.).

Townsend's Solitaire.

*Ptiliogonys townsendii* AUD. Orn. Biog. V. 1839, 206, pl. 419,  
fig. 2.

*Myiadestes townsendi* CAB. Wieg. Archiv, 1847, i. 208.

[B 235, C 121, R 25, C 169.]

GEOG. DIST.—Western United States, from the Plains westward to the Pacific coast, north to British Columbia, and south, in winter, to the southern border of Arizona and northern Lower California. Breeds from the mountains of New Mexico, southern Arizona, and central California northward.

SUBFAMILY **TURDINÆ**. THRUSHES.

GENUS **TURDUS** LINNÆUS.

SUBGENUS **HYLOCICHLA** BAIRD.

*Hylocichla* BAIRD, Rev. Am. B. I. June, 1864, 12. Type, *Turdus mustelinus* GMEL.

755. **Turdus mustelinus** GMEL.

Wood Thrush.

*Turdus mustelinus* GMEL. S. N. I. ii. 1788, 817.



[B 148, C 3, R 1, C 6.]

GEOG. DIST.— Eastern United States to the Plains, north to southern Michigan, Ontario, and Massachusetts, south, in winter, to Guatemala and Cuba. Breeds from Virginia, Kentucky, and Kansas northward.

756. **Turdus fuscescens** STEPH.

Wilson's Thrush.

*Turdus fuscescens* STEPH. Gen. Zool. X. i. 1817, 182.

[B 151, C 6, R 2, C 7.]

GEOG. DIST.— Eastern United States to the Plains, north to Manitoba, Ontario, Anticosti, and Newfoundland. Breeds from northern New Jersey and the northern part of the Lake States northward; winters sparingly in Florida, but chiefly south of the United States.

756 a. **Turdus fuscescens salicicola** (RIDGW.).

Willow Thrush.

*Hylocichla fuscescens salicicola* RIDGW. Pr. U. S. Nat. Mus. IV. April 6, 1882, 374.*Turdus fuscescens salicicola* COUES, Key, ed. 2, 1884, 246.

[B —, C —, R —, C —.]

GEOG. DIST.— Rocky Mountain region, north to British Columbia, east to Dakota, occasionally to Illinois, casually to South Carolina; in winter south to southern Brazil.

757. **Turdus aliciae** BAIRD.

Gray-cheeked Thrush.

*Turdus aliciae* BAIRD, B. N. Am. 1858, 217.

[B 154, C 5a, R 3, C 12.]

GEOG. DIST.— Eastern North America, west to the Plains, Alaska, and eastern Siberia, north to the Arctic coast, south, in winter, to Costa Rica. Breeds chiefly north of the United States.

757 a. **Turdus aliciae bicknelli** (RIDGW.).

Bicknell's Thrush.

*Hylocichla aliciae bicknelli* RIDGW. Pr. U. S. Nat. Mus. IV. April 6, 1882, 377.*Turdus aliciae bicknelli* COUES, Key, ed. 2, 1884, 248.

[B 154, *part*, C 5a, *part*, R 3, *part*, C 12, *part*.]

GEOG. DIST.—Mountainous parts of the northeastern States (Catskills, White Mountains, etc.) and Nova Scotia, migrating south in winter.

758. ***Turdus ustulatus* NUTT.**

**Russet-backed Thrush.**

*Turdus ustulatus* NUTT. Man. Orn. Land. B. ed. 2, 1840, 830  
(*cestulatus*, err. typ. p. 400).

[B 152, C 5b, R 4, C 11.]

GEOG. DIST.—Pacific coast region of North America, from Alaska to California, south in winter to northern Lower California, Mexico, and Guatemala.

758 a. ***Turdus ustulatus swainsonii* (CAB.).**

**Olive-backed Thrush.**

*Turdus swainsonii* CAB. Fauna Per. 1845-46, 187.

*Turdus ustulatus* β. *swainsoni* RIDGW. Field & Forest, II. May,  
1877, 195.

[B 153, C 5, R 4a, C 13.]

GEOG. DIST.—Eastern North America, and westward to the Upper Columbia River and East Humboldt Mountains, straggling to the Pacific coast. Southward in winter to Cuba, Guatemala, Nicaragua, Colombia, Ecuador and Peru. Casual in Bermuda. Breeds in the northern Alleghanies, the Catskills, the mountainous parts of southern New England, southern Sierra Nevada, and northward.

759. ***Turdus aonalaschkæ* GMEL.**

**Dwarf Hermit Thrush.**

*Turdus aonalaschkæ* GMEL. S. N. I. ii. 1788, 808.

[B 150, C 4b, R 5, C 8.]

GEOG. DIST.—Pacific Coast region, from Alaska to Lower California and western Mexico, east during migrations, to Nevada and Arizona. Breeds from the southern Sierra Nevada in California northward.

759 *a.* **Turdus aonalaschkæ auduboni** (BAIRD).

Audubon's Hermit Thrush.

*Turdus auduboni* BAIRD, Rev. Am. B. June, 1864, 16.*Turdus aonalaschkæ auduboni* RIDGW. Pr. U. S. Nat. Mus. III.

March 27, 1880, 1.

[B 149a, C 4a, R 5a, C 9.]

GEOG. DIST.—Rocky Mountain region, from near the northern border of the United States south to the highlands of Mexico and Guatemala, west to the mountains of Arizona and southern California.

759 *b.* **Turdus aonalaschkæ pallasii** (CAB.).

Hermit Thrush.

*Turdus pallasii* CAB. Wieg. Archiv. 1847, i. 205.*Turdus aonalaschkæ pallasii* RIDGW. Pr. U. S. Nat. Mus. III.

March 27, 1880, 1.

[B 149, C 4, R 5b, C 10.]

GEOG. DIST.—Eastern North America, breeding from the northern Alleghanies, the mountainous parts of southern New England, southern New York, and northern Michigan, etc., northward, and wintering from the Northern States southward.

SUBGENUS **TURDUS** LINNÆUS.

*Turdus* LINN. S. N. ed. 10, I. 1758, 168. Type, by elimination,  
*T. viscivorus* LINN.

[760.] **Turdus iliacus** LINN.

Red-winged Thrush.

*Turdus iliacus* LINN. S. N. ed. 10, I. 1758, 168.

[B —, C —, R 6, C 4.]

GEOG. DIST.—Northern parts of the Old World; accidental in Greenland.

GENUS **MERULA** LEACH.

*Merula* LEACH, Syst. Cat. Brit. Mam. & B. 1816, 20. Type,  
*Turdus merula* LINN.

761. **Merula migratoria** (LINN.).

American Robin.

*Turdus migratorius* LINN. S. N. ed. 12, I. 1766, 292.*Merula migratoria* SWAINS. Phil. Mag. I. 1827, 368.[B 155, *part*, C 1, *part*, R 7, C 1.]

GEOG. DIST.—Eastern North America to the Rocky Mountains, including eastern Mexico and Alaska. Breeds from Virginia and Kansas northward to the Arctic coast; winters from southern Canada and the Northern States (irregularly) southward. Casual in Bermuda. Accidental in Europe.

761 a. **Merula migratoria propinqua** RIDGW.

Western Robin.

*T[urdus] propinquus* RIDGW. Bull. Nutt. Orn. Club, II. Jan. 1877, 9.

*Merula migratoria propinqua* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880, 166.

[B 155, *part*, C 1, *part*, R 7a, C 2.]

GEOG. DIST.—Western United States, from the eastern base of the Rocky Mountains westward, and south to northern Lower California and the tablelands of Mexico.

762. **Merula confinis** (BAIRD).

St. Lucas Robin.

*Turdus confinis* BAIRD, Rev. Am. B. I. June, 1864, 29.

*Merula confinis* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880, 166.

[B —, C 1a, R 8, C 3.]

GEOG. DIST.—Cape region of Lower California; accidental at Hayward, California.

GENUS **HESPEROCICHLA** BAIRD.

*Hesperocichla* BAIRD, Rev. Am. B. I. June, 1864, 12. Type,  
*Turdus naevius* GMEL.

763. **Hesperocichla nævia** (GMEL.).

Varied Thrush.

*Turdus nævius* GMEL. S. N. I. ii. 1788, 817.*Hesperocichla nævia* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24,  
1880, 166.

[B 156, C 2, R 9, C 5.]

GEOG. DIST.—Pacific coast of North America, from Bering Strait to southern California. Accidental on Guadalupe Island, Lower California, and in New Jersey, Long Island, and Massachusetts.

GENUS **CYANECULA** BREHM.*Cyanecula* BREHM, Isis, 1828, 1280. Type, *Motacilla suecica* LINN.[764.] **Cyanecula suecica** (LINN.).

Red-spotted Bluethroat.

*Motacilla suecica* LINN. S. N. ed. 10, I. 1758, 187.*Cyanecula suecica* BREHM, Isis, 1828, 1280.

[B —, C —, R 20, C 31.]

GEOG. DIST.—Northern parts of the Old World; casual in Alaska.

GENUS **SAXICOLA** BECHSTEIN.*Saxicola* BECHST. Orn. Taschb. 1803, 216. Type, *Motacilla œnanthe* LINN.765. **Saxicola œnanthe** (LINN.).

Wheatear.

*Motacilla œnanthe* LINN. S. N. ed. 10, I. 1758, 186.*Saxicola œnanthe* BECHST. Orn. Taschb. 1803, 217.

[B 157, C 15, R 21, C 26.]

GEOG. DIST.—Europe, North Africa, Asia, Alaska, Greenland, and Labrador, straggling southward to Nova Scotia, Maine, Long Island, and Bermuda. Accidental at New Orleans, La.

GENUS **SIALIA** SWAINSON.

*Sialia* SWAINS. Phil. Mag. I. May, 1827, 369. Type, *Motacilla sialis* LINN.

766. ***Sialia sialis*** (LINN.).

Bluebird.

*Motacilla sialis* LINN. S. N. ed. 10, I. 1758, 187.

*Sialia sialis* HALDEM. TREGO'S Geog. Penn. 1843, 77.

[B 158, C 16, R 22, C 27.]

GEOG. DIST.—Eastern United States to the eastern base of the Rocky Mountains, north to Manitoba, Ontario, and Nova Scotia, south, in winter, from the Middle States to the Gulf States and Cuba. Bermuda, resident.

766 a. ***Sialia sialis azurea*** (SWAINS.).

Azure Bluebird.

*Sialia azurea* BAIRD, Rev. Am. B. July, 1884, 62.

*Sialia sialis* var. *azurea* B. B. & R. Hist. N. Am. B. I. Jan. 1874, 62.

[B —, C —, R —, C —.]

GEOG. DIST.—Southern Arizona and eastern Mexico.

767. ***Sialia mexicana occidentalis*** (TOWNS.).

Western Bluebird.

*Sialia occidentalis* TOWNS. Journ. Acad. Nat. Sci. Phila. VII. 1837, 188.

*Sialia mexicana occidentalis* RIDGW. Auk, XI. April, 1894, 151, 154.

[B 159, *part*, C 17, *part*, R 23, *part*, C 28, *part*.]

GEOG. DIST.—Pacific coast, from British Columbia to southern California, east to western Nevada, and casually, during migrations, to New Mexico.



767 a. ***Sialia mexicana bairdi*** RIDGW.

Chestnut-backed Bluebird.

*Sialia mexicana bairdi* RIDGW. Auk, XI. April, 1894, 151, 157.[B 159, *part*, C 17, *part*, R 23, *part*, C 28, *part*.]

GEOG. DIST.— Rocky Mountain district, south to northern Mexico.

767 b. ***Sialia mexicana anabelæ*** ANTHONY.

San Pedro Bluebird.

*Sialia mexicana anabelæ* ANTHONY, Proc. Cal. Acad. Sci. 2d Ser. II. Oct. 1889, 79.

[B —, C —, R —, C —.]

GEOG. DIST.— San Pedro Martir Mountains, Lower California.

768. ***Sialia arctica*** SWAINS.

Mountain Bluebird.

*Erythaca (Sialia) arctica* SWAINS. in SW. & RICH. Fauna Bor. Am. II. 1831, 209, pl. 39.

[B 160, C 18, R 24, C 29.]

GEOG. DIST.— Western North America (chiefly the interior), from the western parts of the Plains to the Pacific, north to Great Slave Lake, south to northern Mexico.



# HYPOTHETICAL LIST.<sup>1</sup>

---

## FAMILY **PODICIPIDÆ**.

### 1. **Æchmophorus clarkii** (LAWR.).

Clark's Grebe.

*Podiceps clarkii* LAWR. in BAIRD'S B. N. Am. 1858, 895.

*Æchmophorus clarkii* COUES, Pr. Ac. Nat. Sci. Phila. 1862, 229.

[B 705, C 608a, R 730, C 846.]

Probably the female of *Æ. occidentalis* (LAWR.). (Cf. HENSHAW, Bull. Nutt. Orn. Club, VI. 1881, pp. 214-218; B. B. & R., Water B. N. Am. II. p. 423; and especially BRYANT, Auk, II. 1885, pp. 313, 314.)

## FAMILY **ALCIDÆ**.

### 2. **Cepphus motzfeldi** (BENICK.).

Black-winged Guillemot.

*Uria motzfeldi* BENICK. Isis, Aug. 1824, 889.

*Cepphus motzfeldi* STEJN. Pr. U. S. Nat. Mus. VII. Aug. 5, 1884,  
210.

[B—, C—, R—, C—.]

North American, but its specific validity not satisfactorily established. (Cf. STEJN. *l. c.*, and Water B. N. Am. II. 1884, pp. 497, 498.)

<sup>1</sup> Consisting of species which have been recorded as North American, but whose status as North American birds is doubtful, either from lack of positive evidence of their occurrence within the prescribed limits of the present Check-List, or from absence of satisfactory proof of their validity as species.

3. **Cepphus carbo** PALL.

Sooty Guillemot.

*Cepphus carbo* PALL. Zoog. Rosso-As. II. 1826, 350.

[B 728, C 633, R 762, C 873.]

No evidence of its occurrence in North America. (Cf. STEJN. Proc. U. S. Nat. Mus. VII. 1884, pp. 225-227.)

FAMILY **LARIDÆ**.GENUS **CREAGRUS** BON.

*Creagrus* BON. Naumannia, 1854, 211. Type, *Larus furcatus* NEBOUX.

4. **Creagrus furcatus** (NEB.).

Swallow-tailed Gull.

*Larus furcatus* NEB. Voy. 'Venus,' Atlas, pl. 10 (1846).*Creagrus furcatus* BONAP. Naumannia, 1854, 213.

[B 679, C 559, R 678, C 791.]

In all probability erroneously accredited to North America. At least nine examples are now known,—the type, said to be from Monterey, Cal., three from the Galapagos, one from the coast of Peru, and four from Malpelo Island (Lat. 3° 59' N., Long. 81° 34½' W.).

FAMILY **DIOMEDEIDÆ**.4.1. **Diomedea exulans** LINN.

Wandering Albatross.

*Diomedea exulans* LINN. S. N. ed. 10, I. 1758, 132.

[B—, C—, R—, C—.]

GEOG. DIST.—Southern oceans; said to have occurred in Tampa Bay, Florida, and off coast of Washington. (Cf. COVES, Auk, II. Oct. 1885, 387; XII. April, 1895, 178; RIDGWAY, Man. N. Am. B. 1887, 51. The Florida record is open to question, and the others are unsatisfactory.)

FAMILY **PROCELLARIIDÆ.**5. **Puffinus kuhlii** (BOIE).

Cinereous Shearwater.

*Procellaria kuhlii* BOIE, Isis, 1835, 257.*Puffinus kuhlii* BONAP. Consp. II. 1856, 202.

[B 651, C 596, R 708, C 831.]

An Eastern Atlantic species, of which no American specimens are known to exist in collections.

6. **Oceanodroma hornbyi** (GRAY).

Hornby's Petrel.

*Thalassidroma hornbyi* GRAY, P. Z. S. 1853, 62.*Oceanodroma hornbyi* BONAP. Consp. II. 1856, 195.

[B 641, C 592, R 727, C 827.]

A very distinct species, of which only one specimen has been obtained, the alleged locality being the "northwest coast of America."

FAMILY **PHALACROCORACIDÆ.**7. **Phalacrocorax perspicillatus** PALL.

Pallas's Cormorant.

*Phalacrocorax perspicillatus* PALL. Zoog. Rosso-As. II. 1826, 305.

[B 621, C 533, R 648, C 756.]

Believed, on good evidence, to be now extinct, as it unquestionably is in the locality (Bering Island) where originally discovered. Only four specimens are known to exist in collections, — two in the St. Petersburg Museum and one each in the Leyden and British Museums. Even if existing, it has no valid claim to a place in the North American fauna. (Cf. STEJN. Pr. U. S. Nat. Mus. VI. 1883, p. 65.)

FAMILY **ARDEIDÆ.**9. **Ardea wuerdemanni** BAIRD.

Würdemann's Heron.

*Ardea wuerdemanni* BAIRD, B. N. Am. 1858, 669.

[B 488, C 450, R 486, *part*, C 656, *part*.]

Believed to be either the colored phase of *A. occidentalis* AUD., or an abnormal specimen of *A. wardi* RIDGW. (Cf. RIDGW. Bull. U. S. Geol. & Geog. Surv. Terr. IV. No. 1, 1878, pp. 229-236; Bull. Nutt. Orn. Club, VII. 1882, pp. 1-6; Auk, I. 1884, pp. 161-163; Water B. N. Am. I. 1884, pp. 7-13.)

10. ***Ardea (Dichromanassa) pealei*** BONAP.

Peale's Egret.

*Ardea pealei* BONAP. Ann. Lyc. N. Y. II. 1826, 154.

[B 482, C 355, *part*, R 491, *part*, C 661, *part*.]

Supposed to be the white phase of *A. rufescens* GMEL., but possibly entitled to recognition as a local or geographical race.

FAMILY **SCOLOPACIDÆ.**

11. ***Tringa (Actodromas) cooperi*** BAIRD.

Cooper's Sandpiper.

*Tringa cooperi* BAIRD, B. N. Am. 1858, 716.

[B 527, C 422, R 535, C 618.]

Known only from the single specimen from which the species was originally described, taken on Long Island, in May, 1833, and still extant in the National Museum. The status of the species is in doubt.

11.1. ***Numenius arquatus*** (LINN.).

European Curlew.

*Scolopax arquata* LINN. Syst. Nat. I. 1758, 145.

*Numenius arquatus* LATH. Gen. Syn. Suppl. I. 1787, 291.

[B —, C —, R —, C —.]

GEOG. DIST.—Northern Europe, migrating to southern Africa. Recorded as occurring on Long Island, N. Y. (Cf. MARSHALL and DUTCHER, Auk, IX. Oct. 1892, 390-392.) While there is no question as to the proper identification of the specimen, the evidence that it was taken on Long Island is not considered entirely satisfactory.



FAMILY **CATHARTIDÆ.**GENUS **GYPAGUS** VIEILLOT.

*Gypagus* VIEILL. Analyse, 1816, 21. Type, by elimination,  
*Vultur papa* LINN.

12. **Gypagus papa** (LINN.).  
 King Vulture.

*Vultur papa* LINN. S. N. ed. 10, I. 1758, 86.

*Gypagus papa* VIEILL. Nouv. Dict. d'Hist. Nat. XXXVI. 1819,  
 456.

[B —, C —, R —, C —.]

GEOG. DIST.—Tropical continental America. Recorded as occurring on the Rio Verde, Arizona, but its identity not satisfactorily determined. (Cf. COVES, Bull. Nutt. Orn. Club, VI. 1881, p. 248.)

13. **Cathartes burrovianus** CASS.  
 Burroughs's Turkey Vulture.

*Cathartes burrovianus* CASS. Pr. Ac. Nat. Sci. Phila. II. 1845, 212.

[B 4, C —, R —, C —.]

GEOG. DIST.—Mexico and eastern South America. Reported as having been seen near Brownsville, Texas. (Cf. DRESSER, Ibis, 1865, p. 322.)

FAMILY **FALCONIDÆ.**

14. **Buteo cooperi** CASS.  
 Cooper's Henhawk.

*Buteo cooperi* CASS. Pr. Ac. Nat. Sci. Phila. VIII. 1856, 253.

[B 29, C 349, R 437, C 514.]

Probably the light phase of *B. harlani* AUD. (Cf. RIDGW. Auk, I. 1884, pp. 253, 254; *ibid.*, II. 1885, pp. 165, 166.)

FAMILY **PSITTACIDÆ**.GENUS **RHYNCHOPSITTA** BONAPARTE.

*Rhynchopsitta* BONAP. Rev. et Mag. Zool. VI. 1854, 149. Type,  
*Macrocerus pachyrhynchus* SWAINS.

16. **Rhynchopsitta pachyrhyncha** (SWAINS.).

Thick-billed Parrot.

*Macrocerus pachyrhynchus* SWAINS. Phil. Mag. I. 1827, 439.

*Rhynchopsitta pachyrhyncha* BONAP. Rev. et Mag. Zool. VI. 1854,  
149.

[B 64, C —, R 391, C —.]

GEOG. DIST.—Mexico. There is said to be a specimen in “the collection of the Philadelphia Academy of Natural Sciences, labelled Rio Grande, Texas, J. W. Audubon,” but there is doubt as to whether the specimen was really taken within the limits of the United States. (Cf. BAIRD, Birds N. Am. 1858, p. 66, foot-note; RIDGW. Man. N. Am. B. 1887, 269.) Its occurrence in Texas is not improbable but the evidence is unsatisfactory.

FAMILY **CAPRIMULGIDÆ**.16.1. **Chordeiles virginianus sennetti** (COUES).

Sennett's Nighthawk.

*Chordiles popetue sennetti* COUES, Auk, V. Jan. 1888, 37.

*Chordeiles virginianus sennetti* CHAMBERLAIN, Systematic Table  
of Canadian Birds, 1888, Appendix A. p. 14.

[B 115, *part*, C 267*a*, *part*, R 367*a*, *part*, C 400, *part*.]

GEOG. DIST.—“Dakota to Texas, in any treeless country.”

FAMILY **FRINGILLIDÆ**.17. **Acanthis brewsterii** RIDGW.

Brewster's Linnet.

*Ægiothus (flaviostris* var.) *brewsterii* RIDGW. Am. Nat. July,  
1872, 433.

*Acanthis brewsterii* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B —, C 147, R 180, C 211.]

The type-specimen, taken at Waltham, Mass., remains unique. It cannot be identified with any known species, but may be a hybrid between *Acanthis linaria* and *Spinus pinus*. (Cf. BREWST. Bull. Nutt. Orn. Club, VI. 1881, p. 225.)

18. **Spiza townsendii** (AUD.).

Townsend's Bunting.

*Emberiza townsendii* AUD. Orn. Biog. II. 1834, 183.

*Spiza townsendii* RIDGW. Pr. U. S. Nat. Mus. III. Aug. 24, 1880, 182.

[B 379, C 192, R 255, C 288.]

The original specimen, taken May 11, 1833, in Chester County, Pa., by Mr. J. K. Townsend, remains unique. Its peculiarities cannot be accounted for by hybridism, nor probably by individual variation.

FAMILY **VIREONIDÆ**.

GENUS **HYLOPHILUS** TEMMINCK.

*Hylophilus* TEMM. Pl. Col. III. Livr. 29, 1823, text, and pl. 173, fig. 1. Type, *H. thoracicus* TEMM.

19. **Hylophilus decurtatus** (BONAP.).

Short-winged Hylophilus.

*Sylvicola decurtata* BONAP. P. Z. S. 1837, 118.

*Hylophilus decurtatus* BAIRD, Rev. Am. B. I. 1866, 380.

[B—, C—, R—, C—.]

GEOG. DIST.—Mexico and Central America, to Isthmus of Panama. Southern Texas? (*Helinai brevipennis* GIRAUD, Ann. Lyc. N. Y. 1850, 40. "Mexico and Texas.")

FAMILY **MNIOTILTIDÆ**.

20. **Helminthophila lawrencei** (HERRICK).

Lawrence's Warbler.

*Helminthophaga lawrencei* HERRICK, Pr. Ac. Nat. Sci. Phila. 1874, 220, pl. 15.

*Helminthophila lawrencei* RIDGW. Bull. Nutt. Orn. Club. VII. Jan. 1882, 53.

[B —, C —, R 80, C 99.]

About a dozen specimens have been taken, chiefly in New Jersey and southern Connecticut. Supposed to be a hybrid between *H. pinus* and *H. chrysoptera*. (Cf. RIDGW. Ibis, 1876, p. 169; BREWST. Bull. Nutt. Orn. Club, VI. 1881, pp. 218-225; EAMES, Auk, VI. 1889, pp. 305-310.)

21. ***Helminthophila leucobronchialis*** (BREWST.).

**Brewster's Warbler.**

*Helminthophaga leucobronchialis* BREWST. Bull. Nutt. Orn. Club, I. Jan. 1876, 1, plate.

*Helminthophila leucobronchialis* RIDGW. Bull. Nutt. Orn. Club, VII. Jan. 1882, 53.

[B —, C —, R 82, C 100.]

Known from numerous specimens, taken in Southern New England, Lower Hudson Valley, New Jersey, Virginia, Michigan, etc. Supposed to be a hybrid between *H. pinus* and *H. chrysoptera*. (Cf. BREWST. Bull. Nutt. Orn. Club, VI. 1881, pp. 218-225; RIDGW. Auk, II. Oct. 1885, pp. 359-363; EAMES, Auk, VI. 1889, pp. 305-310.)

22. ***Helminthophila cincinnatiensis*** (LANGD.).

**Cincinnati Warbler.**

*Helminthophaga cincinnatiensis* LANGD. Jour. Cinc. Soc. N. H. July, 1880, 119, 120, pl. 4.

*Helminthophila cincinnatiensis* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354.

[B —, C —, R —, C 101.]

One specimen taken near Cincinnati, Ohio. Probably a hybrid between *H. pinus* and *Geothlypis (Oporornis) formosa*. (Cf. RIDGW. Bull. Nutt. Orn. Club, V. 1880, p. 237.)

23. ***Dendroica (Perissoglossa ?) carbonata*** (AUD.).

**Carbonated Warbler.**

*Sylvia carbonata* AUD. Orn. Biog. I. 1831, 308, pl. 60.

*Dendroica carbonata* BAIRD, B. N. Am. 1858, 287.

*Perissoglossa carbonata* B. B. & R. Hist. N. Am. B. I. Jan. 1874, 214.

[B 207, C —, R 91, C —.]

Known only from Audubon's plate and description of two specimens killed near Henderson, Kentucky, in May, 1811.

24. **Dendroica montana** (WILS.).

Blue Mountain Warbler.

*Sylvia montana* WILS. Am. Orn. V. 1812, 113, pl. 44, fig. 2.

*Dendroica montana* BAIRD, B. N. Am. 1858, 278.

[B 199, C —, R 112, C —.]

Known only from the works of Wilson and Audubon. Taken in the Blue Mountains of Virginia. Not as yet satisfactorily identified with any other species.

25. **Sylvania (?) microcephala** RIDGW.

Small-headed Warbler.

*Sylvania microcephala* RIDGW. Pr. U. S. Nat. Mus. VIII. 1885, 354. (= *Muscicapa minuta* WILS. Am. Orn. VI. 1812, 62, pl. 1, fig. 5, nec GMEL., 1788.)

[B 212, C —, R 126, C —.]

Known only from the works of Wilson and Audubon. Claimed to have been taken in New Jersey and Kentucky.

FAMILY **SYLVIIDÆ.**

26. **Regulus cuvierii** AUD.

Cuvier's Kinglet.

*Regulus cuvierii* AUD. Orn. Biog. I. 1832, 288, pl. 55.

[B 163, C —, R 32, C —.]

Known only from Audubon's description and figure of the original specimen, killed in June, 1812, on the banks of the Schuylkill River, in Pennsylvania.





THE  
FOSSIL BIRDS OF NORTH AMERICA.

---

**A. — JURASSIC.**

1. **Laopteryx priscus** MARSH.

*Laopteryx priscus* MARSH, Am. Journ. Sci. XXI. 1881, 341.

Upper Jurassic beds of Wyoming.

**B. — CRETACEOUS.**<sup>1</sup>

2. **Apatornis celer** MARSH.

*Ichthyornis celer* MARSH, Am. Journ. Sci. V. 1873, 74.

*Apatornis celer* MARSH, Am. Journ. Sci. V. 1873, 162.

Middle Cretaceous of western Kansas.

3. **Baptornis advenus** MARSH.

*Baptornis advenus* MARSH, Am. Journ. Sci. XIV. 1877, 86.

Cretaceous of western Kansas, in the same beds with *Odontornithes* and *Pteranodontia*.

4. **Cimolopteryx rarus** MARSH.

*Cimolopteryx rarus* MARSH, Am. Journ. Sci. XLIV. 1892, 175,  
pl. iii, fig. 2.

Laramie beds of Wyoming.

5. **Cimolopteryx retusus** MARSH.

*Cimolopteryx retusus* MARSH, Am. Journ. Sci. XLIV. 1892, 175.

Laramie beds of Wyoming.

<sup>1</sup>The genera alphabetically arranged.

6. **Coniornis altus** MARSH.

*Coniornis altus* MARSH, Am. Journ. Sci. XLV. 1893, 82.

Cretaceous of Montana, near mouth of Judith River.

7. **Graculavus velox** MARSH.

*Graculavus velox* MARSH, Am. Journ. Sci. III. 1872, 363.

Greensand of the middle marl bed, or Upper Cretaceous, near Hornerstown, New Jersey.

8. **Graculavus pumilus** MARSH.

*Graculavus pumilus* MARSH, Am. Journ. Sci. III. 1872, 364.

Greensand of the middle marl bed, or Upper Cretaceous, near Hornerstown, New Jersey.

9. **Hesperornis regalis** MARSH.

*Hesperornis regalis* MARSH, Am. Journ. Sci. III. 1872, 56.

Pteranodon beds of western Kansas.

10. **Hesperornis crassipes** MARSH.

*Lestornis crassipes* MARSH, Am. Journ. Sci. XI. 1876, 509.

*Hesperornis crassipes* MARSH, Odontornithes, 1880, 196, figs.  
40 a-d, pls. vii, xvii.

Yellow chalk of the Pteranodon beds, western Kansas.

11. **Hesperornis gracilis** MARSH.

*Hesperornis gracilis* MARSH, Am. Journ. Sci. XI. 1876, 510.

Yellow chalk of the Pteranodon beds, western Kansas.

12. **Ichthyornis dispar** MARSH.

*Ichthyornis dispar* MARSH, Am. Journ. Sci. IV. 1872, 344.

Pteranodon beds, Middle Cretaceous, northwestern Kansas.

**13. *Ichthyornis agilis* MARSH.**

*Graculavus agilis* MARSH, Am. Journ. Sci. V. 1873, 230.

*Ichthyornis agilis* MARSH, Odontornithes, 1880, 197.

Pteranodon beds, Middle Cretaceous, western Kansas.

**14. *Ichthyornis anceps* MARSH.**

*Graculavus anceps* MARSH, Am. Journ. Sci. III. 1872, 364.

*Ichthyornis anceps* MARSH, Odontornithes, 1880, 198.

Gray shale of the Middle Cretaceous, Smoky Hill River, western Kansas.

**15. *Ichthyornis lentus* MARSH.**

*Graculavus lentus* MARSH, Am. Journ. Sci. XIV. 1877, 253.

*Ichthyornis lentus* MARSH, Odontornithes, 1880, 198.

Middle Cretaceous beds, near Fort McKinney, Texas.

**16. *Ichthyornis tener* MARSH.**

*Ichthyornis tener* MARSH, Odontornithes, 1880, 198, pl. xxx. fig. 8.

Pteranodon beds, Middle Cretaceous, Wallace County, Kansas.

**17. *Ichthyornis validus* MARSH.**

*Ichthyornis validus* MARSH, Odontornithes, 1880, 198, pl. xxx.  
figs. 11-14.

Yellow chalk of the Middle Cretaceous, near Solomon River, northwestern Kansas.

**18. *Ichthyornis victor* MARSH.**

*Ichthyornis victor* MARSH, Am. Journ. Sci. XI. 1876, 511.

Middle Cretaceous of Kansas, in various localities.

**19. *Laornis edwardsianus* MARSH.**

*Laornis edwardsianus* MARSH, Pr. Ac. Nat. Sci. Phila. 1870, 5.

Middle marl bed, Upper Cretaceous, Birmingham, New Jersey.

20. **Palæotringa littoralis** MARSH.

*Palæotringa littoralis* MARSH, Pr. Ac. Nat. Sci. Phila. 1870, 5.

Greensand of the Upper Cretaceous, near Hornerstown, New Jersey.

21. **Palæotringa vagans** MARSH.

*Palæotringa vagans* MARSH, Am. Journ. Sci. III. 1872, 365.

Greensand of the Upper Cretaceous, near Hornerstown, New Jersey.

22. **Palæotringa vetus** MARSH.

*Scolopax* MORTON, Syn. Organic Remains of the Cret. U. S.  
1834, 32.

*Palæotringa vetus* MARSH, Pr. Ac. Nat. Sci. Phila. 1870, 5.

Lower marl bed of the Cretaceous formation, near Arneytown, New Jersey.

23. **Telmatornis priscus** MARSH.

*Telmatornis priscus* MARSH, Pr. Ac. Nat. Sci. Phila. 1870, 5.

Middle marl bed of the Upper Cretaceous, near Hornerstown, New Jersey.

24. **Telmatornis affinis** MARSH.

*Telmatornis affinis* MARSH, Pr. Ac. Nat. Sci. Phila. 1870, 5.

Middle marl beds of the Upper Cretaceous, near Hornerstown, New Jersey.

## C. — TERTIARY.

## SUBCLASS RATITÆ.

25. **Gastornis giganteus** (COPE).

*Diatryma gigantea* COPE, Pr. Ac. Nat. Sci. Phila. 1876, 11.

*Gastornis giganteus* COUES, Key N. A. Birds, 2d ed. 1884, 825.

Wahsatch Epoch, Eocene of New Mexico.

26. **Barornis regens** MARSH.

*Barornis regens* MARSH, Am. Journ. Sci. XLVIII. 1894, 344.

Eocene of New Jersey, near Squantum, N. J.

## SUBCLASS CARINATÆ.

## ORDER PYGOPODES.

27. **Uria antiqua** (MARSH).

*Catarractes antiqua* MARSH, Am. Journ. Sci. XLIX. 1870, 213.

*Uria antiqua* COUES, in A. O. U. Check-List, ed. 1, 1886, 363.

Miocene of North Carolina.

28. **Uria affinis** (MARSH).

*Catarractes affinis* MARSH, Am. Journ. Sci. IV. 1872, 259.

*Uria affinis* COUES, in A. O. U. Check-List, ed. 1, 1886, 363.

Post-pliocene of Maine.

## ORDER LONGIPENNES.

29. **Larus robustus** SHUFELDT.

*Larus robustus* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI. 1892,  
398, pl. xv, fig. 1, 2.

Pliocene of Oregon.

30. **Larus oregonus** SHUFELDT.

*Larus oregonus* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI. 1892,  
398.

Pliocene of Oregon.

ORDER TUBINARES.

31. **Puffinus conradii** MARSH.

*Puffinus conradii* MARSH, Am. Journ. Sci. XLIX. 1870, 212.

Miocene of Maryland.

ORDER STEGANOPODES.

32. **Sula loxostyla** COPE.

*Sula loxostyla* COPE, Tr. Amer. Philos. Soc. XIV. 1870, 236.

Miocene of North Carolina.

33. **Phalacrocorax idahensis** (MARSH).

*Graculus idahensis* MARSH, Am. Journ. Sci. XLIX. 1870, 216.

*Phalacrocorax idahensis* COUES, Key N. A. Birds, 2d ed. 1884,  
824.

Pliocene of Idaho.

34. **Phalacrocorax macropus** (COPE).

*Graculus macropus* COPE, Bull. U. S. Geol. Surv. Terr. IV. No.  
2, 1878, 386.

*Phalacrocorax macropus* COUES, Key N. A. Birds, 2d ed. 1884,  
824.

Pliocene of Oregon.



## ORDER ANSERES.

35. **Anser condoni** SHUFELDT.

*Anser condoni* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI. 1892,  
406.

Pliocene of Oregon.

36. **Branta hypsibatus** (COPE).

*Anser hypsibatus* COPE, Bull. U. S. Geol. Surv. Terr. IV. No. 2,  
1878, 387.

*Branta hypsibates* COUES, in A. O. U. Check-List, ed. 1, 1886,  
364.

37. **Branta propinqua** SHUFELDT.

*Branta propinqua* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI.  
1892, 407, pl. xv, fig. 17.

Pliocene of Oregon.

38. **Cygnus paloregonus** COPE.

*Cygnus paloregonus* COPE, Bull. U. S. Geol. Surv. Terr. IV. No.  
2, 1878, 388.

Pliocene of Oregon.

## ORDER ODONTOGLOSSÆ.

39. **Phœnicopterus copei** SHUFELDT.

*Phœnicopterus copei* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI.  
1892, 410, pl. xv, fig. 41-43, pl. xvii, fig. 28, 29, 38.

Pliocene of Oregon.

## ORDER HERODIONES.

40. **Ardea paloccidentalis** SHUFELDT.

*Ardea paloccidentalis* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI.  
1892, 411, pl. xvii. fig. 31.

Pliocene of Oregon.

## ORDER PALUDICOLÆ.

41. **Grus haydeni** MARSH.

*Grus haydeni* MARSH, Am. Journ. Sci. XLIX. 1870, 214.

Pliocene of Nebraska.

42. **Grus proavus** MARSH.

*Grus proavus* MARSH, Am. Journ. Sci. IV. 1872, 261.

Post-pliocene of New Jersey.

43. **Fulica minor** SHUFELDT.

*Fulica minor* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI. 1892,  
412, pl. xvii. fig. 32.

Pliocene of Oregon.

44. **Aletornis nobilis** MARSH.

*Aletornis nobilis* MARSH, Am. Journ. Sci. IV. 1872, 256.

Eocene of Wyoming.

45. **Aletornis pernix** MARSH.

*Aletornis pernix* MARSH, Am. Journ. Sci. IV. 1872, 256.

Eocene of Wyoming.

46. **Aletornis venustus** MARSH.

*Aletornis venustus* MARSH, Am. Journ. Sci. IV. 1872, 257.

Eocene of Wyoming.

47. **Aletornis gracilis** MARSH.

*Aletornis gracilis* MARSH, Am. Journ. Sci. IV. 1872, 258.

Eocene of Wyoming.

48. **Aletornis bellus** MARSH.

*Aletornis bellus* MARSH, Am. Journ. Sci. IV. 1872, 258.

Eocene of Wyoming.

## ORDER LIMICOLÆ.

49. **Charadrius sheppardianus** COPE.

*Charadrius sheppardianus* COPE, Bull. U. S. Geol. Surv. Terr.  
VI. No. 1, 1881, 83.

(Formation and locality not given.)

## ORDER GALLINÆ.

50. **Palæotetrix gilli** SHUFELDT.

*Palæotetrix gilli* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI. 1892,  
415, pl. xvii. fig. 37.

Pliocene of Oregon.

51. **Pediocætes lucasi** SHUFELDT.

*Pediocætes lucasi* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI. 1892,  
414, pl. xvii. fig. 30.

Pliocene of Oregon.

52. **Pediocætes nanus** SHUFELDT.

*Pediocætes nanus* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI.  
1892, 414, pl. xvii. fig. 36, 37.

Pliocene of Oregon.

53. **Meleagris antiquus** MARSH.

*Meleagris antiquus* MARSH, Am. Journ. Sci. II. 1871, 126.

Miocene of Colorado.

54. **Meleagris altus** MARSH.

*Meleagris altus* MARSH, Pr. Ac. Nat. Sci. Phila. 1870, 11.

Post-pliocene of New Jersey.

55. **Meleagris celer** MARSH.

*Meleagris celer* MARSH, Am. Journ. Sci. 1872, 261.

Post-pliocene of New Jersey.

## ORDER RAPTORES.

56. **Palæoborus umbrosus** COPE.

*Cathartes umbrosus* COPE, Pr. Ac. Nat. Sci. Phila. 1874, 151.

*Palæoborus umbrosus* COUES, Key N. A. Birds, 2d ed. 1884, 822.

Pliocene of New Mexico.

57. **Bubo leptosteus** MARSH.

*Bubo leptosteus* MARSH, Am. Journ. Sci. II. 1871, 126.

Lower Tertiary of Wyoming.

58. **Aquila danana** MARSH.

*Aquila danana* MARSH, Am. Journ. Sci. II. 1871, 125.

Pliocene of Nebraska.

59. **Aquila pliogryps** SHUFELDT.

*Aquila pliogryps* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI.  
1892, 416, pl. xvii. fig. 33.

Pliocene of Oregon.

60. **Aquila sodalis** SHUFELDT.

*Aquila sodalis* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI. 1892,  
417.

Pliocene of Oregon.

## ORDER COCCYGES.

61. **Uintornis lucaris** MARSH.

*Uintornis lucaris* MARSH, Am. Journ. Sci. IV. 1872, 259.

Lower Tertiary formation of Wyoming.

## ORDER PASSERES.

62. **Corvus annectens** SHUFELDT.

*Corvus annectens* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI.  
1892, 419, pl. xv. fig. 14-16.

Pliocene of Oregon.

63. **Scolecophagus affinis** SHUFELDT.

*Scolecophagus affinis* SHUFELDT, Journ. Ac. Nat. Sci. Phila. XI.  
1892, 418, pl. xv. fig. 10.

Pliocene of Oregon.

64. **Palæospiza bella** ALLEN.

*Palæospiza bella* ALLEN, Bull. U. S. Geol. Surv. Terr. IV. No. 2,  
1878, 443, pl. i. figs. 1, 2.

Insect-bearing shales of Florissant, Colorado.





## INDEX.

---

- ACANTHIS, 217.  
     brewsterii, 330.  
     hornemannii, 217.  
     hornemannii exilipes, 217.  
     linaria, 217.  
     linaria holbœllii, 217.  
     linaria rostrata, 218.  
 Accipiter, 128.  
     atricapillus, 129.  
     atricapillus striatulus, 129.  
     cooperi, 128.  
     velox, 128.  
 Accipitrinæ, 126.  
 Actitis, 96.  
     macularia, 97.  
 Actochelidon, 23.  
 Actodromas, 87.  
 Æchmophorus, 1.  
     clarkii, 325.  
     occidentalis, 1.  
 Ægialitis, 100.  
     dubia, 101.  
     hiaticula, 101.  
     meloda, 101.  
     meloda circumcincta, 101.  
     mongola, 102.  
     montana, 103.  
     nivosa, 102.  
     semipalmata, 100.  
     vocifera, 100.  
     wilsonia, 102.  
 Aëronautæ, 172.  
     melanoleucus, 172.  
 Æsalon, 138.  
 Æstrelata, 34.  
     fisheri, 34.  
     hasitata, 34.  
     scalaris, 34.  
 Agelaius, 204.  
     gubernator, 205.  
     phœniceus, 204.  
     phœniceus bryanti, 205.  
     phœniceus sonoriensis, 205.  
     tricolor, 205.  
 Aix, 52.  
 Aix sponsa, 52.  
 Ajaja, 66.  
     ajaja, 67.  
 Alauda, 191.  
     arvensis, 191.  
 Alaudidæ, 191.  
 Albatross, Black-footed, 28.  
     Short-tailed, 28.  
     Sooty, 29.  
     Wandering, 326.  
     Yellow-nosed, 28.  
 Alca, 12.  
     torda, 12.  
 Alcedinidæ, 156.  
 Alcidiæ, 5, 325.  
 Alcinæ, 11.  
 Alcyones, 156.  
 Aletornis bellus, 343.  
     gracilis, 342.  
     nobilis, 342.  
     pernix, 342.  
     venustus, 342.  
 Alle, 13.  
     alle, 13.  
 Allinæ, 13.  
 Amazilia, 177.  
     cerviniventris, 177.  
     fuscicaudata, 177.  
 Ammodramus, 223, 227.  
     bairdii, 226.  
     beldingi, 225.  
     caudacutus, 227.  
     caudacutus nelsoni, 227.  
     caudacutus subvirgatus, 228.  
     henslowii, 226.  
     henslowii occidentalis, 227.  
     leconteii, 227.  
     maritimus, 228.  
     maritimus peninsulæ, 228.  
     maritimus sennetti, 228.  
     nigrescens, 229.  
     princeps, 223.  
     rostratus, 225.

- Ammodramus rostratus guttatus*, 225.  
     *sandwichensis*, 224.  
     *sandwichensis alaudinus*, 224.  
     *sandwichensis bryanti*, 224.  
     *sandwichensis savanna*, 224.  
     *savannarum passerinus*, 226.  
     *savannarum perpallidus*, 226.  
*Ampelidæ*, 260.  
*Ampelinæ*, 260.  
*Ampelis*, 260.  
     *cedrorum*, 260.  
     *garrulus*, 260.  
*Amphispiza*, 237.  
     *belli*, 238.  
     *belli cinerea*, 238.  
     *belli nevadensis*, 238.  
     *bilineata*, 237.  
*Anas*, 48.  
     *americana*, 49.  
     *boschas*, 48.  
     *carolinensis*, 50.  
     *crecca*, 50.  
     *cyanoptera*, 51.  
     *discors*, 50.  
     *fulvigula*, 48.  
     *fulvigula maculosa*, 49.  
     *obscura*, 48.  
     *penelope*, 49.  
     *strepera*, 49.  
*Anatidæ*, 47.  
*Anatinæ*, 48.  
*Ancylocheilus*, 90.  
*Anhinga*, 41.  
     *anhinga*, 41.  
*Anhingidæ*, 41.  
*Ani*, 153.  
     *Groove-billed*, 153.  
*Anorthura*, 300.  
*Anous*, 27.  
     *stolidus*, 27.  
*Anser*, 61.  
     *albifrons*, 61.  
     *albifrons gambeli*, 61.  
     *condoni*, 340.  
*Anseres*, 47.  
*Anserinæ*, 60.  
*Anthus*, 289.  
     *cervinus*, 290.  
     *pensilvanicus*, 289.  
     *pratensis*, 290.  
     *spragueii*, 290.  
*Antrostomus*, 168.  
     *carolinensis*, 168.  
     *vociferus*, 168.  
     *vociferus macromystax*, 169.  
*Apatornis celer*, 335.  
*Aphelocoma*, 196.  
     *californica*, 197.  
         *californica hypoleuca*, 197.  
         *californica obscura*, 197.  
         *cyanotis*, 197.  
         *floridana*, 196.  
         *insularis*, 197.  
         *sieberii arizonæ*, 198.  
         *woodhousei*, 196.  
*Aphriza*, 103.  
     *virgata*, 103.  
*Aphrizidæ*, 103.  
*Aphrizinæ*, 103.  
*Aquila*, 135.  
     *chrysaetos*, 135.  
     *danana*, 344.  
     *pliogryps*, 344.  
     *sodalis*, 344.  
*Aramidæ*, 76.  
*Aramus*, 76.  
     *giganteus*, 76.  
*Archibuteo*, 134.  
     *ferrugineus*, 135.  
     *lagopus*, 134.  
     *lagopus sancti-johannis*, 134.  
*Arctonetta*, 56.  
     *fischeri*, 56.  
*Ardea*, 70.  
     *candidissima*, 72.  
     *cinerea*, 71.  
     *cærulea*, 73.  
     *egretta*, 71.  
     *herodias*, 71.  
     *occidentalis*, 71.  
     *paloccidentalis*, 341.  
     *pealei*, 328.  
     *rufescens*, 72.  
     *tricolor ruficollis*, 72.  
     *virescens*, 73.  
     *virescens frazari*, 73.  
     *wardi*, 71.  
     *wuerdemanni*, 327.  
*Ardeidæ*, 69, 327.  
*Ardeinæ*, 70.  
*Ardetta*, 70.  
     *exilis*, 70.  
     *neoxena*, 70.  
*Arenaria*, 103.  
     *interpres*, 103.  
     *melanocephala*, 104.  
*Arenariinæ*, 103.  
*Arquatella*, 87.  
*Asio*, 142.  
     *accipitrinus*, 142.  
     *wilsonianus*, 142.  
*Astur*, 128.  
*Asturina*, 134.  
     *plagiata*, 134.

- Asyndesmus, 165.  
 Auk, Great, 13.  
     Razor-billed, 12.  
 Auklet, Cassin's, 7.  
     Crested, 7.  
     Least, 8.  
     Paroquet, 7.  
     Rhinoceros, 6.  
     Whiskered, 8.  
 Auriparus, 312.  
     flaviceps, 313.  
 Avocet, American, 83.  
 Aythya, 52.  
     affinis, 53.  
     americana, 52.  
     collaris, 53.  
     marila nearctica, 53.  
     vallisneria, 53.  
  
 BALDPATE, 49.  
 Baptonis advenus, 335.  
 Barornis regens, 339.  
 Bartramia, 96.  
     longicauda, 96.  
 Basileuterus, 288.  
     belli, 288.  
     culicivorus, 288.  
 Basilinna, 178.  
     leucotis, 178.  
     xantusi, 178.  
 Becard, Xantus's, 179.  
 Bird, Man-of-War, 46.  
     Red-billed Tropic, 39.  
     Surf, 103.  
     Yellow-billed Tropic, 39.  
 Bittern, American, 70.  
     Cory's Least, 70.  
     Least, 70.  
 Blackbird, Bicolored, 205.  
     Brewer's, 209.  
     Red-winged, 204.  
     Rusty, 209.  
     Tricolored, 205.  
     Yellow-headed, 204.  
 Bluebird, 322.  
     Azure, 322.  
     Chestnut-backed, 323.  
     Mountain, 323.  
     San Pedro, 322.  
     Western, 322.  
 Bluethroat, Red-spotted, 321.  
 Bobolink, 202.  
 Bob-white, 106.  
     Florida, 106.  
     Masked, 107.  
     Texan, 107.  
  
 Bonasa, 112.  
     umbellus, 112.  
     umbellus sabini, 112.  
     umbellus togata, 112.  
     umbellus umbelloides, 112.  
 Booby, 40.  
     Blue-faced, 40.  
     Blue-footed, 40.  
     Brewster's, 40.  
     Red-footed, 40.  
 Botaurinae, 69.  
 Botaurus, 69.  
     lentiginosus, 70.  
 Brachyramphus, 9.  
     craveri, 10.  
     hypoleucus, 9.  
     kittlitzii, 9.  
     marmoratus, 9.  
 Brant, 63.  
     Black, 63.  
 Branta, 62.  
     bernica, 63.  
     canadensis, 62.  
     canadensis hutchinsii, 62.  
     canadensis minima, 63.  
     canadensis occidentalis, 62.  
     hypsibatus, 341.  
     leucopsis, 63.  
     nigricans, 63.  
     propinqua, 341.  
 Bubo, 147.  
     leptosteus, 344.  
     virginianus, 148.  
     virginianus arcticus, 148.  
     virginianus saturatus, 148.  
     virginianus subarcticus, 148.  
 Bubonidae, 142.  
 Budytes, 289.  
     flavus leucostratus, 289.  
 Buffle-head, 54.  
 Bullfinch, Cassin's, 212.  
 Bulweria, 34.  
     bulweri, 35.  
 Bunting, Beautiful, 252.  
     Indigo, 252.  
     Lark, 254.  
     Lazuli, 252.  
     Painted, 252.  
     Townsend's, 331.  
     Varied, 252.  
 Burrica, 213.  
 Bush-Tit, 311.  
     Californian, 311.  
     Grinda's, 312.  
     Lead-colored, 312.  
     Lloyd's, 312.  
     Santa Rita, 312.

- Buteo*, 130.  
     *abbreviatus*, 132.  
     *albicaudatus sennetti*, 132.  
     *borealis*, 130.  
     *borealis calurus*, 130.  
     *borealis harlani*, 131.  
     *borealis kriderii*, 130.  
     *borealis lucasanus*, 131.  
     *brachyurus*, 133.  
     *buteo*, 130.  
     *cooperi*, 329.  
     *latissimus*, 133.  
     *lineatus*, 131.  
     *lineatus alleni*, 131.  
     *lineatus elegans*, 132.  
     *swainsoni*, 132.  
*Buteola*, 133.  
*Butorides*, 73.  
*Buzzard*, European, 130.  
  
*CALAMOSPIZA*, 254.  
     *melanocorys*, 254.  
*Calcarius*, 221.  
     *lapponicus*, 221.  
     *ornatus*, 222.  
     *pictus*, 221.  
*Calidris*, 91.  
     *arenaria*, 91.  
*Callichelidon*, 259.  
     *cyaneoviridis*, 259.  
*Callipepla*, 108.  
     *californica*, 109.  
     *californica vallicola*, 109.  
     *gambeli*, 109.  
     *squamata*, 108.  
     *squamata castanogastris*, 108.  
*Caliothrus*, 203.  
     *robustus*, 204.  
*Calothorax*, 177.  
     *lucifer*, 177.  
*Calypte*, 175.  
     *anna*, 175.  
     *costæ*, 175.  
*Campephilus*, 157.  
     *principalis*, 158.  
*Camptolaimus*, 56.  
     *labradorius*, 56.  
*Canachites*, 111.  
*Canvas-back*, 53.  
*Caprimulgi*, 168.  
*Caprimulgidæ*, 168, 330.  
*Caracara*, Audubon's, 141.  
     *Guadalupe*, 141.  
*Cardellina*, 287.  
*Cardellina rubrifrons*, 287.  
*Cardinal*, 249.  
  
*Cardinal*, Arizona, 249.  
     *Gray-tailed*, 249.  
     *Saint Lucas*, 249.  
     *Texas*, 250.  
*Cardinalis*, 248.  
     *cardinalis*, 249.  
     *cardinalis canicaudus*, 249.  
     *cardinalis igneus*, 249.  
     *cardinalis superbus*, 249.  
*Carinatae*, 339.  
*Carpodacus*, 212.  
     *amplus*, 214.  
     *cassini*, 213.  
     *mexicanus frontalis*, 213.  
     *mexicanus ruberrimus*, 214.  
     *purpureus*, 213.  
     *purpureus californicus*, 213.  
*Catbird*, 292.  
*Catharista*, 125.  
     *atrata*, 125.  
*Cathartes*, 125.  
     *aura*, 125.  
     *burrovianus*, 329.  
*Cathartidæ*, 124, 329.  
*Catherpes*, 296.  
     *mexicanus*, 297.  
     *mexicanus conspersus*, 297.  
     *mexicanus punctulatus*, 297.  
*Centrocerus*, 117.  
     *urophasianus*, 117.  
*Centronyx*, 225.  
*Centurus*, 165.  
*Ceophlæus*, 164.  
     *pileatus*, 164.  
*Cepphi*, 3.  
*Cephus*, 10.  
     *carbo*, 326.  
     *columba*, 11.  
     *grylle*, 10.  
     *mandtii*, 10.  
     *motzfeldi*, 325.  
*Cerorhinca*, 6.  
     *monocerata*, 6.  
*Certhia*, 303.  
     *familiaris alticola*, 303.  
     *familiaris americana*, 303.  
     *familiaris occidentalis*, 304.  
*Certhiidae*, 303.  
*Ceryle*, 156.  
     *alcyon*, 156.  
     *americana septentrionalis*, 157.  
     *torquata*, 157.  
*Chachalaca*, 119.  
*Chætura*, 172.  
     *pelagica*, 172.  
     *vauxii*, 172.  
*Chæturinæ*, 171.

- Chamaea*, 310.  
     *fasciata*, 310.  
     *fasciata henshawi*, 311.  
*Chamaeinæ*, 310.  
*Charadriidæ*, 98.  
*Charadrius*, 99.  
     *apricarius*, 99.  
     *dominicus*, 99.  
     *dominicus fulvus*, 100.  
     *sheppardianus*, 343.  
     *squatarola*, 99.  
*Charitonetta*, 54.  
     *albeola*, 54.  
*Chat*, Long-tailed, 285.  
     Yellow-breasted, 285.  
*Chaulelasmus*, 49.  
*Chelidon*, 258.  
     *erythrogaster*, 258.  
*Chen*, 60.  
     *cærulescens*, 61.  
     *hyperborea*, 60.  
     *hyperborea nivalis*, 60.  
     *rossii*, 61.  
*Chickadee*, 307.  
     *California*, 310.  
     *Carolina*, 308.  
     *Chestnut-backed*, 310.  
     *Columbian*, 310.  
     *Hudsonian*, 309.  
     *Kowak*, 309.  
     *Long-tailed*, 308.  
     *Mexican*, 309.  
     *Mountain*, 309.  
     *Oregon*, 308.  
     *Plumbeous*, 308.  
     *Siberian*, 309.  
*Chloroceryle*, 157.  
*Chondestes*, 229.  
     *grammacus*, 229.  
     *grammacus strigatus*, 229.  
*Chordeiles*, 170.  
     *acutipennis texensis*, 171.  
     *virginianus*, 170.  
     *virginianus chapmani*, 171.  
     *virginianus henryi*, 170.  
     *virginianus sennetti*, 330.  
*Chuck-will's-widow*, 168.  
*Ciceronia*, 8.  
*Ciconiæ*, 68.  
*Ciconiidæ*, 68.  
*Ciconiina*, 69.  
*Cimolopteryx* *rarus*, 335.  
     *retusus*, 335.  
*Cinclidæ*, 291.  
*Cinclus*, 291.  
     *mexicanus*, 291.  
*Circus*, 127.  
     *Circus hudsonius*, 128.  
*Cistothorus*, 301.  
     *marianæ*, 303.  
     *palustris*, 302.  
     *palustris griseus*, 302.  
     *palustris paludicola*, 302.  
     *stellaris*, 301.  
*Clamatores*, 179.  
*Clangula*, 55.  
     *hyemalis*, 55.  
*Clivicola*, 259.  
     *riparia*, 259.  
*Coccothraustes*, 211.  
     *vespertinus*, 211.  
     *vespertinus montanus*, 211.  
*Coccyges*, 153.  
*Coccyginæ*, 153.  
*Coccyzus*, 154.  
     *americanus*, 154.  
     *americanus occidentalis*, 155.  
     *erythrophthalmus*, 155.  
     *minor*, 154.  
     *minor maynardi*, 154.  
*Cœligena*, 173.  
     *clemenciæ*, 174.  
*Cœreba*, 267.  
     *bahamensis*, 268.  
*Cœrebidæ*, 267.  
*Colaptes*, 166.  
     *auratus*, 166.  
     *cafer*, 167.  
     *cafer saturator*, 167.  
     *chrysoides*, 167.  
     *rufipileus*, 167.  
*Colinus*, 106.  
     *ridgwayi*, 107.  
     *virginianus*, 106.  
     *virginianus floridanus*, 106.  
     *virginianus texanus*, 107.  
*Columba*, 119.  
     *fasciata*, 119.  
     *fasciata vioscæ*, 120.  
     *flavirostris*, 120.  
     *leucocephala*, 120.  
*Columbæ*, 119.  
*Columbidæ*, 119.  
*Columbigallina*, 122.  
     *passerina pallescens*, 123.  
     *passerina terrestris*, 122.  
*Colymbus*, 1.  
     *auritus*, 2.  
     *dominicus*, 2.  
     *holbœlli*, 1.  
*Colymbus nigricollis californicus*, 2.  
*Compsohalieus*, 43.  
*Compsothlypis*, 273.  
     *americana*, 273.

- Compsothlypis nigrilora*, 273.  
*Coniornis altus*, 336.  
*Contopus*, 185.  
     *borealis*, 185.  
     *pertinax*, 186.  
     *richardsonii*, 186.  
     *richardsonii peninsulæ*, 187.  
     *virens*, 186.  
*Conurus*, 152.  
     *carolinensis*, 152.  
*Coot*, American, 81.  
     European, 81.  
*Cormorant*, 42.  
     Baird's, 44.  
     Brandt's, 44.  
     Double-crested, 42.  
     Farallone, 43.  
     Florida, 42.  
     Mexican, 43.  
     Pallas's, 327.  
     Pelagic, 44.  
     Red-faced, 45.  
     Violet-green, 44.  
     White-crested, 43.  
*Corvidæ*, 194.  
*Corvinæ*, 199.  
*Corvus*, 199.  
     *americanus*, 200.  
     *americanus floridanus*, 200.  
     *caurinus*, 201.  
     *corax principalis*, 200.  
     *corax sinuatus*, 200.  
     *cryptoleucus*, 200.  
     *ossifragus*, 201.  
*Cotingidæ*, 179.  
*Coturnicops*, 79.  
*Coturniculus*, 226.  
*Cowbird*, 203.  
     Dwarf, 203.  
     Red-eyed, 204.  
*Cracidæ*, 119.  
*Crake*, Corn, 80.  
     Spotted, 78.  
*Crane*, Little Brown, 75.  
     Sandhill, 75.  
     Whooping, 75.  
*Creagrus*, 326.  
     *furcatus*, 326.  
*Creciscus*, 79.  
*Creepers*, Bahama Honey, 268.  
     Brown, 303.  
     California, 304.  
     Mexican, 303.  
     Rocky Mountain, 303.  
*Crex*, 80.  
     *crex*, 80.  
*Crossbill*, American, 214.  
     Mexican, 215.  
*Crossbill*, White-winged, 215.  
*Crotophaga*, 153.  
     *ani*, 153.  
     *sulcirostris*, 153.  
*Crotophaginae*, 153.  
*Crow*, American, 200.  
     Fish, 201.  
     Florida, 200.  
     Northwest, 201.  
*Crymophilus*, 82.  
     *fulicarius*, 82.  
*Cuckoo*, Black-billed, 155.  
     California, 155.  
     Kamchatkan, 155.  
     Mangrove, 154.  
     Maynard's, 154.  
     Yellow-billed, 154.  
*Cuculi*, 153.  
*Cuculidæ*, 153.  
*Cuculus*, 155.  
     *canorus telephonus*, 155.  
*Curlew*, Bristle-thighed, 98.  
     Eskimo, 97.  
     European, 328.  
     Hudsonian, 97.  
     Long-billed, 97.  
*Cyanecula*, 321.  
     *suecica*, 321.  
*Cyanocephalus*, 201.  
     *cyanocephalus*, 202.  
*Cyanocitta*, 194.  
     *cristata*, 194.  
     *cristata florincola*, 195.  
     *stelleri*, 195.  
     *stelleri annectens*, 196.  
     *stelleri frontalis*, 195.  
     *stelleri macrolopha*, 195.  
*Cyclorhynchus*, 7.  
     *psittaculus*, 7.  
*Cygninæ*, 65.  
*Cygnus paloregonus*, 341.  
*Cymodroma*, 38.  
     *grallaria*, 38.  
*Cypseli*, 171.  
*Cypseloides*, 171.  
     *niger*, 171.  
*Cyrtonyx*, 109.  
     *montezumæ*, 110.  
*Cyrtopelicanus*, 45.  
  
*DAFILA*, 51.  
     *acuta*, 51.  
*Daption*, 35.  
     *capensis*, 35.  
*Dendragapus*, 110.  
     *canadensis*, 111.  
     *franklinii*, 111.



- Dendragapus obscurus*, 110.  
     *obscurus fuliginosus*, 110.  
     *obscurus richardsonii*, 111.  
*Dendrocygna*, 64.  
     *autumnalis*, 64.  
     *fulva*, 64.  
*Dendroica*, 273, 274.  
     *æstiva*, 274.  
     *æstiva sonorana*, 274.  
     *auduboni*, 275.  
     *blackburniæ*, 277.  
     *bryanti castaneiceps*, 275.  
     *cærulea*, 276.  
     *cærulescens*, 275.  
     *carbonata*, 332.  
     *castanea*, 277.  
     *chrysoparia*, 278.  
     *coronata*, 275.  
     *discolor*, 281.  
     *dominica*, 277.  
     *dominica albilora*, 278.  
     *graciæ*, 278.  
     *kirtlandi*, 280.  
     *maculosa*, 276.  
     *montana*, 333.  
     *nigrescens*, 278.  
     *occidentalis*, 279.  
     *olivacea*, 274.  
     *palmarum*, 280.  
     *palmarum hypochrysea*, 280.  
     *pensylvanica*, 276.  
     *striata*, 277.  
     *tigrina*, 273.  
     *townsendi*, 279.  
     *vigorsii*, 280.  
     *virens*, 279.  
*Dichromanassa*, 72.  
*Dickcissel*, 254.  
*Diomedea*, 28.  
     *albatrus*, 28.  
     *exulans*, 326.  
     *nigripes*, 28.  
*Diomedeidæ*, 28, 326.  
*Dipper*, American, 291.  
*Dolichonyx*, 202.  
     *oryzivorus*, 202.  
*Dove*, Ground, 122.  
     *Inca*, 123.  
     *Mexican Ground*, 123.  
     *Mourning*, 121.  
     *White-fronted*, 122.  
     *White-winged*, 122.  
     *Zenaida*, 121.  
*Dovekie*, 13.  
*Dowitcher*, 85.  
     *Long-billed*, 86.  
*Dryobates*, 158.  
*Dryobates arizonæ*, 161.  
     *borealis*, 160.  
     *nuttallii*, 161.  
     *pubescens*, 159.  
     *pubescens gairdnerii*, 159.  
     *pubescens oreæcus*, 160.  
     *scalaris bairdi*, 160.  
     *scalaris lucasanus*, 160.  
     *villosus*, 158.  
     *villosus audubonii*, 158.  
     *villosus harrisii*, 159.  
     *villosus hyloscopus*, 159.  
     *villosus leucomelas*, 158.  
*Duck*, American Scaup, 53.  
     *Black*, 48.  
     *Florida*, 48.  
     *Harlequin*, 55.  
     *Labrador*, 56.  
     *Lesser Scaup*, 53.  
     *Masked*, 60.  
     *Mottled*, 49.  
     *Ring-necked*, 53.  
     *Ruddy*, 59.  
     *Rufous-crested*, 52.  
     *Steller's*, 55.  
     *Wood*, 52.  
*Dunlin*, 89.  
*Dysporus*, 41.  
*Dytes*, 2.  
  
*EAGLE*, Bald, 136.  
     *Golden*, 135.  
     *Gray Sea*, 136.  
     *Harpy*, 135.  
*Ectopistes*, 120.  
     *migratorius*, 120.  
*Egret*, American, 71.  
     *Peale's*, 328.  
     *Reddish*, 72.  
*Eider*, American, 57.  
     *King*, 58.  
     *Northern*, 57.  
     *Pacific*, 57.  
     *Spectacled*, 56.  
*Elanoides*, 126.  
     *forficatus*, 126.  
*Elanus*, 126.  
     *leucurus*, 126.  
*Embernagra*, 245.  
     *rufivirgata*, 245.  
*Empidonax*, 187.  
     *cineritius*, 187.  
     *difficilis*, 187.  
     *flaviventris*, 187.  
     *fulvifrons*, 189.  
     *fulvifrons pygmæus*, 189.

- Empidonax griseus*, 189.  
     *hammondi*, 189.  
     *minimus*, 188.  
     *traillii*, 188.  
     *traillii alnorum*, 188.  
     *virescens*, 188.  
     *wrightii*, 189.  
*Eniconetta*, 56.  
     *stelleri*, 56.  
*Ereunetes*, 90.  
     *occidentalis*, 91.  
     *pusillus*, 90.  
*Erionetta*, 57.  
*Ergaticus*, 287.  
     *ruber*, 288.  
*Erismatura*, 59.  
     *rubida*, 59.  
*Euethia*, 253.  
     *bicolor*, 253.  
     *canora*, 253.  
*Eugenes*, 173.  
     *fulgens*, 173.  
*Euphonia*, 254.  
     *elegantissima*, 255.  
*Euphonia*, Blue-headed, 255.  
*Eurynorhynchus*, 90.  
     *pygmæus*, 90.  
  
**FALCO**, 136.  
     *columbarius*, 138.  
     *columbarius suckleyi*, 139.  
     *dominicensis*, 140.  
     *fusco-cærulescens*, 139.  
     *islandus*, 136.  
     *mexicanus*, 137.  
     *peregrinus anatum*, 138.  
     *peregrinus pealei*, 138.  
     *regulus*, 139.  
     *richardsonii*, 139.  
     *rusticolus*, 137.  
     *rusticolus gyrfalco*, 137.  
     *rusticolus obsoletus*, 137.  
     *sparverius*, 140.  
     *sparverius deserticolus*, 140.  
     *sparverius peninsularis*, 140.  
     *tinnunculus*, 140.  
*Falcon*, Aplomado, 139.  
     *Peale's*, 138.  
     *Prairie*, 137.  
*Falcones*, 126.  
*Falconidæ*, 126, 329.  
*Falconinæ*, 136.  
*Finch*, California Purple, 213.  
     *Cassin's Purple*, 213.  
     *Guadalupe House*, 214.  
     *House*, 213.  
     *Finch*, Purple, 213.  
         *St. Lucas House*, 214.  
*Flamingo*, American, 66.  
*Flicker*, 166.  
     *Gilded*, 167.  
     *Guadalupe*, 167.  
     *Northwestern*, 167.  
     *Red-shafted*, 167.  
*Florida*, 73.  
*Flycatcher*, Alder, 188.  
     *Arizona Crested*, 183.  
     *Ash-throated*, 183.  
     *Beardless*, 190.  
     *Buff-breasted*, 189.  
     *Coues's*, 186.  
     *Crested*, 183.  
     *Derby*, 182.  
     *Fork-tailed*, 179.  
     *Fulvous*, 189.  
     *Giraud's*, 182.  
     *Gray*, 189.  
     *Green-crested*, 188.  
     *Hammond's*, 189.  
     *Lawrence's*, 184.  
     *Least*, 188.  
     *Mexican Crested*, 183.  
     *Nutting's*, 183.  
     *Olivaceous*, 184.  
     *Olive-sided*, 185.  
     *Ridgway's*, 190.  
     *St. Lucas*, 187.  
     *Scissor-tailed*, 180.  
     *Sulphur-bellied*, 182.  
     *Traill's*, 188.  
     *Vermilion*, 190.  
     *Western*, 187.  
     *Wright's*, 189.  
     *Yellow-bellied*, 187.  
*Fratercula*, 5.  
     *arctica*, 5.  
     *arctica glacialis*, 5.  
     *corniculata*, 6.  
*Fraterculinæ*, 5.  
*Fregata*, 46.  
     *aquila*, 46.  
*Fregatidæ*, 46.  
*Fringillidæ*, 211.  
*Fulica*, 81.  
     *americana*, 81.  
     *atra*, 81.  
     *minor*, 342.  
*Fulicinæ*, 81.  
*Fuligula*, 53.  
*Fulmar*, 30.  
     *Giant*, 29.  
     *Lesser*, 30.  
     *Pacific*, 30.

- Fulmar, Rodgers's, 30.  
     Slender-billed, 31.  
 Fulmarus, 30.  
     glacialis, 30.  
     glacialis glupischa, 30.  
     glacialis minor, 30.  
     glacialis rodgersii, 30.  
     glacialoides, 31.  
  
 GADWALL, 49.  
 Galeoscoptes, 291.  
     carolinensis, 291.  
 Gallinæ, 106.  
 Gallinago, 85.  
     delicata, 85.  
     gallinago, 85.  
 Gallinula, 81.  
     galeata, 81.  
 Gallinule, Florida, 81.  
     Purple, 80.  
 Gallinulinæ, 80.  
 Gannet, 41.  
 Garrulinæ, 194.  
 Garzetta, 72.  
 Gastornis giganteus, 338.  
 Gavia, 15.  
     alba, 15.  
 Gelocheidon, 22.  
     nilotica, 22.  
 Geococcyx, 153.  
     californianus, 154.  
 Geothlypis, 282, 283.  
     agilis, 282.  
     beldingi, 284.  
     formosa, 282.  
     macgillivrayi, 283.  
     philadelphia, 283.  
     trichas, 283.  
     trichas ignota, 284.  
     trichas occidentalis, 284.  
     poliocephala ralphi, 284.  
 Geotrygon, 123.  
     martinica, 123.  
     montana, 124.  
 Glaucidium, 150.  
     gnoma, 150.  
     gnoma californicum, 151.  
     hoskinsii, 151.  
     phalænoides, 151.  
 Glaucionetta, 54.  
     clangula americana, 54.  
     islandica, 54.  
 Glottis, 93.  
 Gnatcatcher, Black-tailed, 315.  
     Blue-gray, 315.  
     Plumbeous, 315.  
     Gnatcatcher, Western, 315.  
 Godwit, Black-tailed, 92.  
     Hudsonian, 92.  
     Marbled, 91.  
     Pacific, 92.  
 Golden-eye, American, 54.  
     Barrow's, 54.  
 Goldfinch, American, 218.  
     Arizona, 219.  
     Arkansas, 218.  
     Black-headed, 219.  
     Lawrence's, 219.  
     Mexican, 219.  
     Western, 218.  
 Goose, American White-fronted, 61.  
     Barnacle, 63.  
     Blue, 61.  
     Cackling, 63.  
     Canada, 62.  
     Emperor, 64.  
     Greater Snow, 60.  
     Hutchins's, 62.  
     Lesser Snow, 60.  
     Ross's Snow, 61.  
     White-cheeked, 62.  
     White-fronted, 61.  
 Goshawk, American, 129.  
     Mexican, 134.  
     Western, 129.  
 Grackle, Boat-tailed, 211.  
     Bronzed, 210.  
     Florida, 210.  
     Great-tailed, 210.  
     Purple, 210.  
 Graculavus pumilus, 336.  
     velox, 336.  
 Grassquit, 253.  
     Melodius, 253.  
 Grebe, American Eared, 2.  
     Clark's, 325.  
     Holboell's, 1.  
     Horned, 2.  
     Pied-billed, 3.  
     St. Domingo, 2.  
     Western, 1.  
 Green-shank, 93.  
 Grosbeak, Black-headed, 251.  
     Blue, 251.  
     Evening, 211.  
     Pine, 212.  
     Rose-breasted, 251.  
     Western Blue, 250.  
     Western Evening, 211.  
 Grouse, Canada, 111.  
     Canadian Ruffed, 112.  
     Columbian Sharp-tailed, 116.  
     Dusky, 110.

- Grouse, Franklin's, 111.  
 Gray Ruffed, 112.  
 Oregon Ruffed, 112.  
 Prairie Sharp-tailed, 117.  
 Richardson's, 111.  
 Ruffed, 112.  
 Sage, 117.  
 Sharp-tailed, 116.  
 Sooty, 110.
- Grues, 75.
- Gruidæ, 75.
- Grus, 75.  
 americana, 75.  
 canadensis, 75.  
 haydeni, 342.  
 mexicana, 75.  
 proavus, 342.
- Guara, 67.  
 alba, 67.  
 rubra, 67.
- Guillemot, Black, 10.  
 Black-winged, 325.  
 Mandt's, 10.  
 Pigeon, 11.  
 Sooty, 326.
- Guiraca, 251.  
 cærulea, 251.  
 cærulea euryncha, 251.
- Gull, American Herring, 19.  
 Bonaparte's, 21.  
 California, 19.  
 Franklin's, 20.  
 Glaucous, 16.  
 Glaucous-winged, 17.  
 Great Black-backed, 18.  
 Herring, 18.  
 Heermann's, 20.  
 Iceland, 17.  
 Ivory, 15.  
 Kumlien's, 17.  
 Laughing, 20.  
 Little, 21.  
 Mew, 20.  
 Nelson's, 17.  
 Pallas's, 89.  
 Point Barrow, 16.  
 Ring-billed, 19.  
 Ross's, 21.  
 Sabine's, 22.  
 Short-billed, 20.  
 Siberian, 18.  
 Slaty-backed, 18.  
 Swallow-tailed, 326.
- Gull, Vega, 19.  
 Western, 18.
- Gypagus, 329.  
 papa, 329.
- Gyr Falcon, 137.  
 Black, 137.  
 Gray, 137.  
 White, 136.
- HABIA, 250.  
 ludoviciana, 250.  
 melanocephala, 251.
- Hæmatopodidæ, 104.
- Hæmatopus, 104.  
 bachmani, 105.  
 frazari, 105.  
 ostralegus, 104.  
 palliatus, 104.
- Haliplana, 26.
- Halocyptena, 35.  
 microsoma, 35.
- Halizætus, 136.  
 albicilla, 136.  
 leucocephalus, 136.
- Harporhynchus, 292, 294.  
 bendirei, 293.  
 cinereus, 294.  
 cinereus mearnsi, 294.  
 crissalis, 295.  
 curvirostris, 293.  
 curvirostris palmeri, 293.  
 lecontei, 294.  
 longirostris sennetti, 293.  
 redivivus, 294.  
 rufus, 293.
- Hawk, American Rough-legged, 134.  
 American Sparrow, 140.  
 Desert Sparrow, 140.  
 Broad-winged, 133.  
 Cooper's, 128.  
 Cuban Sparrow, 140.  
 Duck, 138.  
 Florida Red-shouldered, 131.  
 Harlan's, 131.  
 Harris's, 129.  
 Krider's, 130.  
 Marsh, 128.  
 Mexican Black, 133.  
 Pigeon, 138.  
 Red-bellied, 132.  
 Red-shouldered, 131.  
 Red-tailed, 130.  
 Rough-legged, 134.  
 Sennett's White-tailed, 132.  
 Sharp-shinned, 128.  
 Short-tailed, 133.  
 St. Lucas Sparrow, 140.  
 Swainson's, 132.  
 Zone-tailed, 132.
- Heleodytes, 295.

- Heleodytes brunneicapillus*, 295.  
     *brunneicapillus affinis*, 296.  
     *brunneicapillus bryanti*, 295.  
*Helinaia*, 269.  
     *swainsonii*, 269.  
*Helminthophila*, 269.  
     *bachmani*, 270.  
     *celata*, 272.  
     *celata lutescens*, 272.  
     *celata sordida*, 272.  
     *chrysoptera*, 270.  
     *cincinnatiensis*, 332.  
     *lawrencei*, 331.  
     *leucobronchialis*, 332.  
     *lucia*, 270.  
     *peregrina*, 272.  
     *pinus*, 270.  
     *rufigapilla*, 271.  
     *rufigapilla gutturalis*, 271.  
     *virginia*, 271.  
*Helmitherus*, 269.  
     *vermivorus*, 269.  
*Helodromas*, 93.  
*Hen, Attwater's Prairie*, 115.  
     *Heath*, 115.  
     *Lesser Prairie*, 116.  
     *Prairie*, 115.  
*Henhawk, Cooper's*, 329.  
*Herodias*, 71.  
*Herodii*, 69.  
*Herodiones*, 66.  
*Heron, Black-crowned Night*, 74.  
     *European Blue*, 71.  
     *Frazar's Green*, 73.  
     *Great Blue*, 71.  
     *Great White*, 71.  
     *Green*, 73.  
     *Little Blue*, 73.  
     *Louisiana*, 72.  
     *Snowy*, 72.  
     *Ward's*, 71.  
     *Würdemann's*, 327.  
     *Yellow-crowned Night*, 74.  
*Hesperiphona*, 211.  
*Hesperocichla*, 320.  
     *naevia*, 321.  
*Hesperornis crassipes*, 336.  
     *gracilis*, 336.  
     *regalis*, 336.  
*Heteractitis*, 95.  
     *incana*, 95.  
*Hierofalco*, 136.  
*Himantopus*, 83.  
     *mexicanus*, 84.  
*Hirundinidæ*, 256.  
*Histrionicus*, 55.  
     *histrionicus*, 55.  
*Hummingbird, Allen's*, 176.  
     *Anna's*, 175.  
     *Black-chinned*, 174.  
     *Blue-throated*, 174.  
     *Broad-billed*, 178.  
     *Broad-tailed*, 175.  
     *Buff-bellied*, 177.  
     *Calliope*, 176.  
     *Costa's*, 175.  
     *Flores's*, 175.  
     *Lucifer*, 177.  
     *Rieffer's*, 177.  
     *Rivoli*, 173.  
     *Ruby-throated*, 174.  
     *Rufous*, 176.  
     *Violet-throated*, 174.  
     *White-eared*, 178.  
     *Xantus's*, 178.  
*Hydranassa*, 72.  
*Hydrochelidon*, 26.  
     *leucoptera*, 27.  
     *nigra surinamensis*, 26.  
*Hylocichla*, 316.  
*Hylophilus*, 331.  
     *decurtatus*, 331.  
*Hylophilus, Short-winged*, 331.  
  
*IACHE*, 178.  
     *latirostris*, 178.  
*Ibides*, 66.  
*Ibididæ*, 67.  
*Ibis, Glossy*, 68.  
     *Scarlet*, 67.  
     *White*, 67.  
     *White-faced Glossy*, 68.  
     *Wood*, 69.  
*Ichthyornis agilis*, 336.  
     *anceps*, 337.  
     *dispar*, 337.  
     *lentus*, 337.  
     *tener*, 337.  
     *validus*, 337.  
     *victor*, 337.  
*Icteria*, 284.  
     *virens*, 285.  
     *virens longicauda*, 285.  
*Icteridæ*, 202.  
*Icterus*, 206.  
     *audubonii*, 207.  
     *bullocki*, 208.  
     *cucullatus*, 207.  
     *cucullatus nelsoni*, 208.  
     *galbula*, 208.  
     *icterus*, 207.  
     *parisorum*, 207.  
     *spurius*, 208.

- Ictinia, 127.  
     mississippiensis, 127.  
 Iornornis, 80.  
     martinica, 80.
- JABIRU, 69.  
 Jacana, 105.  
     spinosa, 105.  
 Jaçana, Mexican, 105.  
 Jacanidæ, 105.  
 Jaeger, Parasitic, 14.  
     Pomarine, 14.  
     Long-tailed, 15.
- Jay, Alaskan, 199.  
     Arizona, 198.  
     Belding's, 197.  
     Black-headed, 196.  
     Blue, 194.  
     Blue-eared, 197.  
     Blue-fronted, 195.  
     California, 197.  
     Canada, 198.  
     Florida, 196.  
     Florida Blue, 195.  
     Green, 198.  
     Labrador, 199.  
     Long-crested, 195.  
     Oregon, 199.  
     Piñon, 202.  
     Rocky Mountain, 199.  
     Santa Cruz, 197.  
     Steller's, 195.  
     Woodhouse's, 196.  
     Xantus's, 197.
- Junco, 234.  
     aikeni, 234.  
     annectens, 235.  
     bairdi, 237.  
     caniceps, 236.  
     phæonotus dorsalis, 236.  
     phæonotus palliatus, 236.  
     hyemalis, 234.  
     hyemalis carolinensis, 235.  
     hyemalis oregonus, 234.  
     hyemalis pinosus, 235.  
     hyemalis shufeldti, 235.  
     hyemalis thurberi, 235.  
     insularis, 237.  
     ridgwayi, 236.  
     townsendi, 237.
- Junco, Arizona, 236.  
     Baird's, 237.  
     Carolina, 235.  
     Gray-headed, 236.  
     Guadalupe, 237.  
     Oregon, 234.
- Junco, Pink-sided, 235.  
     Point Pinos, 235.  
     Red-backed, 236.  
     Ridgway's, 236.  
     Slate-colored, 234.  
     Shufeldt's, 236.  
     Thurber's, 235.  
     Townsend's, 237.  
     White-winged, 234.
- KESTREL, 140.  
 Killdeer, 100.  
 Kingbird, 180.  
     Arkansas, 181.  
     Cassin's, 181.  
     Couch's, 181.  
     Gray, 180.
- Kingfisher, Belted, 156.  
     Ringed, 157.  
     Texan, 157.
- Kingleet, Cuvier's, 333.  
     Dusky, 314.  
     Golden-crowned, 313.  
     Ruby-crowned, 314.  
     Western Golden-crowned, 314.
- Kite, Everglade, 127.  
     Mississippi, 127.  
     Swallow-tailed, 126.  
     White-tailed, 126.
- Kittiwake, 15.  
     Pacific, 16.  
     Red-legged, 16.
- Knot, 86.
- LAGOPUS, 113.  
     lagopus, 113.  
     lagopus alleni, 113.  
     leucurus, 114.  
     rupestris, 113.  
     rupestris atkhensis, 114.  
     rupestris nelsoni, 114.  
     rupestris reinhardi, 114.  
     welchi, 114.
- Laniidæ, 261.  
 Lanius, 261.  
     borealis, 261.  
     ludovicianus, 262.  
     ludovicianus gambeli, 262.  
     ludovicianus excubitorides, 262.
- Lanivireo, 297.
- Laopteryx priscus, 335.
- Laornis edwardsianus, 337.
- Lapwing, 98.
- Laridæ, 15, 326.
- Larinæ, 15.



- Lark, Desert Horned, 192.  
   Dusky Horned, 193.  
   Horned, 191.  
   Mexican Horned, 192.  
   Pallid Horned, 192.  
   Prairie Horned, 192.  
   Ruddy Horned, 193.  
   Scorched Horned, 193.  
   Sonoran Horned, 193.  
   Streaked Horned, 193.  
   Texan Horned, 192.
- Larus, 16.  
   affinis, 18.  
   argentatus, 18.  
   argentatus smithsonianus, 19.  
   atricilla, 20.  
   barrovianus, 16.  
   brachyrhynchus, 20.  
   cachinnans, 19.  
   californicus, 89.  
   canus, 20.  
   delawarensis, 19.  
   franklinii, 20.  
   glaucescens, 17.  
   glaucus, 16.  
   heermanni, 20.  
   kumlieni, 17.  
   leucopterus, 17.  
   marinus, 18.  
   minutus, 21.  
   nelsoni, 17.  
   occidentalis, 18.  
   oregonus, 339.  
   philadelphia, 21.  
   robustus, 339.  
   schistisagus, 18.  
   vegæ, 19.
- Leptopelicanus, 46.
- Leptotila, 121.  
   fulviventris brachyptera, 122.
- Leucosticte, 215.  
   atrata, 216.  
   australis, 216.  
   griseonucha, 215.  
   tephrocotis, 215.  
   tephrocotis littoralis, 216.
- Leucosticte, Aleutian, 215.  
   Black, 216.  
   Brown-capped, 216.  
   Gray-crowned, 215.  
   Hepburn's, 216.
- Limicolæ, 82.
- Limosa, 91.  
   fedoa, 91.  
   hæmastica, 92.  
   lapponica baueri, 92.  
   limosa, 92.
- Limpkin, 76.
- Linnet, Brewster's, 330.
- Longipennes, 13.
- Longspur, Chestnut-collared, 222.  
   Lapland, 221.  
   McCown's, 222.  
   Smith's, 221.
- Loon, 3.  
   Black-throated, 4.  
   Pacific, 4.  
   Red-throated, 4.  
   Yellow-billed, 4.
- Lophodytes, 47.  
   cucullatus, 48.
- Lophophanes, 306.
- Lophortyx, 109.
- Loxia, 214.  
   curvirostra minor, 214.  
   curvirostra stricklandi, 215.  
   leucoptera, 215.
- Lunda, 5.  
   cirrhata, 5.
- MACROCHIRES, 168.
- Macrorhamphus, 85.  
   griseus, 85.  
   scelopaceus, 86.
- Magpie, American, 194.  
   Yellow-billed, 194.
- Mallard, 48.
- Man-o'-War Bird, 46.
- Mareca, 49.
- Martin, Cuban, 257.  
   Purple, 257.  
   Western, 257.
- Meadowlark, 206.  
   Mexican, 206.  
   Western, 206.
- Megalestris, 13.  
   skua, 14.
- Megaquiscalus, 210.
- Megascops, 145.  
   asio, 145.  
   asio aikenii, 147.  
   asio bendirei, 146.  
   asio cineraceus, 146.  
   asio floridanus, 145.  
   asio kennicottii, 146.  
   asio macfarlanei, 147.  
   asio maxwelliæ, 146.  
   asio trichopsis, 145.  
   flammeola, 147.  
   flammeola idahoensis, 147.
- Melanerpes, 164.  
   aurifrons, 166.  
   carolinus, 166.

- Melanerpes erythrocephalus*, 164.  
   *formicivorus angustifrons*, 165.  
   *formicivorus bairdi*, 165.  
   *torquatus*, 165.  
   *uropygialis*, 166.  
*Melanitta*, 58.  
*Meleagrinae*, 117.  
*Meleagris*, 117.  
   *altus*, 344.  
   *antiquus*, 343.  
   *celer*, 344.  
   *gallopavo*, 118.  
   *gallopavo ellioti*, 118.  
   *gallopavo mexicana*, 118.  
   *gallopavo osceola*, 118.  
*Melopelia*, 122.  
   *leucoptera*, 122.  
*Melospiza*, 240.  
   *cinerea*, 243.  
   *fasciata*, 241.  
   *fasciata clementæ*, 243.  
   *fasciata fallax*, 241.  
   *fasciata graminea*, 242.  
   *fasciata guttata*, 242.  
   *fasciata heermanni*, 241.  
   *fasciata montana*, 241.  
   *fasciata rivularis*, 242.  
   *fasciata rufina*, 242.  
   *fasciata samuelis*, 241.  
   *georgiana*, 244.  
   *insignis*, 243.  
   *lincolni*, 243.  
   *lincolni striata*, 243.  
*Merganser*, 47.  
   *americanus*, 47.  
   *serrator*, 47.  
*Merganser, American*, 47.  
   Hooded, 48.  
   Red-breasted, 47.  
*Merginae*, 47.  
*Merlin*, 139.  
   Black, 139.  
   Richardson's, 139.  
*Merula*, 319.  
   *confinis*, 320.  
   *migratoria*, 320.  
   *migratoria propinqua*, 320.  
*Methriopterus*, 292.  
*Micropalama*, 86.  
   *himantopus*, 86.  
*Micropallas*, 151.  
   *whitneyi*, 151.  
*Micropodidae*, 171.  
*Micropodinae*, 172.  
*Milvulus*, 179.  
   *forficatus*, 180.  
   *tyrannus*, 179.  
*Miminae*, 291.  
*Mimus*, 291.  
   *polyglottos*, 292.  
*Mniotilta*, 268.  
   *varia*, 268.  
*Mniotiltidae*, 268, 331.  
*Mockingbird*, 292.  
*Molothrus*, 203.  
   *ater*, 203.  
   *ater obscurus*, 203.  
*Motacilla*, 288.  
   *alba*, 289.  
   *ocularis*, 289.  
*Motacillidae*, 288.  
*Murre*, 11.  
   Brünnich's, 12.  
   California, 11.  
   Pallas's, 12.  
*Murrelet, Ancient*, 8.  
   Craveri's, 10.  
   Kittlitz's, 9.  
   Marbled, 9.  
   Xantus's, 9.  
*Myadestes*, 316.  
   *townsendii*, 316.  
*Myadestinae*, 316.  
*Mycteria*, 69.  
   *americana*, 69.  
*Myiarchus*, 182.  
   *cinerascens*, 183.  
   *cinerascens nuttingi*, 183.  
   *crinitus*, 183.  
   *lawrenceii*, 184.  
   *lawrenceii olivascens*, 184.  
   *mexicanus*, 183.  
   *mexicanus magister*, 183.  
*Myiozetetes*, 182.  
   *texensis*, 182.  
*Myiodynastes*, 182.  
   *luteiventris*, 182.  
  
*NEOCORYS*, 290.  
*Netta*, 52.  
   *rufina*, 52.  
*Nettion*, 50.  
*Nighthawk*, 170.  
   Florida, 171.  
   Sennett's, 330.  
   Texan, 171.  
   Western, 170.  
*Noddy*, 27.  
*Nomonyx*, 60.  
   *dominicus*, 60.  
*Nucifraga*, 201.  
   *columbianus*, 201.  
*Numenius*, 97.

*Numenius arquatus*, 328.  
     *borealis*, 97.  
     *hudsonicus*, 97.  
     *longirostris*, 97.  
     *phæopus*, 98.  
     *tahitiensis*, 98.  
*Nutcracker*, Clarke's, 201.  
*Nuthatch*, Brown-headed, 305.  
     Florida White-breasted, 304.  
     Pygmy, 305.  
     Red-breasted, 305.  
     Slender-billed, 304.  
     White-breasted, 304.  
     White-naped, 305.  
*Nuttallornis*, 185.  
*Nyctala*, 144.  
     *acadica*, 145.  
     *tengmalmi richardsoni*, 144.  
*Nyctanassa*, 74.  
*Nyctea*, 149.  
     *nyctea*, 149.  
*Nycticorax*, 74.  
     *nycticorax nævius*, 74.  
     *violaceus*, 74.  
*Nyctidromus*, 170.  
     *albicollis merrilli*, 170.  
  
 OCEANITES, 37.  
     *oceanicus*, 37.  
*Oceanitinæ*, 37.  
*Oceanodroma*, 36.  
     *furcata*, 36.  
     *homochroa*, 37.  
     *hornbyi*, 327.  
     *leucorhoa*, 36.  
     *macrodactyla*, 37.  
     *melania*, 37.  
*Ochthodromus*, 102.  
*Odontoglossæ*, 66.  
*Oidemia*, 58.  
     *americana*, 58.  
     *deglandi*, 59.  
     *fusca*, 58.  
     *perspicillata*, 59.  
*Oid-squaw*, 55.  
*Olor*, 65.  
     *buccinator*, 65.  
     *columbianus*, 65.  
     *cygnus*, 65.  
*Oporornis*, 282.  
*Oreortyx*, 107.  
     *pictus*, 107.  
     *pictus confinis*, 108.  
     *pictus plumiferus*, 107.  
*Oriole*, Arizona Hooded, 208.  
     Audubon's, 207.

*Oriole*, Baltimore, 208.  
     Bullock's, 208.  
     Hooded, 207.  
     Orchard, 208.  
     Scott's, 207.  
*Ornithion*, 190.  
     *imberbe*, 190.  
     *imberbe ridgwayi*, 190.  
*Oroscoptes*, 291.  
     *montanus*, 291.  
*Ortalis*, 119.  
     *vetula maccali*, 119.  
*Oscines*, 238.  
*Osprey*, American, 141.  
*Ossifraga*, 29.  
     *gigantea*, 29.  
*Otocoris*, 191.  
     *alpestris*, 191.  
     *alpestris adusta*, 193.  
     *alpestris arenicola*, 192.  
     *alpestris chrysolæma*, 192.  
     *alpestris giraudi*, 192.  
     *alpestris leucolæma*, 192.  
     *alpestris merrilli*, 193.  
     *alpestris pallida*, 193.  
     *alpestris praticola*, 192.  
     *alpestris rubea*, 193.  
     *alpestris strigata*, 193.  
*Oven-bird*, 281.  
*Owl*, American Barn, 142.  
     Aiken's Screech, 147.  
     American Hawk, 149.  
     American Long-eared, 142.  
     Arctic Horned, 148.  
     Barred, 143.  
     Burrowing, 150.  
     California Pygmy, 151.  
     California Screech, 146.  
     Dusky Horned, 148.  
     Dwarf Screech, 147.  
     Elf, 151.  
     Ferruginous Pygmy, 151.  
     Flammulated Screech, 147.  
     Florida Barred, 143.  
     Florida Burrowing, 150.  
     Florida Screech, 145.  
     Great Gray, 144.  
     Great Horned, 148.  
     Hawk, 149.  
     Hoskins's Pygmy, 151.  
     Kennicott's Screech, 146.  
     Lapp, 144.  
     MacFarlane's Screech, 147.  
     Mexican Screech, 146.  
     Pygmy, 150.  
     Richardson's, 144.  
     Rocky Mountain Screech, 145.

- Owl, Saw-whet, 145.  
 Screech, 145.  
 Short-eared, 142.  
 Snowy, 149.  
 Spotted, 143.  
 Texas Screech, 145.  
 Western Horned, 148.
- Oxyechus, 100.
- Oyster-catcher, 104.  
 American, 104.  
 Black, 105.  
 Frazer's, 105.
- PALÆOBORUS umbrosus, 344.
- Palæospiza bella, 345.
- Palæotetrix gilli, 343.
- Palæotringa littoralis, 337.  
 vagans, 338.  
 vetus, 338.
- Paludicolæ, 75.
- Pandion, 141.  
 haliaëtus carolinensis, 141.
- Pandioninæ, 141.
- Parabuteo, 129.  
 uncinctus harrisi, 129.
- Parauque, Merrill's, 170.
- Paridæ, 304.
- Parinæ, 332.
- Paroquet, Carolina, 152.
- Parrot, Thick-billed, 330.
- Partridge, California, 109.  
 Chestnut-bellied Scaled, 108.  
 Gambel's, 109.  
 Massena, 110.  
 Mountain, 107.  
 Plumed, 107.  
 San Pedro, 108.  
 Scaled, 108.  
 Valley, 109.
- Parus, 306.  
 atricapillus, 307.  
 atricapillus occidentalis, 308.  
 atricapillus septentrionalis, 308.  
 atricristatus, 306.  
 bicolor, 306.  
 bicolor texensis, 306.  
 carolinensis, 308.  
 carolinensis agilis, 308.  
 cinctus obtectus, 309.  
 gambeli, 309.  
 hudsonicus, 309.  
 hudsonicus columbianus, 310.  
 hudsonicus stoneyi, 309.  
 inornatus, 306.  
 inornatus cineraceus, 307.  
 inornatus griseus, 307.
- Parus meridionalis, 309.  
 rufescens, 310.  
 rufescens neglectus, 310.  
 wollweberi, 307.
- Passerculus, 223.
- Passerella, 244.  
 iliaca, 244.  
 iliaca megarhyncha, 245.  
 iliaca schistacea, 245.  
 iliaca unalaschensis, 244.
- Passeres, 179.
- Passerina, 251.  
 amœna, 252.  
 ciris, 252.  
 cyanea, 252.  
 versicolor, 252.  
 versicolor pulchra, 252.
- Pavoncella, 95.  
 pugnax, 95.
- Pediocætes, 116.  
 lucasi, 343.  
 nanus, 343.  
 phasianellus, 116.  
 phasianellus campestris, 117.  
 phasianellus columbianus, 116.
- Pelagodroma, 38.  
 marina, 38.
- Pelecanidæ, 45.
- Pelecanus, 45.  
 californicus, 46.  
 erythrorhynchos, 45.  
 fuscus, 46.
- Pelican, American White, 45.  
 Brown, 46.  
 California Brown, 46.
- Pelidna, 89.
- Pelionetta, 59.
- Pendulinus, 207.
- Penelopes, 118.
- Penelopina, 118.
- Perdicinæ, 106.
- Perisoreus, 198.  
 canadensis, 198.  
 canadensis capitalis, 199.  
 canadensis fumifrons, 199.  
 canadensis nigricapillus, 199.  
 obscurus, 199.
- Perissoglossa, 273.
- Petrel, Ashy, 37.  
 Black, 37.  
 Black-capped, 34.  
 Bulwer's, 35.  
 Fisher's, 34.  
 Fork-tailed, 36.  
 Guadalupe, 37.  
 Hornby's, 327.  
 Leach's, 36.

- Petrel, Least, 35.  
     Pintado, 35.  
     Scaled, 34.  
     Stormy, 36.  
     White-bellied, 38.  
     White-faced, 38.  
     Wilson's, 37.  
 Petrochelidon, 257.  
     fulva, 258.  
     lunifrons, 257.  
 Peucaea, 238.  
     æstivalis, 238.  
     æstivalis bachmanii, 239.  
     arizonæ, 239.  
     carpalis, 240.  
     cassini, 239.  
     mexicana, 239.  
     ruficeps, 240.  
     ruficeps boucardi, 240.  
     ruficeps eremœca, 240.  
 Peucedramus, 274.  
 Pewee, Large-billed, 187.  
     Western Wood, 186.  
     Wood, 186.  
 Phaëthon, 39.  
     æthereus, 39.  
     flavirostris, 39.  
 Phaëthontidæ, 39.  
 Phainopepla, 261.  
     nitens, 261.  
 Phalacrocoracidæ, 42, 327.  
 Phalacrocorax, 42.  
     carbo, 42.  
     dilophus, 42.  
     dilophus albociliatus, 43.  
     dilophus cinctus, 43.  
     dilophus floridanus, 42.  
     idahensis, 340.  
     macropus, 340.  
     mexicanus, 43.  
     pelagicus, 44.  
     pelagicus resplendens, 44.  
     pelagicus robustus, 44.  
     penicillatus, 44.  
     perspicillatus, 327.  
     urile, 45.  
 Phalænoptilus, 169.  
     nuttallii, 169.  
     nuttallii californicus, 169.  
     nuttallii nitidus, 169.  
 Phalarope, Northern, 82.  
     Red, 82.  
     Wilson's, 83.  
 Phalaropodidæ, 82.  
 Phalaropus, 82.  
     lobatus, 82.  
     tricolor, 83.  
 Phalerinæ, 6.  
 Phaleris, 8.  
 Phasiani, 106.  
 Phasianidæ, 117.  
 Philacte, 64.  
     canagica, 64.  
 Philohela, 84.  
     minor, 84.  
 Phœbe, 184.  
     Black, 185.  
     Say's, 185.  
 Phœbetria, 29.  
     fuliginosa, 29.  
 Phœnicopteridæ, 66.  
 Phœnicopterus, 66.  
     copei, 341.  
     ruber, 66.  
 Phylloperseus, 313.  
     borealis, 313.  
 Pica, 194.  
     nuttalli, 194.  
     pica hudsonica, 194.  
 Pici, 157.  
 Picidæ, 157.  
 Picoides, 161.  
     americanus, 162.  
     americanus alascensis, 162.  
     americanus dorsalis, 162.  
     arcticus, 162.  
 Picicorvus, 201.  
 Pigeon, Band-tailed, 119.  
     Passenger, 120.  
     Red-billed, 120.  
     Viosca's, 119.  
     White-crowned, 120.  
 Pinicola, 212.  
     enucleator, 212.  
 Pintail, 51.  
 Pipilo, 245.  
     alberti, 248.  
     chlorurus, 247.  
     consobrinus, 247.  
     erythrophthalmus, 246.  
     erythrophthalmus alleni, 246.  
     fuscus albigula, 248.  
     fuscus crissalis, 248.  
     fuscus mesoleucus, 247.  
     fuscus senicula, 248.  
     maculatus arcticus, 246.  
     maculatus megalonyx, 246.  
     maculatus oregonus, 247.  
 Pipit, American, 289.  
     Meadow, 291.  
     Red-throated, 290.  
     Sprague's, 290.  
 Piranga, 255.  
     erythromelas, 255.

- Piranga hepatica*, 256.  
*ludoviciana*, 255.  
*rubra*, 256.  
*rubra cooperi*, 256.  
*rubriceps*, 255.  
*Pitangus*, 181.  
*derbianus*, 182.  
*Plataleidae*, 66.  
*Platypsaris*, 179.  
*albiventris*, 179.  
*Plautus*, 12.  
*impennis*, 13.  
*Plectrophenax*, 220.  
*hyperboreus*, 221.  
*nivalis*, 220.  
*nivalis townsendi*, 220.  
*Plegadis*, 68.  
*autumnalis*, 68.  
*guarauna*, 68.  
*Plover*, American Golden, 99.  
*Belted Piping*, 101.  
*Black-bellied*, 99.  
*Golden*, 99.  
*Little Ring*, 101.  
*Mongolian*, 102.  
*Mountain*, 103.  
*Pacific Golden*, 100.  
*Piping*, 101.  
*Ring*, 101.  
*Semipalmated*, 100.  
*Snowy*, 102.  
*Wilson's*, 102.  
*Podasocys*, 102.  
*Podiceps*, 2.  
*Podicipedes*, 1.  
*Podicipidae*, 1, 325.  
*Podilymbus*, 3.  
*podiceps*, 3.  
*Polioptila*, 314.  
*cærulea*, 315.  
*cærulea obscura*, 315.  
*californica*, 315.  
*plumbea*, 315.  
*Polioptilinæ*, 314.  
*Polyborus*, 141.  
*cheriway*, 141.  
*lutosus*, 141.  
*Poocætes*, 222.  
*gramineus*, 222.  
*gramineus affinis*, 223.  
*gramineus confinis*, 223.  
*Poor-will*, 169.  
*Dusky*, 169.  
*Frosted*, 169.  
*Porzana*, 78.  
*carolina*, 79.  
*coturniculus*, 80.  
*Porzana jamaicensis*, 79.  
*noveboracensis*, 79.  
*porzana*, 78.  
*Priocella*, 31.  
*Priofinus*, 33.  
*Procellaria*, 36.  
*pelagica*, 36.  
*Procellariidae*, 29, 327.  
*Procellariinae*, 29.  
*Progne*, 256.  
*cryptoleuca*, 257.  
*subis*, 257.  
*subis hesperia*, 257.  
*Protonotaria*, 268.  
*citrea*, 268.  
*Psaltiriparus*, 311.  
*lloydi*, 312.  
*minimus*, 311.  
*minimus californicus*, 311.  
*minimus grindæ*, 312.  
*plumbeus*, 312.  
*santaritæ*, 312.  
*Pseudogryphus*, 124.  
*californianus*, 125.  
*Psittaci*, 152.  
*Psittacidae*, 152, 330.  
*Ptarmigan*, Allen's, 113.  
*Nelson's*, 114.  
*Reinhardt's*, 114.  
*Rock*, 113.  
*Turner's*, 114.  
*Welch's*, 114.  
*White-tailed*, 114.  
*Willow*, 113.  
*Ptiliogonatinae*, 261.  
*Ptychoramphus*, 6.  
*aleuticus*, 7.  
*Puffin*, 5.  
*Horned*, 6.  
*Large-billed*, 5.  
*Tufted*, 5.  
*Puffinus*, 31.  
*auduboni*, 32.  
*borealis*, 31.  
*cinereus*, 33.  
*conradii*, 340.  
*creatopus*, 32.  
*gavia*, 32.  
*griseus*, 33.  
*kuhlii*, 327.  
*major*, 31.  
*puffinus*, 32.  
*stricklandi*, 32.  
*tenuirostris*, 33.  
*Pygopodes*, 1.  
*Pyrocephalus*, 190.  
*rubineus mexicanus*, 190.



- Pyrrhula, 212.  
     cassini, 212.  
 Pyrrhuloxia, 249.  
     sinuata, 250.  
     sinuata beckhami, 250.  
     sinuata peninsulæ, 250.  
 Pyrrhuloxia, Arizona, 250.  
     Saint Lucas, 250.
- QUAIL-DOVE, Blue-headed, 124.  
     Key West, 123.  
     Ruddy, 124.
- Querquedula, 50.
- Quiscalus, 209.  
     macrourus, 210.  
     major, 211.  
     quiscula, 210.  
     quiscula æneus, 210.  
     quiscula aglæus, 210.
- RAIL, Belding's, 77.  
     Black, 79.  
     California Clapper, 77.  
     Caribbean Clapper, 78.  
     Clapper, 77.  
     Farallone, 80.  
     Florida Clapper, 78.  
     King, 76.  
     Louisiana Clapper, 77.  
     Virginia, 78.  
     Yellow, 79.
- Ralli, 76.
- Rallidæ, 76.
- Rallinæ, 76.
- Rallus, 76.  
     beldingi, 77.  
     elegans, 76.  
     crepitans, 77.  
     crepitans saturatus, 77.  
     longirostris caribæus, 78.  
     obsoletus, 77.  
     scottii, 78.  
     virginianus, 78.
- Raptore, 124.
- Raven, American, 200.  
     Northern, 200.  
     White-necked, 200.
- Ratitæ, 338.
- Recurvirostra, 83.  
     americana, 83.
- Recurvirostridæ, 83.
- Redhead, 52.
- Redpoll, 217.  
     Greenland, 217.
- Redpoll, Greater, 218.  
     Hoary, 217.  
     Holbøll's, 217.
- Redstart, American, 286.  
     Painted, 287.  
     Red-bellied, 287.
- Red-tail, St. Lucas, 131.  
     Western, 130.
- Regulina, 313.
- Regulus, 313.  
     calendula, 314.  
     cuvieri, 333.  
     obscurus, 314.  
     satrapa, 313.  
     satrapa olivaceus, 314.
- Rhodostethia, 21.  
     rosea, 21.
- Rhynchodon, 138.
- Rhynchofalco, 139.
- Rhynchophanes, 222.  
     mccownii, 222.
- Rhynchopsitta, 330.  
     pachyrhynca, 330.
- Rissa, 15.  
     brevirostris, 16.  
     tridactyla, 15.  
     tridactyla pollicaris, 16.
- Road-runner, 154.
- Robin, American, 320.  
     St. Lucas, 320.  
     Western, 320.
- Rostrhamus, 127.  
     sociabilis, 127.
- Rough-leg, Ferruginous, 135.
- Ruff, 95.
- Rynchopidæ, 27.
- Rynchops, 27.  
     nigra, 27.
- SALPINCTES, 296.  
     guadeloupensis, 296.  
     obsoletus, 296.
- Sanderling, 91.
- Sandpiper, Aleutian, 87.  
     Baird's, 88.  
     Bartramian, 96.  
     Buff-breasted, 96.  
     Cooper's, 328.  
     Curlew, 90.  
     Green, 94.  
     Least, 89.  
     Pectoral, 88.  
     Pribilof, 87.  
     Purple, 87.  
     Red-backed, 89.  
     Semipalmated, 90.

- Sandpiper, Sharp-tailed, 88.  
     Solitary, 94.  
     Spoonbill, 90.  
     Spotted, 97.  
     Stilt, 86.  
     Western, 91.  
     Western Solitary, 94.  
     White-rumped, 88.  
 Sapsucker, Red-breasted, 163.  
     Red-naped, 163.  
     Williamson's, 163.  
     Yellow-bellied, 163.  
 Sarcorhamphi, 124.  
 Saxicola, 321.  
     œnanthe, 321.  
 Sayornis, 184.  
     nigricans, 185.  
     phœbe, 184.  
     saya, 185.  
 Scardafella, 123.  
     inca, 123.  
 Scolecophagus, 209.  
     affinis, 345.  
     carolinus, 209.  
     cyanocephalus, 209.  
 Scolopacidae, 84, 328.  
 Scolopax, 84.  
     rusticola, 84.  
 Scoter, American, 58.  
     Surf, 59.  
     Velvet, 58.  
     White-winged, 59.  
 Scotiaptex, 144.  
     cinerea, 144.  
     cinerea lapponica, 144.  
 Seed-eater, Sharpe's, 253.  
 Seiurus, 281.  
     aurocapillus, 281.  
     motacilla, 282.  
     noveboracensis, 281.  
     noveboracensis notabilis, 281.  
 Selasphorus, 175.  
     alleni, 176.  
     floresii, 175.  
     platycercus, 175.  
     rufus, 176.  
 Setophaga, 286.  
     miniata, 287.  
     picta, 287.  
     ruticilla, 286.  
 Shearwater, Audubon's, 32.  
     Black-tailed, 33.  
     Black-vented, 32.  
     Cinereous, 327.  
     Cory's, 31.  
     Dark-bodied, 33.  
     Greater, 31.  
 Shearwater, Manx, 32.  
     Pink-footed, 32.  
     Slender-billed, 33.  
     Sooty, 32.  
 Shoveller, 51.  
 Shrike, California, 262.  
     Loggerhead, 262.  
     Northern, 261.  
     White-rumped, 262.  
 Sialia, 322.  
     arctica, 323.  
     mexicana anabelæ, 322.  
     mexicana bairdi, 323.  
     mexicana occidentalis, 322.  
     sialis, 322.  
     sialis azurea, 322.  
 Simorhynchus, 7.  
     cristatellus, 7.  
     pusillus, 8.  
     pygmæus, 8.  
 Siskin, Pine, 220.  
 Sitta, 304.  
     canadensis, 305.  
     carolinensis, 304.  
     carolinensis aculeata, 304.  
     carolinensis atkinsi, 304.  
     pusilla, 305.  
     pygmæa, 305.  
     pygmæa leuconucha, 305.  
 Sittinæ, 304.  
 Skimmer, Black, 27.  
 Skua, 14.  
 Skylark, 191.  
 Snipe, European, 85.  
     Wilson's, 85.  
 Snowflake, 220.  
     McKay's, 221.  
     Pribilof, 220.  
 Solitaire, Townsend's, 316.  
 Somateria, 57.  
     dresseri, 57.  
     mollissima borealis, 57.  
     spectabilis, 58.  
     v-nigra, 57.  
 Sora, 79.  
 Sparrow, Acadian Sharp-tailed, 228.  
     Aleutian Song, 243.  
     Arizona, 239.  
     Bachman's, 239.  
     Baird's, 226.  
     Belding's Marsh, 225.  
     Bell's, 238.  
     Bischoff's Song, 242.  
     Black-chinned, 234.  
     Black-throated, 237.  
     Boucard's, 240.  
     Brewer's, 233.

- Sparrow, Brown's Song, 242.  
   Bryant's Marsh, 224.  
   Cassin's, 239.  
   Chipping, 232.  
   Clay-colored, 232.  
   Desert Song, 241.  
   Dusky Seaside, 229.  
   Field, 233.  
   Forbush's, 243.  
   Fox, 244.  
   Gambel's, 231.  
   Golden-crowned, 231.  
   Grasshopper, 226.  
   Gray Sage, 238.  
   Harris's, 230.  
   Heermann's, 241.  
   Henslow's, 226.  
   Intermediate, 230.  
   Ipswich, 223.  
   Large-billed, 225.  
   Lark, 229.  
   Leconte's, 227.  
   Lincoln's, 243.  
   Mexican, 239.  
   Mountain Song, 241.  
   Nelson's, 227.  
   Oregon Vesper, 223.  
   Pine-woods, 238.  
   Rock, 240.  
   Rufous-crowned, 240.  
   Rufous-winged, 240.  
   Rusty Song, 242.  
   Sage, 238.  
   St. Lucas, 225.  
   Samuels's Song, 241.  
   San Clemente Song, 243.  
   Sandwich, 224.  
   Santa Barbara Song, 242.  
   Savanna, 224.  
   Scott's Seaside, 228.  
   Seaside, 228.  
   Sharp-tailed, 227.  
   Slate-colored, 245.  
   Song, 241.  
   Sooty Song, 242.  
   Swamp, 244.  
   Texas, 245.  
   Texas Seaside, 228.  
   Thick-billed, 245.  
   Townsend's, 244.  
   Tree, 231.  
   Vesper, 222.  
   Western Chipping, 232.  
   Western Field, 233.  
   Western Grasshopper, 226.  
   Western Henslow's, 227.  
   Western Lark, 229.  
  
 Sparrow, Western Savanna, 224.  
   Western Tree, 232.  
   Western Vesper, 223.  
   White-crowned, 230.  
   White-throated, 231.  
   Worthen's, 233.  
 Spatula, 51.  
   clypeata, 51.  
 Speotyto, 150.  
   cuniculara floridana, 150.  
   cunicularia hypogæa, 150.  
 Sphyrapicus, 163.  
   ruber, 163.  
   thyroideus, 163.  
   varius, 163.  
   varius nuchalis, 163.  
 Spinus, 218.  
   lawrencei, 219.  
   notatus, 219.  
   pinus, 220.  
   psaltria, 218.  
   psaltria arizonæ, 219.  
   psaltria mexicanus, 219.  
   tristis, 218.  
   tristis pallidus, 218.  
 Spiza, 254.  
   americana, 254.  
   townsendi, 331.  
 Spizella, 231.  
   atrigularis, 234.  
   breweri, 233.  
   monticola, 231.  
   monticola ochracea, 232.  
   pallida, 232.  
   pusilla, 233.  
   pusilla arenacea, 233.  
   socialis, 232.  
   socialis arizonæ, 232.  
   wortheni, 233.  
 Spoonbill, Roseate, 67.  
 Sporophila, 253.  
   morelleti sharpei, 253.  
 Squatarola, 99.  
 Starling, 202.  
 Sturnenas, 124.  
   cyanocephala, 124.  
 Steganopodes, 39.  
 Steganopus, 83.  
 Stelgidopteryx, 260.  
   serripennis, 260.  
 Stellula, 176.  
   calliope, 176.  
 Stercorariidæ, 13.  
 Stercorarius, 14.  
   longicaudus, 15.  
   parasiticus, 14.  
   pomarinus, 14.

- Sterna*, 23, 24.  
     *aleutica*, 25.  
     *anæthetus*, 26.  
     *antillarum*, 26.  
     *dougalli*, 25.  
     *elegans*, 23.  
     *forsteri*, 24.  
     *fuliginosa*, 26.  
     *hirundo*, 24.  
     *maxima*, 23.  
     *paradisæa*, 25.  
     *sandvicensis acuflavida*, 24.  
     *trudeaui*, 24.  
     *tschegrava*, 23.  
*Sternina*, 22.  
*Sternula*, 25.  
*Stilt*, Black-necked, 84.  
*Stint*, Long-toed, 89.  
*Streptoceryle*, 156.  
*Striges*, 142.  
*Strigidæ*, 142.  
*Strix*, 142.  
     *pratincola*, 142.  
*Sturnella*, 206.  
     *magna*, 206.  
     *magna mexicana*, 206.  
     *magna neglecta*, 206.  
*Sturnidæ*, 202.  
*Sturnus*, 202.  
     *vulgaris*, 202.  
*Sula*, 39.  
     *bassana*, 41.  
     *brewsteri*, 40.  
     *cyanops*, 40.  
     *gossi*, 40.  
     *loxostyla*, 340.  
     *piscator*, 40.  
     *sula*, 40.  
*Sulidæ*, 39.  
*Surnia*, 149.  
     *ulula*, 149.  
     *ulula caparoch*, 149.  
*Swallow*, Bahaman, 259.  
     *Bank*, 259.  
     *Barn*, 258.  
     *Cliff*, 257.  
     *Cuban Cliff*, 258.  
     *Rough-winged*, 260.  
     *Tree*, 257.  
     *Violet-green*, 259.  
*Swan*, Trumpeter, 65.  
     *Whistling*, 65.  
     *Whooping*, 65.  
*Swift*, Black, 171.  
     *Chimney*, 172.  
     *Vaux's*, 172.  
     *White-throated*, 173.  
*Sylvania*, 285.  
     *canadensis*, 286.  
     *microcephala*, 333.  
     *mitrata*, 285.  
     *pusilla*, 285.  
     *pusilla pileolata*, 286.  
*Sylviidæ*, 313, 333.  
*Sylviinæ*, 313.  
*Symphemia*, 94.  
     *semipalmata*, 94.  
     *semipalmata inornata*, 95.  
*Synthliboramphus*, 8.  
     *antiquus*, 8.  
*Syrnium*, 143.  
     *nebulosum*, 143.  
     *nebulosum alleni*, 143.  
     *occidentale*, 143.  
  
TACHYICINETA, 258.  
     *bicolor*, 258.  
     *thalassina*, 259.  
*Tachytiorchis*, 132.  
*Tanager*, Cooper's, 256.  
     *Gray's*, 255.  
     *Hepatic*, 256.  
     *Louisiana*, 255.  
     *Scarlet*, 255.  
     *Summer*, 256.  
*Tanagridæ*, 254.  
*Tantalina*, 68.  
*Tantalus*, 68.  
     *loculator*, 69.  
*Tatler*, Wandering, 95.  
*Teal*, Blue-winged, 50.  
     *Cinnamon*, 51.  
     *European*, 50.  
     *Green-winged*, 50.  
*Telmatodytes*, 302.  
*Telmatornis affinis*, 338.  
     *priscus*, 338.  
*Tern*, Aleutian, 25.  
     *Arctic*, 25.  
     *Black*, 26.  
     *Bridled*, 26.  
     *Cabot's*, 24.  
     *Caspian*, 23.  
     *Common*, 24.  
     *Elegant*, 23.  
     *Forster's*, 24.  
     *Gull-billed*, 22.  
     *Least*, 26.  
     *Roseate*, 25.  
     *Royal*, 23.  
     *Sooty*, 26.  
     *Trudeau's*, 24.  
     *White-winged Black*, 27.

- Tetraonidæ, 106.  
 Tetraoninæ, 110.  
 Thalasseus, 23.  
 Thalassogeron, 28.  
     culminatus, 28.  
 Thrasaëtos, 135.  
     harpyia, 135.  
 Thrasher, Bendire's, 293.  
     Brown, 293.  
     California, 294.  
     Crissal, 295.  
     Curve-billed, 293.  
     Leconte's, 294.  
     Mearns's, 294.  
     Palmer's, 293.  
     Sage, 291.  
     St. Lucas, 294.  
     Sennett's, 293.  
 Thrush, Audubon's Hermit, 319.  
     Bicknell's, 317.  
     Dwarf Hermit, 318.  
     Gray-cheeked, 317.  
     Hermit, 319.  
     Olive-backed, 318.  
     Red-winged, 319.  
     Russet-backed, 318.  
     Varied, 321.  
     Willow, 317.  
     Wilson's, 317.  
     Wood, 316.  
 Thryomanes, 298.  
 Thryothorus, 297.  
     bewickii, 298.  
     bewickii bairdi, 299.  
     bewickii spilurus, 299.  
     brevicaudus, 299.  
     leucophrys, 299.  
     ludovicianus, 298.  
     ludovicianus lomitisensis, 298.  
     ludovicianus miamensis, 298.  
 Tinnunculus, 140.  
 Titmouse, Ashy, 307.  
     Black-crested, 306.  
     Bridled, 307.  
     Gray, 307.  
     Plain, 306.  
     Texas Tufted, 306.  
     Tufted, 306.  
 Totanus, 92.  
     flavipes, 93.  
     melanoleucus, 93.  
     nebularius, 93.  
     ochropus, 94.  
     solitarius, 94.  
     solitarius cinnamomeus, 94.  
 Towhee, 246.  
     Abert's, 248.  
     Towhee, Anthony's, 248.  
         Arctic, 246.  
         California, 248.  
         Cañon, 247.  
         Green-tailed, 247.  
         Guadalupe, 247.  
         Oregon, 247.  
         Saint Lucas, 248.  
         Spurred, 246.  
         White-eyed, 246.  
 Tree-duck, Black-bellied, 64.  
     Fulvous, 64.  
 Tringa, 86.  
     acuminata, 88.  
     alpina, 89.  
     alpina pacifica, 89.  
     bairdii, 88.  
     canutus, 86.  
     cooperi, 328.  
     couesi, 87.  
     damacensis, 89.  
     ferruginea, 90.  
     fuscicollis, 88.  
     maculata, 88.  
     maritima, 87.  
     minutilla, 89.  
     ptilocnemis, 87.  
 Trochili, 173.  
 Trochilidæ, 173.  
 Trochilus, 174.  
     alexandri, 174.  
     colubris, 174.  
     violajugulum, 174.  
 Troglodytes, 300.  
     aëdon, 300.  
     aëdon aztecus, 300.  
     aëdon parkmanii, 300.  
     alascensis, 301.  
     hiemalis, 301.  
     hiemalis pacificus, 301.  
 Troglodytidæ, 291.  
 Troglodytinæ, 295.  
 Trogon, 156.  
     ambiguus, 156.  
     Coppery-tailed, 156.  
 Trogones, 156.  
 Trogonidæ, 156.  
 Troupial, 207.  
 Tryngites, 96.  
     subruficollis, 96.  
 Tubinares, 28.  
 Turdidæ, 316.  
 Turdinæ, 316.  
 Turdus, 316.  
     aliciæ, 317.  
     aliciæ bicknelli, 317.  
     aonalaschkæ, 318.

- Turdus aonalaschkæ auduboni*, 319.  
*aonalaschkæ pallasi*, 319.  
*fuscescens*, 317.  
*fuscescens salicicolus*, 317.  
*iliacus*, 319.  
*mustelinus*, 316.  
*ustulatus*, 318.  
*ustulatus swainsoni*, 318.  
 Turkey, Florida Wild, 118.  
     Mexican, 118.  
     Rio Grande Wild, 118.  
     Wild, 118.  
 Turnstone, 103.  
     Black, 104.  
 Tympanuchus, 115.  
     americanus, 115.  
     americanus attwateri, 115.  
     cupido, 115.  
     pallidicinctus, 116.  
 Tyrannidæ, 179.  
 Tyrannus, 180.  
     dominicensis, 180.  
     melancholicus couchii, 181.  
     tyrannus, 180.  
     verticalis, 181.  
     vociferans, 181.  
  
 UINTORNIS lucaris, 345.  
 Uria, 11.  
     affinis, 339.  
     antiqua, 339.  
     lomvia, 12.  
     lomvia arra, 12.  
     troile, 11.  
     troile californica, 11.  
 Urile, 44.  
 Urinator, 3.  
     adamsii, 4.  
     arcticus, 4.  
     imber, 3.  
     lumme, 4.  
     pacificus, 4.  
 Urinatoridæ, 3.  
 Urubitinga, 133.  
     anthracina, 133.  
  
 VANELLUS, 98.  
     vanellus, 98.  
 Verdin, 313.  
 Vireo, 272, 265.  
     atricapillus, 265.  
     bellii, 267.  
     bellii pusillus, 267.  
     calidris barbatulus, 263.  
  
*Vireo flavifrons*, 264.  
*flavoviridis*, 263.  
*gilvus*, 264.  
*huttoni*, 266.  
*huttoni obscurus*, 266.  
*huttoni stephensi*, 266.  
*novaboracensis*, 266.  
*novaboracensis maynardi*, 266.  
*olivaceus*, 263.  
*philadelphicus*, 263.  
*solitarius*, 264.  
*solitarius alticola*, 265.  
*solitarius cassinii*, 264.  
*solitarius lucasanus*, 265.  
*solitarius plumbeus*, 265.  
*vicinior*, 267.  
 Vireo, Anthony's, 266.  
     Bell's, 267.  
     Black-capped, 265.  
     Black-whiskered, 263.  
     Blue-headed, 264.  
     Cassin's, 264.  
     Gray, 267.  
     Hutton's, 266.  
     Key West, 266.  
     Least, 267.  
     Mountain Solitary, 265.  
     Philadelphia, 263.  
     Plumbeous, 265.  
     Red-eyed, 263.  
     St. Lucas Solitary, 265.  
     Stephens's, 266.  
     Warbling, 264.  
     White-eyed, 266.  
     Yellow-green, 263.  
     Yellow-throated, 264.  
 Vireonidæ, 262, 331.  
 Vireosylva, 262.  
 Vulture, Black, 125.  
     Burroughs's Turkey, 329.  
     California, 125.  
     King, 329.  
     Turkey, 125.  
  
 WAGTAIL, Siberian Yellow, 289.  
     Swinhoe's, 289.  
     White, 289.  
 Warbler, Audubon's, 275.  
     Bachman's, 270.  
     Bay-breasted, 277.  
     Bell's, 288.  
     Black and White, 268.  
     Blackburnian, 277.  
     Black-poll, 277.  
     Black-throated Blue, 275.  
     Black-throated Gray, 278.



## Warbler, Black-throated Green, 279.

- Blue Mountain, 333.
- Blue-winged, 270.
- Brasher's, 288.
- Brewster's, 332.
- Calaveras, 271.
- Canadian, 286.
- Cape May, 273.
- Carbonated, 332.
- Cerulean, 276.
- Chestnut-sided, 276.
- Cincinnati, 332.
- Connecticut, 282.
- Dusky, 272.
- Golden-cheeked, 278.
- Golden-winged, 270.
- Grace's, 278.
- Hermit, 279.
- Hooded, 285.
- Kennicott's Willow, 313.
- Kentucky, 282.
- Kirtland's, 280.
- Lawrence's, 331.
- Lucy's, 270.
- Lutescent, 272.
- Macgillivray's, 283.
- Magnolia, 276.
- Mangrove, 275.
- Mourning, 283.
- Myrtle, 275.
- Nashville, 271.
- Olive, 274.
- Orange-crowned, 272.
- Palm, 280.
- Parula, 273.
- Pileolated, 286.
- Pine, 280.
- Prairie, 281.
- Prothonotary, 268.
- Red, 288.
- Red-faced, 287.
- Sennett's, 273.
- Small-headed, 333.
- Sonora Yellow, 274.
- Swainson's, 269.
- Sycamore, 278.
- Tennessee, 272.
- Townsend's, 279.
- Virginia's, 271.
- Wilson's, 285.
- Worm-eating, 269.
- Yellow, 274.
- Yellow Palm, 280.
- Yellow-throated, 277.

## Water-Thrush, 281.

- Grinnell's, 281.
- Louisiana, 282.

## Waxwing, Bohemian, 260.

- Cedar, 260.
- Wheatear, 321.
- Whimbrel, 98.
- Whip-poor-will, 168.
- Stephens's, 169.
- Widgeon, 49.
- Willet, 94.
- Western, 95.
- Woodcock, American, 84.
- European, 84.
- Woodpecker, Alaskan Three-toed, 162.
- Alpine Three-toed, 162.
- American Three-toed, 162.
- Arctic Three-toed, 162.
- Arizona, 161.
- Batchelder's, 160.
- Cabanis's, 159.
- Californian, 164.
- Downy, 159.
- Gairdner's, 159.
- Gila, 166.
- Golden-fronted, 166.
- Hairy, 158.
- Harris's, 159.
- Ivory-billed, 158.
- Lewis's, 164.
- Narrow-fronted, 164.
- Northern Hairy, 158.
- Nuttall's, 161.
- Pileated, 164.
- Red-bellied, 165.
- Red-cockaded, 160.
- Red-headed, 164.
- Saint Lucas, 160.
- Southern Hairy, 158.
- Texan, 160.
- White-headed, 161.
- Williamson's, 163.
- Wren, Alaskan, 301.
- Baird's, 299.
- Bewick's, 298.
- Bryant's Cactus, 295.
- Cactus, 295.
- Cañon, 297.
- Carolina, 298.
- Dotted Cañon, 295.
- Florida, 298.
- Guadalupe, 299.
- Guadalupe Rock, 296.
- House, 300.
- Lomita, 298.
- Long-billed Marsh, 302.
- Marian's Marsh, 303.
- Parkman's, 300.
- Rock, 296.
- St. Lucas Cactus, 296.

Wren, San Clemente, 299.  
     Short-billed Marsh, 301.  
     Tulé, 302.  
     Vigors's, 299.  
     Western House, 300.  
     Western Winter, 301.  
     White-throated, 297.  
     Winter, 301.  
     Worthington's Marsh, 302.  
 Wren-Tit, 310.  
     Pallid, 311.

XANTHOCEPHALUS, 204.  
     xanthocephalus, 204.

Xanthoura, 198.  
     luxuosa, 198.

Xema, 22.  
     sabinii, 22.

Xenopicus, 161.  
     albolarvatus, 161.

YELLOW-LEGS, 93.  
     Greater, 93.  
 Yellow-throat, Belding's, 284.  
     Florida, 284.  
     Maryland, 283.  
     Rio Grande, 284.  
     Western, 284.  
 Yphantes, 208.

ZENAIDA, 121.  
     zenaida, 121.

Zenaidura, 121.  
     macroura, 121.

Zonotrichia, 230.  
     albicollis, 231.  
     coronata, 231.  
     leucophrys, 230.  
     leucophrys gambeli, 231.  
     leucophrys intermedia, 230.  
     querula, 230.









SMITHSONIAN INSTITUTION LIBRARIES



3 9088 00559 0203